

Calcium Antagonistic Activity of *Biophytum petersianum* on Vascular Smooth Muscles of Wistar Rat

SIMÉNOU TITRIKOU, KWASHIE EKLUGADEGBEKEU, AKLESSO MOUZOU, KODJO AKLIKOKOU and MESSANVI GBEASSOR

For author affiliations, see end of text.

Received August 21, 2007; Revised December 22, 2007; Accepted January 15, 2008

This paper is available online at <http://ijpt.iums.ac.ir>

ABSTRACT

The whole plant of *Biophytum petersianum* was extracted with a mixture of water – alcohol (1:1) to evaluate its relaxant effect on aorta rings. In isolated Wistar rat tissue, the hydro-ethanolic extract (0.1, 0.25 and 0.5 mg/ml) non-competitively antagonized calcium chloride and high- K^+ -induced aorta contractions in a concentration-dependent manner. Moreover, the inhibition of noradrenaline-induced contractions in the presence of extract (1 mg/ml) suggests that the extract has an effect on mobilization of intracellular calcium. These results indicate that hypotensive effect of *Biophytum petersianum* may result from inhibition of calcium influx via both voltage- and receptor-operated calcium channels.

Keywords: *Biophytum petersianum*, Aorta, Calcium antagonistic activity, Hypertension

Biophytum petersianum Klotzsch (Oxalidaceae.) is an annual plant which is distributed widely in tropical areas in Africa, Madagascar and Asia. It has been used in traditional medicine for many purposes. In Mali, the leaves are used in wound healing and for its complement fixing activity [1]. It is reported that aqueous extract of the leaves has hypocholesterolemic [2] and hypoglycaemic [3] effects. It also showed insulinotropic activity in non-diabetic and alloxan-diabetic rabbits [4]. Amentoflavone extracted from the roots of *B. petersianum* showed a selective inhibition of cyclooxygenase (COX)-1/COX-2 [5]. In Togo, the whole plant is reported to have antihypertensive effect and previous studies with hydro-ethanolic extract on guinea-pig and Wistar rat confirmed this indication [6]. Hydro-ethanolic extract of the whole plant is also claimed to stimulate corticosterone and aldosterone secretion in rat without any desensitization phenomenon [7].

Calcium plays an important role in the pathophysiological process of hypertension through the increase of heart activity, mainly by activation of smooth muscle contraction under a variety of circumstances [8]. It's well documented that substances that inhibit calcium release in vascular smooth cells prevent vasoconstriction [9] and are beneficial in the treatment of hypertension.

The antihypertensive effect of *B. petersianum* may be explained by inhibition of calcium release into the inside of vascular smooth cells. The present study was designed to examine the possible effect of hydro-alcoholic extract of *B. petersianum* on intracellular mobilization and on extracellular calcium influx through L-Type channels of rat aortic rings in Ca^{2+} -free and high K^+ medium.

MATERIALS AND METHODS

Chemicals

Noradrenaline, Verapamil, Quercetin, Gallic Acid and Sodium carbonate were purchased from Sigma Chemical Co. (St. Louis, MO).

Plant material

Whole plant material was collected in September 2003 in the locality of Kévé (southern region of Togo). A voucher specimen was deposited in the herbarium under the number 1032 in the Department of Botanic, Faculty of Sciences, University of Lomé, Togo.

Animals

Wistar rats of both sex (180-230 g) provided by the Department of Animal Physiology were used in these




Fig. 1. Muscle tension produced by cumulative concentrations of calcium chloride in the absence and presence of hydro-alcoholic extract of *B. petersianum* in rat thoracic aorta. **A:** representative traces of the effect of the cumulative calcium chloride concentrations on rat isolated aorta in a high K^+ medium without calcium in the absence (a) and in the presence of the 0.1 mg/ml (b), 0.25 mg/ml (c) and 0.5 mg/ml (d) extract and 1 μ M verapamil (e). **B:** Effects of the plant extract on the contraction induced by calcium chloride in rat thoracic aorta. Each point represents the mean of 5 preparations. Vertical bars indicate the S.E.M. * $p < 0.001$, when compared with control value.

experiments. They were kept under standard environmental conditions with free access to food and water.

Preparation of extract

The powder of the whole plant (100 g) was macerated during 72 hours in water/ethanol (50% v/v) solution, at room temperature with occasional stirring. After filtration, the filtrate was evaporated in a vacuum rotavapor at 50°C (yield 12 %). The extract was dark brown and soluble in distilled water.

Total phenols determination

The phenolic content in the hydro-alcoholic extract of *Biophytum petersianum* was measured by Folin-Ciocalteu reagent using gallic acid as standard [10]. Briefly, we dissolved the plant extract in methanol/distilled water (v/v, 1:1) to achieve a concentration of 1mg/ml. About 500 μ l of this solution was mixed with

5 ml of Folin-Ciocalteu reagent (1:10 ml distilled water), and 4 ml of sodium carbonate (Na_2CO_3 , 1 M). The mixture was incubated at room temperature for 15 minutes and the absorbance was measured at 765 nm against a methanol blank using a spectrophotometer (Sequoia-Turner, Model 340). Total phenolic content was determined using a six point standard curve of Gallic acid (0–250 mg/L). All tests were carried out in triplicate. Data were converted into milligram gallic acid equivalents per gram of *Biophytum petersianum* extract (GAE/g of BPE).

Total flavonoids determination

Aluminium chloride colorimetric method described by Pourmorad et al. [10] was used to determine flavonoid content in plant extract. The extract was dissolved in methanol to achieve a final concentration of 10 mg/ml. About 500 μ l of this solution was mixed with 1.5ml of




Fig. 2. Effect of noradrenaline (1 μ M) on rat thoracic aorta in Ca^{2+} -free, high K^{+} (60 mM) solution in absence (control) and presence of hydro-alcoholic extract of *B. petersianum* (1 mg/ml). The extract was added 20 min prior to drug addition. **A:** representative traces of the opposing effect of extract on noradrenaline-induced contraction in a depolarized medium without calcium in the absence (a) and in presence (b) of 1 mg/ml extract **B:** Effects of the plant extract on the contraction induced by adrenaline in rat thoracic aorta. Each point represents the mean of 6 preparations. Vertical bars indicate the S.E.M., * $p < 0.001$ when compared with control value.

methanol, 100 μ l of aluminium chloride (0.1 g/ml), 100 μ l of sodium acetate (1 M) and 2.8 ml of distilled water. After 30 minutes incubation at room temperature, the absorbance was measured at 415 nm against methanol blank with a spectrophotometer. Quercetin was chosen as standard and six points standards curve (0 – 1000 μ g/ml) was used for the determination of total flavonoid content. All tests were carried out in replicate ($n = 4$). The data were converted into milligram quercetin equivalents per gram of *Biophytum petersianum* extract (QE/g of BPE).

Aortic preparation

The rats were sacrificed under ether anesthesia. Thoracic aorta was then removed and immediately placed in cold Krebs-Henseleit solution. Composition of the Krebs's solution in mM was as follows: NaCl 118, KCl 4.7, CaCl_2 2.5, KH_2PO_4 2.5, MgSO_4 1.2, NaHCO_3 25, glucose 11.0. The aorta was cleaned of adherent tissue and cut into 5 mm rings, taking care not to damage the endothelium. Each ring was fixed vertically between triangular stainless steel hooks (SA-068, UGO BASILE) under a resting tension of 2 g in 10 ml bath filled with the Krebs's solution and maintained at 37°C. The bath solution was continuously oxygenated and the rings were allowed to equilibrate for 90 minutes before the start of the experiments. Initially, all rings were contracted twice with KCl (60 mM) to confirm the viability

of vessels before performing the experimental protocols. Isometric tension change was measured with TDS105A Variable Force transducer (BIOPAC SYSTEM, MODEL MP100, HAVARD APPARATUS) coupled to a computer (Acq Knowledge III software).

Effects of *B. petersianum* on extracellular Ca^{2+} influx through voltage-sensitive channels

Aortic ring depolarized and contracted by Ca^{2+} was chosen as the model to investigate the effects of *B. petersianum*'s hydro-ethanolic extract on calcium influx through voltage-sensitive channels, as previously described by Hof et al. [9]. The rings were exposed to calcium-free high-potassium depolarizing solution (NaCl of Krebs's solution was replaced by 60 mM of KCl). Vessels were allowed to equilibrate for at least 30 minutes before drug addition and the solution was changed twice at 15-minute intervals. Then, cumulative concentration-response curves to CaCl_2 (0.1–8 mM) were determined in the absence (control) or in the presence of extract (0.1; 0.25 and 0.5 mg/ml) or verapamil (1 μ M), which was used as the reference drug.

Effects of *B. petersianum* on the release of intracellular calcium

The method of Dar et al. was used [11]. Briefly, the contraction was initiated with noradrenaline (1 μ M) in calcium-free high K^{+} physiological solution. Mean am-

plitude of contraction induced by noradrenaline was compared to the contraction obtained in the presence of extract (1 mg/ml) which was introduced 20 minutes prior to noradrenaline addition.

Statistical analysis

Data are expressed as the mean \pm S.E.M and n indicates the number of animals. Statistical analysis was based on analysis of variance (ANOVA) followed by Fisher's LSD test using Systat (Version 11.0). Differences were considered to be significant when $p < 0.05$.

RESULTS

Total phenolic and flavonoid content in *Biophytum petersianum*

The results showed that the total phenolic and flavonoid in *B. petersianum* are respectively 205 mg GAE/g of BPE and 97.5 mg QE/g of BPE.

Effect of hydro-alcoholic extract of *B. petersianum* on Calcium channels activity

In Ca^{2+} -free, high K^+ (60 mM) solution, the cell membrane of aortic smooth muscle was depolarized, but the lack of Ca^{2+} entry in vascular smooth muscle cells prevented vasoconstriction in this high K^+ solution. Cumulative addition of calcium chloride (0.1–8 mM) caused an increase in contraction of the rat aorta (fig. 1). The maximal tension attained at 8 mM Ca^{2+} (1266.6 ± 8.8 mg) was considered as 100 %. When the aortic rings were treated with extract at 0.1; 0.25 and 0.5 mg/ml (20 minutes prior to CaCl_2 addition), the KCl-induced contraction was rightward shifted in non-parallel manner. The plant extract at 0.5 mg/ml significantly reduced the maximal tension induced by calcium chloride (74.8%, $p < 0.001$) which was comparable to that of verapamil at $1\mu\text{M}$ (83.2 %).

To evaluate the effect of *B. petersianum* on the release of intracellular calcium, noradrenaline-induced contraction was measured in presence of extract. Phasic contraction of rat aortic rings induced by noradrenaline in the absence and in the presence of plant extract was 1038 ± 104 mg and 351.66 ± 30.3 mg respectively (fig. 2), showing that contraction maximal amplitude induced by noradrenalin was significantly reduced ($p < 0.001$) by the hydro-alcoholic extract of *B. petersianum*.

DISCUSSION

The major finding in this study was that the hydro-alcoholic extract of *B. petersianum* acts as a non-selective calcium antagonist. Our study demonstrated that pre-treatment with the plant extract prevents, in a concentration-dependent manner, the aortic contractile response to high K^+ . It's well documented that the contractions of smooth muscles which are induced by high K^+ , are dependent of calcium entry into cell through voltage-operated calcium channels [12,13]. This suggested that the plant extract might interfere with Ca^{2+} -

sensitive voltage channels in aortic smooth muscle probably by inhibiting Ca^{2+} influx into the cell.

Noradrenaline-induced responses in free- Ca^{2+} , high K^+ medium, moreover, were significantly reduced in presence of 1 mg/ml of plant extract, suggesting extract also affected the release of Ca^{2+} from intracellular stores.

We have quantified compounds in the extract of *B. petersianum*, such as total phenolic and flavonoid. These findings are consistent with those of Lin et al. [14] and Bucal et al. [15] who found biflavone and flavonoid, and phenolic compounds from *Biophytum petersianum* (syn. *Biophytum sensitivum*) respectively. Several studies have revealed that phenolic compounds and flavonoids have calcium antagonist activity, and vasodilatory actions [11, 16-19]. However, further investigations are necessary to identify these compounds in *B. petersianum* and to confirm their calcium antagonistic effect.

In summary, the present study demonstrates that the hydro-ethanolic extract of *Biophytum petersianum* has calcium antagonistic activity on rat isolated aorta rings. The vasorelaxant effect of *B. petersianum* may be involved in its hypotensive effect. Further experiments are necessary to elucidate the active principles and their possible mechanism of action.

REFERENCES

1. Inngjerdingen KT, Coulibaly A, Diallo D, Michaelsen TE, Paulsen BS. A complement fixing polysaccharide from *Biophytum petersianum* klotzsch, a medicinal plant from Mali, West Africa. *Biomacromolecules* 2006; 7: 48-53.
2. Puri D. Hypcholesterolemic effect of *Biophytum sensitivum* leaf water extract. *Pharmaceut Biol* 2003; 41: 253-8.
3. Puri D, Baral N. Hypoglycemic effect of *Biophytum sensitivum* in the alloxan diabetic rabbits. *Indian J Physiol Pharmacol* 1998; 42: 401-6.
4. Puri D. The insulinotropic activity of a Napalese medicinal plant *Biophytum sensitivum*: preliminary experimental study. *J Ethnopharmacol* 2001; 78: 89-93.
5. Bucar F, Jachak SM, Noreem Y, Kartnig T, Perera P, Bohlin L, Schubert-Zsilavecz M. Amentoflavone from *Biophytum sensitivum* and its effect on COX-1/COX-2 catalysed prostaglandin biosynthesis. *Planta Med* 1998; 64: 373-4.
6. Titrikou S, Aklikokou AK, Gbeassor M. Effets de l'extrait de *Biophytum petersianum* (Oxalidaceae), Klotzsch, sur le système cardiovasculaire de cobaye. *Pharm Méd Trad Afr* 1998; 10: 32-42.
7. Kodjo KM, Contesse V, Do Rego JL, Aklikokou K, Titrikou S, Gbeassor M, Vaudry H. In vitro effects of crude extracts of *Parikia biglobosa* (Mimosaceae), *Stereospermum kunthianum* (Bignoniaceae) and *Biophytum petersianum* (Oxalidaceae) on corticosteroid secretion in rat. *J Steroid Biochem Mol Biol* 2006; 100: 202-8.
8. Triggle DJ, Swamy VC. Pharmacology of agents that affect calcium: Agonists and Antagonists. *Chest* 1980; 78: 174-9.
9. Hop RP, Vuorela HJ. Assessing calcium antagonism on vascular smooth muscle: a comparison of three methods. *J Pharmacol Methods* 1983; 9: 41-52.
10. Pourmorad F, Hosseini-mehr SJ, Shahabimajd N. Antioxydant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *Afr J Biotechnol* 2006; 5: 1142-5.
11. Dar A, Channa S. Calcium antagonistic activity of *Bacopa monniera* on vascular and intestinal smooth muscles of rabbit and guinea-pig. *J Ethnopharmacol* 1999; 66: 166-74.

12. Yeh JL, Liou SF, Liang JC, Lee CH, Chiu CC, Lin YT and Chen IJ. Labetidipinedilol-C: A third-generation dihydropyridine-type calcium channel antagonist displaying K⁺ Channel opening, NO-dependant and adrenergic antagonist activities. *J Cardiovasc Pharmacol* 2005; 46: 130-40.
13. Abreu IC, Marhino ASS, Paes AMA, Freire SMF, Olea RSG, Borges MOR and Borges ACR. Hypotensive and vasorelaxant effects of ethanolic extract from *Jatropha gossypifolia* L. in rats. *Fitoterapia* 2003; 74: 650-7.
14. Lin Y-L, Wang W-Y. Chemical constituents of *Biophytum sensitivum*. *Chin Pharmaceut J* 2003; 55: 71-5.
15. Bucar F, Jachak SM, Kartnig T, Schubert-Zsilavecz M. Phenolic compounds from *Biophytum sensitivum*. *Pharmazie* 1998; 53: 651-3.
16. Soobrattee MA, Neergheen VS, Luximon-Ramma A, Aruoma OI and Bahorun T. Phenolics as potential antioxidant therapeutic agents: Mechanism and actions. *Mut Res* 2005; 579: 200-13.
17. Attaguile G, Perticone G, Mania G, Savoca F, Pennisi G and Salomone S. *Citrus incanus* and *Citrus monspeliensis* inhibit the contractile response in isolated rat smooth muscle. *J Ethnopharmacol* 2004; 92: 245-50.
18. Geleijnse JM, Launer LJ, Van der Kuip DAM, Hofman A, and Witteman JCM. Inverse association of tea and flavonoid intakes with incident myocardial infarction: the Rotterdam Study. *Am J Clin Nutr* 2002; 75: 880-6.
19. Sesso HD, Gaziano JM, Liu S and Buring JE. Flavonoid intake and the risk of cardiovascular disease in women. *Am J Clin Nutr* 2003; 77: 1400-8.ref

CURRENT AUTHOR ADDRESSES

Siménou Titrikou, Département de Physiologie/Pharmacologie, Faculté des Sciences, Université de Lomé B.P. 1515 Lomé – TOGO.

Kwashie Eklu-Gadegbeku, Département de Physiologie/Pharmacologie, Faculté des Sciences, Université de Lomé B.P. 1515 Lomé – TOGO.

Aklesso Mouzou, Département de Physiologie/Pharmacologie, Faculté des Sciences, Université de Lomé B.P. 1515 Lomé – TOGO.

Kodjo Aklikokou, Département de Physiologie/Pharmacologie, and Centre de Recherche et de Formation sur les Plantes Médicinales (CERFOPLAM), Faculté des Sciences, Université de Lomé B.P. 1515 Lomé – TOGO.

Messanvi Gbeassor, Département de Physiologie/Pharmacologie, and Centre de Recherche et de Formation sur les Plantes Médicinales (CERFOPLAM), Faculté des Sciences, Université de Lomé B.P. 1515 Lomé – TOGO. E-mail: gbeassor@tg.refer.org (Corresponding author)