

## RESEARCH ARTICLE

## Regulatory role of Calcium Channel Blockers on spontaneous muscular activity of *Gastrothylax crumenifer*, a rumen amphistome

PAWAN KUMAR VERMA, DINESH KUMAR, S. K. TANDAN, ANIL KUMAR SRIVASTAVA and RAJINDER RAINA

For author affiliations, see end of text.

Received December 30, 2006; Revised May 8, 2007; Accepted August 5, 2007

This paper is available online at <http://ijpt.iuums.ac.ir>

### ABSTRACT

Major proportion of intracellular calcium ( $\text{Ca}^{2+}$ ) is achieved through opening of calcium channels present in the plasma membrane which play an important role in regulating neuromuscular coordination and release of neurotransmitters from nerve terminals. Blockade of these calcium channels adversely affects contractile process and release of neurotransmitters in majority of the neuromuscular preparations in vitro. In present study, the cumulative addition of verapamil ( $10^{-7}$ - $10^{-3}$  M) caused marked excitation in amplitude, baseline tension and frequency of spontaneous muscular activity of *Gastrothylax crumenifer* a rumen amphistome. Diltiazem ( $10^{-6}$ - $10^{-3}$  M) caused a significant and concentration-dependent increase in amplitude and frequency of spontaneous muscular activity of isometrically mounted rumen amphistome. It also caused significant rise in baseline tension at  $10^{-5}$  to  $10^{-3}$  M concentrations. Addition of nifedipine ( $10^{-7}$ - $10^{-3}$  M) elicited significant and concentration-dependent rise in amplitude and baseline tension, as compared to control values without significantly effecting frequency of spontaneous contraction.

**Keywords:** comma separated keywords

*Gastrothylax crumenifer* is a common rumen amphistome belonging to class trematode. The infestation of this parasite in ruminants results in decreased growth, production and reproductive performance of productive animals besides decreasing the quality and quantity of animal products [1]. Synthetic anthelmintics currently in use have long been considered the only effective way of controlling these parasitic infections. Injudicious and frequent use of these anthelmintics has resulted in the development of resistance. Furthermore, residual toxicity and adverse reactions in animals have been associated with the available synthetic anthelmintics [2,3]. Therefore, there is a need to develop specific drug(s) targeting various macro-molecular components of these parasites.

Neuromuscular system of helminthes is an important area for target identification and drug development. Acetylcholine, a major inhibitory neurotransmitter of trematodes has been demonstrated immunocytochemically to be present in peripheral and central nervous system [4]. Calcium ions ( $\text{Ca}^{++}$ ) play an important role in neurotransmitter release from the nerve terminals and neuromuscular coordination [5]. Bathing

medium free of  $\text{Ca}^{++}$  ions reduces the spontaneous muscular activity of *Schistoma mansoni* [6]. Similarly increasing external  $\text{Ca}^{++}$  ions concentration in the medium mimics the inhibitory effect of ACh on spontaneous muscular activity of split preparation of adult *Fasciola hepatica* and *Hymenolepis diminuta* [7,8]. However, calcium channels blockers, diltiazem and verapamil, cause marked stimulation followed by paralysis of *Schistoma mansoni* and *Fasciola gigantica*, respectively. [9,10]. Contractions induced by calcium-dependent depolarization have been observed in dispersed muscle fibres of *Schistoma mansoni* [11] whereas nifedipine, a calcium channel blocker, blocks these contractions [12].

Calcium currents have been recorded from muscle fibers of *Bdelloura candida*. However,  $\text{Ca}^{++}$  currents could not be recorded from muscle fibers of *S. mansoni* [13] and *F. gigantica* [14]. Recently, a number of components vital for  $\text{Ca}^{++}$  storage and release involving Ryanodine receptors (RyR) present in sarcoplasmic reticulum have been demonstrated in the genera of *Schistosomes* [15,16].

Table 1. Effects of verapamil, diltiazem, nifedipine, and on amplitude (g) baseline tension (g) and frequency (per 5 min) of spontaneous muscular activity of *Gastrothylax crumenifer*

Observations	Concentrations					
	Control	10 <sup>-7</sup> M	10 <sup>-6</sup> M	10 <sup>-5</sup> M	10 <sup>-4</sup> M	10 <sup>-3</sup> M
<b>Diltiazem</b>						
Amplitude (g)	0.49±0.03	0.51±0.03	0.56±0.04**	0.61±0.06**	0.64±0.06**	0.68±0.05***
Baseline tension (g)	0.20±0.02	0.21±0.02	0.23±0.03	0.24±0.02**	0.27±0.03**	0.29±0.02***
Frequency/5min.	43±4.09	47.5±6.14	49.5±5.20*	50±5.42*	54±4.11*	55.5±5.85*
<b>Verapamil</b>						
Amplitude (g)	0.44±0.05	0.45±0.05	0.47±0.04	0.55±0.03	0.58±0.03*	0.75±0.08***
Baseline tension (g)	0.15±0.01	0.17±0.02	0.19±0.02*	0.22±0.02**	0.24±0.02***	0.27±0.03***
Frequency/5min.	46.5±3.94	48.5±5.41	49.5±3.18	50±2.65	51±5.38**	53.5±4.69*
<b>Nifedipine</b>						
Amplitude (g)	0.58±0.05	0.67±0.04*	0.73±0.08*	0.75±0.06***	0.77±0.09***	0.80±0.09***
Baseline tension (g)	0.22±0.02	0.24±0.02	0.25±0.02	0.26±0.03*	0.29±0.03**	0.30±0.03***
Frequency/5min.	54±4.39	58.5±3.54	61±5.19	60±4.53	57±3.47	55±4.44

\*= $p<0.05$ ; \*\*= $p<0.01$ ; \*\*\*= $p<0.001$ ; as compared to the controls

The present study was planned to investigate the role of different groups of voltage sensitive calcium channel blockers on spontaneous muscular activity of isometrically mounted parasitic rumen amphistome, *Gastrothylax crumenifer*.

## MATERIAL AND METHODS

### Collection of rumen amphistomes

Mature and healthy *Gastrothylax crumenifer* were collected from the rumen of freshly slaughtered goats at local abattoir in warm (38±1°C) Hank's Balanced Salt Solution (HBSS) in an insulated container and brought to the laboratory. They were kept in the BOD incubator at 38±1°C until further use. The amphistomes (*G. crumenifer*) were identified before experimentation.

### Tissue preparation and mechanical recording

The spontaneously active whole mature amphistome was mounted isometrically in HBS solution at 38±1°C as per the method described for *Gigantocotyle explanatum* [17]. In short, the amphistome was mounted with the help of two fine hooks. One hook was inserted 1-2 mm caudal to anterior sucker and fixed to the tip of aeration tube and another hook was pierced through the surface of acetabulum and connected to the isometric force transducer. The spontaneous muscular activity of isometrically mounted amphistome was recorded in Chart Window 4 Software programme. (Powerlab, AD Instruments, Australia).

### Experimental protocol

Graded molar concentrations (10<sup>-7</sup>- 10<sup>-3</sup> M) of different groups of calcium channel blockers; verapamil (Phenylalkylamine derivative) (Sigma, USA), diltiazem (benzothiazepine derivative) (Sigma, USA) and nifedipine (dihydropyridine derivative) (Sigma, USA) were applied after equilibration of the fluke to examine their effects on spontaneous muscular activity of *G. crumenifer*.

### Data collection and statistical analysis

Three attributes, viz., the amplitude (average of all peaks per five minutes), baseline tension (average of all minimum levels of contractions used for measuring amplitude) and frequency (total number of contractions in

95 min) of spontaneous muscular contractions were utilized to evaluate the effect of different concentrations of different groups of calcium channel blockers. Experiment with each drug was repeated at least six times on fresh isometrically mounted *G. crumenifer*.

The results are presented as mean ± standard error of mean and paired 't' test was applied to measure the level of significance [18].

## RESULTS

The isometrically mounted amphistomes exhibited rhythmic phasic contractile activity continuously for 24 hours. The mean amplitude, baseline tension and frequency of the rhythmicity recorded every 15 min after applying the tension of 200 mg, were 0.42 ± 0.03 g (n=6), 0.13 ± 0.01 g (n=6) and 45 ± 2.90 contractions/5 min time period (n=6), respectively. The isometrically mounted amphistomes exhibited apparently uniform pattern of spontaneous muscular activity for a period of 24 h. There was no significant difference in amplitude (0.38 ± 0.02 g; n=6), baseline tension (0.11 ± 0.02 g; n=6) and frequency (41 ± 3.44/ 5 min; n=6) of spontaneous contractions recorded 2 h after mounting as compared with those recorded 15 min after applying the tension to the amphistome. The representative recording is given in Fig. 1.

### Effect of calcium channel blockers on spontaneous muscular activity of *G. Crumenifer*

Diltiazem, a benzothiazepine derivative causes a significant and concentration-dependent (10<sup>-6</sup> to 10<sup>-3</sup> M) increase in amplitude and frequency of spontaneous muscular activity as compared with control amplitude (0.49 ± 0.03 g) and frequency (43.0 ± 4.09/ 5 min). It also caused marked rise in baseline tension at 10<sup>-5</sup> to 10<sup>-3</sup> M concentrations in a concentration-dependent manner as shown in Table 1 and Fig 2a and 2b.

Effects of verapamil, a phenylalkylamine derivative on control amplitude, baseline tension and frequency of spontaneous contractions of isometrically mounted rumen fluke is shown in Table 1, and Fig 3a and 3b. Verapamil in cumulative concentrations at an increment of 1 log unit produced significant excitation in the amplitude (at 10<sup>-4</sup> and 10<sup>-3</sup> M), baseline tension (10<sup>-6</sup> to 10<sup>-3</sup> M) and frequency (at 10<sup>-4</sup> and 10<sup>-3</sup> M) of spontaneous muscular activity of the rumen fluke.




Fig. 1. Time-dependent control recording of spontaneous muscular activity of *G. crumenifer* (0 to 120 min)




Fig. 2a. Effects of different concentrations ( $10^{-7}$ M to  $10^{-3}$ M) of Diltiazem on spontaneous muscular activity of *G. crumenifer*




Fig. 3a. Effect of different concentrations ( $10^{-7}$ M to  $10^{-3}$ M) of Verapamil on spontaneous muscular activity of *G. crumenifer*




Fig. 4a. Effects of different concentrations ( $10^{-7}$ M to  $10^{-3}$ M) of Nifedipine on spontaneous muscular activity of *G. crumenifer*




Fig. 2b. Effect of different concentrations ( $10^{-7}$  M to  $10^{-3}$  M) of Diltiazem on amplitude (g) baseline tension (g) and frequency (per 5 min) of spontaneous muscular activity of *G. crumenifer*




Fig. 4b. Effect of different concentrations ( $10^{-7}$  M to  $10^{-3}$  M) of Nifedipine on amplitude (g) baseline tension (g) and frequency (per 5 min) of spontaneous muscular activity of *G. crumenifer*




Fig. 3b. Effect of different concentrations ( $10^{-7}$  M to  $10^{-3}$  M) of Verapamil on amplitude (g) baseline tension (g) and frequency (per 5 min) of spontaneous muscular activity of *G. crumenifer*

176 with a variation in the intensity. Out of these different  
177 groups dihydropyridine derivatives are producing pro-  
178 nounced contractile responses as compared to other  
179 blockers. The results are in agreement with the earlier  
180 reports on *S. mansoni* [9] and *F. gigantica* [10]. How-  
181 ever, it has been shown that Ca<sup>2+</sup>-free bathing medium  
182 reduced the spontaneous muscular activity of *S. man-*  
183 *soni* [6], while increasing external calcium ion concen-  
184 tration, mimicked the inhibitory effects of ACh on spon-  
185 taneous muscular activity of split-preparation of adult *F.*  
186 *hepatica* [7]. The probable mechanism may be that  
187 these calcium channel blockers inhibit the release of  
188 inhibitory neurotransmitters at the nerve terminals as it  
189 is well documented that the release of neurotransmitter  
190 at nerve terminal requires Ca<sup>2+</sup> [5] or these calcium  
191 channel blockers may be producing a direct stimulatory  
192 effect on trematode neuromuscular system. Verapamil  
193 and diltiazem have significant stimulatory effects on  
194 muscular contraction whereas Nifedipine did not have  
195 significant effect on the frequency of the activity. This  
196 may be due to reversal of action of sodium-calcium ex-  
197 changer proteins resulting from blocking of calcium  
198 channels [19]. It is likely that similar mechanism operates  
199 in muscular tissue of *G. crumenifer* and may be respon-  
200 sible for Nifedipine not to significantly increase the fre-  
201 quency of spontaneous muscular activity in *G. cru-*  
202 *menifer*. It will be interesting to study the exact mecha-  
203 nism for this excitatory effect on spontaneous muscular  
204 activity with calcium channel blockers in trematodes at  
205 molecular level.

## DISCUSSION

164 Intracellular calcium (Ca<sup>2+</sup>) is responsible for the  
165 muscular contraction and release of neurotransmitters  
166 from nerve terminals in mammals. Removal of extra-  
167 cellular Ca<sup>2+</sup> and /or blockade of calcium channels ad-  
168 versely affect contractile process and release of neuro-  
169 transmitters in majority of the neuromuscular prepara-  
170 tions *in vitro*. In the present study, however, calcium  
171 channel blockers from different groups elicited an exci-  
172 tatory response in amplitude and baseline tension of  
173 muscular activity of *G. crumenifer*. The effect of the  
174 calcium channel blockers on amplitude and baseline  
175 tension in all groups show similar excitatory patterns

## REFERENCES

- 206  
207 Prasad, A., Varma, T.K. On the prevalence and community  
208 dominance among paramphistomes infecting domestic rumi-  
209 nants, *Journal of Veterinary Parasitology*, 1999; 13(2), 129-133.  
210 Sangster, N.C., Gill, J. Pharmacology of anthelmintic resistance,  
211 *Parasitology Today*, 1997; 15, 141-146.  
212 Prichard, R.K. Is anthelmintics resistance a concern for heart-  
213 worm control? What can we learn from the human filariasis con-  
214 trol program?, *Veterinary Parasitology*, 2005; 133: 243-253.  
215 Maule, A.G., Halton, D.W., Johnston, C.F., Show, C., Fair-  
216 weather, I. Serotonergic, cholinergic and peptidergic compo-

- 217 nents of the nervous system in the monogenean parasite, *Dicli-*253  
218 *dophora merlangi*: a cytochemical study, *Parasitology*, 1990; 254  
219 100, 255-273. 255
2205. Katz, B., Miledi, R. The timing of calcium action during neuro-256  
221 muscular transmission, *Journal of Physiology*, 1967; 189, 535-257  
222 544. 258
2236. Wolde Mussie, E., Vande Waa, J., Pax, R. A., Fetterer, R., Ben-259  
224 nett, J. L. *Schistosoma mansoni*: Calcium efflux and effects of 260  
225 calcium-free media on responses of the adult male musculature 261  
226 to praziquantel and agents inducing contractions. *Experimental* 262  
227 *Parasitology*, 1992; 53, 270-278. 263
2287. Sukhdeo, S.C., Sangster, N.C., Mettrick, D.F. Effects of cho-264  
229 linergic drugs on longitudinal muscle contraction of *Fasciola he-*265  
230 *patica*, *Journal of Parasitology*, 1986; 72, 492-497. 266
2318. Thompson, C. S., Mettrick, D. R. Neuromuscular physiology of 267  
232 *Hymenolepis diminuta* and *H. microstoma* (Cestoda), 1984; 268  
233 *Parasitology*, 89, 567-578. 269
2349. Senft, W.A., Gibler, W.B., Guterman, J.J. Influence of calcium 270  
235 perturbing agents on schistosomes : Comparison of effects of 271  
236 praziquantel and verapamil on worm tegument, *Journal Experi-* 272  
237 *mental Zoology*, 1986; 239, 25-36. 273
23810. Kumar, D., Tripathi, H.C. Calcium dependence of stimulatory 273  
239 effect of calcium channel blockers on *Fasciola gigantica*, *Journal* 274  
240 *of Veterinary Parasitology*, 2000; 14(1), 35-38. 275
24111. Day, T. A., Bennett, J. L., Pax, R. A. Serotonin and its require- 276  
242 ment for the maintenance of contractility in muscle fibres iso- 277  
243 lated from *Schistosoma mansoni*, *Parasitology*, 1994a; 108, 425- 278  
244 432. 279
24512. Day, T.A., Maule, A.G., Shaw, C., Halton, D.W., Moore, S., 280  
246 Bennett, J.L., Pax, R.A. Platyhelminth FMRF amide, related 281  
247 peptides (FaRPs) contract *Schistosoma mansoni* (Trematoda : 282  
248 Digenea) muscle fibres in vitro, *Parasitology*, 1994b; 109(4), 283  
249 455-459. 284
25013. Blair, K. L., Anderson, P. A. V. Physiological and pharmaco- 285  
251 logical properties of muscle cells isolated from the flatworm, 286  
252 *Bdelloura candida* (Tricladia), *Parasitology*, 1994; 109, 325-335. 286
- Kumar, D., Chandra, S., Tripathi, H.C. In vitro motility re-  
cording of *Fasciola gigantica*, *Journal of Veterinary Parasitol-*  
*ogy*, 1995; 9, 31-36.
- Talla, E. De, Mendonca, R. L., Degand, I., Goffeau, A., Ghis-  
lain, M. *Schistosoma mansoni* Ca<sup>++</sup>-ATPase SMA2 restores vi-  
ability to yeast Ca<sup>2+</sup>-ATPase-dependent strains and functions in  
calcineurin-mediated Ca<sup>2+</sup> tolerance. *Journal of Biological*  
*Chemistry*, 1998; 273, 27831-27840.
- Silva, C. L. M., Cunha, V. M. N., Mendonca, Silva, D. L., Noel,  
F. Evidence for ryanodine receptors in *Schistosoma mansoni*,  
*Biochemical Pharmacology*, 1998; 56, 997-1003.
- Ahmad, M., Nizami, W.A. In vitro effect of some anthelmintics  
on the motility of *Gigantocotyle explanatum*, *Japanes Journal of*  
*Parasitology*, 1990; 39(6), 529-534.
- Snedecor, G. W., Cochran, W. J. *Statistical Methods*, 1989; 61p,  
Oxford IBH Co., Bombay.
- Kohomoto, O. Levi, A. J. and J. H. Relation between reverse  
sodium-calcium exchange and sarcoplasmic reticulum calcium  
release in guinea pig ventricular cells, *Circulation research*,  
1994; 74, 550-554.

## CURRENT AUTHOR ADDRESSES

Pawan Kumar Verma, Junior Scientist  
Division of Pharmacology and  
Toxicology F.V.Sc. & A.H. R.S.Pura, Jammu, J&K, India. E-  
mail: drpawankv@yahoo.co.in (Corresponding author)

Dinesh Kumar, Senior Scientist, Division of Pharmacology and Toxi-  
cology, Indian Veterinary Research Institute, Izatnagar, Bareilly -  
243 122 (UP), India.

S. K. Tandan, Senior Scientist, Division of Pharmacology and Toxi-  
cology, Indian Veterinary Research Institute, Izatnagar, Bareilly -  
243 122 (UP), India.

Anil Kumar Srivastava, Dean, Faculty of Veterinary Sciences and  
Animal Husbandry, SKUAST-J, R.S. Pura, Jammu (J&K) India.

Rajinder Raina, Head, Division of Pharmacology and Toxicology,  
F.V.Sc. & A.H., SKUAST-J, R.S. Pura, Jammu (J&K), India.