

# Electrical Lesion of Substantia Nigra Pars Compacta: An Alternative and Convenient Way to Generate the Animal Model of Parkinson's Disease

HADI FATHI MOGHADDAM and MEHDI SHAFIEE ARDESTANI

*For author affiliations, see end of text.*

Received September 8, 2007; Revised August 29, 2008; Accepted September 21, 2008

This paper is available online at <http://ijpt.iuums.ac.ir>

## ABSTRACT

The unilateral or bilateral lesions of the Substantia nigra pars compacta (SNc) by chemical neurotoxins such as 6-hydroxydopamine (6-OHDA) or 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) are main method for generating the parkinson's animal models. But these neurotoxins are expensive, dangerous, difficult to use and inaccessible conveniently due to the international economical sanctions in Iran. The present study was designed to find a new animal model for Parkinson's disease. Forty eight animals were included in this study and divided into six groups. One group was selected as the control and also two groups as the sham of the SNc lesion. Other groups were subjected to lesion of SNc by 6-OHDA, MPTP and electrical lesion. Using histological studies and Murprogo's method precision, the effectiveness of the electrical lesion was compared with that of chemical lesions. Our results showed the same extent of lesion was provided with electrical insult and neurotoxins. Due to the technical convenience, low cost, accessibility and the same potency of electrical- and neurotoxins-induced lesion, this model can be used as an alternative method for creating the Parkinson's models.

**Keywords:** 6-hydroxydopamine (6-OHDA), Electrical lesion, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP), Substantia nigra pars compacta, Parkinson's animal model

Parkinson's disease (PD) is a degenerative neuro-dopaminergic disease in nigrostriatum pathway of human, animal. The resultant loss of nerve terminals accompanied by dopamine deficiency in this pathway is responsible for most of the movement disorders [1]. Several reports indicate SNc damages are induced by several agents such as free radicals, 6-hydroxydopamine (6-OHDA), 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP), reserpine, phenithiazine and butyrophenones [2,3]. The damage has also been observed in encephalitis and aging [4,5]. Too many efforts have been done to achieve the effective treatment of Parkinson's disease. Many of these researches have been carried out in animal models induced by chemical neurotoxins such as MPTP [4].

SNc lesion in animals like rat leads to rigidity due to decreasing the inhibitory dopamin effects on the caudate nucleus and putamen, as the main rigidity-inductor neurotransmitter-releasing areas, located in the striatum. Because of these changes in the brain of the lesioned rat, rigidity was occurred on the limbs on both the sides [6]. The unilateral or bilateral lesion of the SNc by

chemical neurotoxins such as 6-OHDA or MPTP is the most commonly and the main used methods for creating the PD animal models. The use of MPTP is highly dangerous. The compound is able to be absorbed from the skin, gastrointestinal tract and blood brain barrier. Furthermore, MPTP as well as 6-OHDA destroys any catecholaminergic parts in the brain; which leads to complications [7]. Also, preparation and use of these compounds are very expensive and time consuming. These disadvantages of 6-OHDA and MPTP were encouraged us to find another possible, low cost, technically-convenient and less time-consuming method. Mimicking some previously-published works, we used electrical lesion of substantia nigra pars compacta [8]. Electrical lesion can destroy SNc and create an animal model of PD.

## MATERIALS AND METHODS:

### Animals

Forty eight male albino Wistar rats (200-250g) were used in the present study. The animals were purchased

**Table 1.** Mean of rigidity grades as  $M \pm S.E.M$  in investigated groups (N=8)

Groups	Time (min)							
	0	20	40	60	90	120	180	240
Control	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0
Sham of electrical lesion	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0
Sham of 6-OHDA and MPTP lesion	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0
Electrical lesion	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0
6-OHDA lesion	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0
MPTP lesion	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0	3.5±0

from Pasteur Institute of Iran and then housed in groups of ten in stainless steel cages, handled daily, and were provided food and water ad libitum. A 12-h light/12-h dark cycle was maintained, and the animals were tested during the light cycle. The experiments were carried out in accordance with recommendations from the declaration of Helsinki and the internationally-accepted principles in the use of experimental animals. In this study, we divided animals into six groups, each eight rats.

#### Drugs and Solvents

6-OHDA HBr and MPTP were purchased from Sigma (Chemical Co., St. Louis, MO, USA) and ascorbic acid was purchased from Merck (Germany).

#### General principles of the Surgery

Each rat was anesthetized separately by injection of 75 mg/kg ketamine combined with 8 mg/kg Xylazin intraperitoneally. Then, we prepared the rats for surgery and placed them in the stereotaxic apparatus (Stoelting, IL, USA). The left SNc, region of the nigrostriatum was targeted. Stereotaxic coordinators for the left SNc region were set at -4.8 mm posterior and -1.6 mm lateral to bregma and 8.1 mm for 6-OHDA and MPTP or 8.2 mm for electrical lesion ventral to the surface of the skull according to the atlas of Paxinos and Watson. [9]. It should be notified that we used unilateral lesion of SNc in all investigated lesioned groups.

#### General procedure of the electrical lesion

The stainless steel electrode was placed in the left SNc and destroyed the area by Electrical Lesion Maker (Siemens Company, Germany), using electrical current (1mA, 10 seconds). Then, they were kept in individual cages for 7-10 days after the surgery for recovery.

#### General procedure of the 6-OHDA lesion

The skull was exposed and a hole was drilled through the skull in the area overlying the left SNc. A guide-cannula was lowered into the brain and fixed to the cranium with miniature screws and acrylic dental cement and the incision was closed with sutures. Surgery was performed using sterile instruments and aseptic conditions. Then 6-OHDA HBr (10µl of a solution based on 66.7 µg, base form, in ascorbic acid 0.1%) was injected into the left SNc. This procedure has been described in detail previously [10]. Rats were allowed to

recover from the surgery for 7–10 days in individual transparent test cages.


#### General procedure of the MPTP lesion was as follows:

The skull was exposed and a hole was drilled through the skull in the area overlying the left SNc. A guide-cannula was lowered into the brain and fixed to the cranium with miniature screws and acrylic dental cement and the incision was closed with sutures. Surgery was performed using sterile instruments and aseptic conditions. Then MPTP (5 µl of a solution based on 2 mg MPTP in sterile 0.9% saline) was injected into the left SNc. This procedure has been described in detail previously [11]. Rats were allowed to recover from the surgery for 7–10 days in individual transparent test cages.

#### Rigidity evaluations:

At the time of study, all animals exhibited rigidity, a loss of vocalisation, diminished blinking, incoordination and a course action tremor. Murprogo's Method [12] was used to measure the rigidity of all subjected animals at the times: 0, 20, 40, 60, 90, 120, 180 and 240 min. The wood-platforms with the steps of 3 and 9 cm were used in this study. For the procedure of behavior experiments, the animal was put on the bench. when the animal did not move by touch, it received the score of 0.5. Then, the right hand of the animal was placed on the wood-platform with the height of 3 cm. If the animal did not take its hand off the platform after at least 10 seconds, it received the score of 0.5. Rigidity evaluation was repeated for the left hand of the animal as well. In the next stage of the procedure, the right hand of the animal was placed on the wood-platform with the height of 9 cm, so that any other parts of the animal did not touch the platform. The animal was given score of 1, if it did not take its hand off the platform after 10 seconds. Finally, the test was repeated in the same way for the left hand. Each animal that had full rigidity (PD) was given a total score of 3.5. The results of the Murprogo's test are mentioned in Table 1.

The effects of neurotoxin vehicles on the rigidity or cellular damage have not been investigated because the previous studies have proven the inertness of saline or ascorbic acid in the models (3, 7, 11). Also, unnecessary use of animals is in contrast with internationally-accepted principles in the use of experimental animals.



**Fig 1.** Histological study about accuracy of the electrical lesion of SNc in comparison with 6-OHDA- and MPTP-induced lesion and control groups. (A) SNc was painted in control and sham groups, without microscopic magnification. (B) SNc was painted in control and sham groups, with microscopic magnification (4X). (C) Destroyed SNc in Parkinsonian rats using electrical lesion didn't paint, without microscopic magnification. (D) Destroyed SNc in Parkinsonian rats using electrical lesion didn't paint, with microscopic magnification (4X). (E) Destroyed SNc in Parkinsonian rats using 6-OHDA lesion didn't paint, without microscopic magnification. (F) Destroyed SNc in Parkinsonian rats using 6-OHDA didn't paint, with microscopic magnification (4X). (G) Destroyed SNc in Parkinsonian rats using MPTP didn't paint, without microscopic magnification. (H) Destroyed SNc in Parkinsonian rats using MPTP didn't paint, with microscopic magnification (4X).

#### *Evaluations the electrical method*

##### *In vitro investigation of electrical lesion method*

Estimation of violence and duration of the lesion was accepted empirically in vitro by determination of clot-dimensions and comparison with the atlas of Paxi-

nos and Watson in electrocardiograph gel caused by electrical maker.

##### *In vivo investigation*

The experiments including the Murprogo's method and histological studies were carried out to evaluate and

compare the electrical lesion with 6-OHDA- and MPTP-induced lesion. After the rigidity test, each suspected animal to PD was decapitated and the brain was removed and kept in a 10% formalin solution. Selected brains were cut on a cryostat as 50  $\mu$ m thick coronal sections, mounted on glass slides, and stained with H&E. Sections were examined under a light-microscope to find the accuracy of lesion of the left SNc. Finally, the histological results were compared together. The Fig. 1 shows the accuracy and the precision of the lesion.

## RESULTS

### *Effects of different types of the SNc lesion on rigidity*

All of SNc-lesioned rats were given a total score of 3.5 for rigidity test. It is noteworthy that electrical lesion of SNc produced animal model of PD similar to the model produced by 6-OHDA and MPTP.

### *Histological comparisons between lesion methods*

Histological studies showed the same potency, precision and accuracy of SNc lesion for electrical lesion in comparison with chemical neurotoxins, as shown in Fig. 1.

## DISCUSSION

According to the results of this study, electrical lesion of SNc can be employed as an alternative and useful method for creating the PD models with low cost and conveniently-technical procedure because it does not need to use cannula or preparation of dangerous neurotoxins. Also, at this time our country suffers a lot of economical sanctions and access to chemical neurotoxins is not easy. Therefore, the electrical lesion method at this time can solve the neuroscientists' problems in creating PD model as an alternative method.

## ACKNOWLEDGEMENTS

This study has been supported by research council Tehran University of Medical Sciences and Jondishapour University of Medical Sciences. The authors sincerely would like to thank Dr. Northman, professor

of neuropathology, for providing 6-OHDA and MPTP as well as consulting in histological studies.

## REFERENCES

1. McGeer PL, McGeer EG. Innate immunity, local inflammation, and degenerative disease. *Sci Aging Knowledge Environ* 2002; 29: 3.
2. Reichman H, Lestienne P, Jelliner K. Parkinson's disease and the electron transport chain post mortem brain. In: Narabayashi H, Nagatsu T, Yanayisava N, Mizuno N. Parkinson's Disease: from basic research to treatment. New York: Raven company, 1993.
3. Chio D, Rothman S. The role of glutamate neurotoxicity in hypoxic-ischemic death. *Ann Neurosci*. 1990; 13: 171-8.
4. Simon R, Aminof A, Greenberg D. Clinical neurology. 4<sup>th</sup> ed. Stamford: Appleton and Lange, 1999.
5. Schapira H, Mann V, Cooper J. Anatomic and disease specificity of NADH co G1 reductase (complex) deficiency Parkinson's disease. *J Neurochem* 1990; 55: 2142-5.
6. Katzung BG. Basic and clinical pharmacology. New York: Appleton and Lange, 2007.
7. Gerlach M, Riederer P. Animal models of Parkinson's disease: an empirical comparison with the phenomenology of the disease in man. *Neural Transm* 1996; 103: 987-1020.
8. Fathi Moghaddam H, Kesmati M, Kargar HM. The effect of paraventricular nucleus lesion on conditioned place preference (CPP) in presence or absence of alpha-2 adrenergic agonist (clonidine) in male rats. *Acta Physiol Hung*. 2006; 93: 33-40.
9. Paxinos G, Watson C. The rat brain in stereotaxic coordinates. San Diego: Academic Press, 1997.
10. Kostrzewa RM, Gong L. Supersensitized D1 receptors mediate enhanced oral activity after neonatal 6-OHDA. *Pharmacol Biochem Behav* 1991; 39, 677-82.
11. Costa S, Irvani MM, Pearce RK, Jenner P. Glial cell line-derived neurotrophic factor concentration dependently improves disability and motor activity in MPTP-treated common marmosets. *Eur J Pharmacol* 2001; 412: 45-50.
12. Murprogo C. Effect of antiparkinson drug on a phenothiazine-induced catatonia reaction. *Arc Int Pharma Co Dyn* 1962; 137: 48-90.

## CURRENT AUTHOR ADDRESSES

Mehdi Shafiee Ardestani, Department of Medicinal Chemistry and Radiopharmacy, Tehran University of Medical Sciences, Faculty of Pharmacy, Tehran, Iran. E-mail: mehdishafieea@aol.com (Corresponding author)

Hadi Fathi Moghaddam, Department of Physiology and Physiology Research Center, Jondishapour University of Medical Sciences, Ahwaz, Iran.