

1 RESEARCH ARTICLE

2 Toxicity of Plant Derived Molluscicides in Attractant 3 Food Pellets against Snail, *Lymnaea Acuminata*

4 FARINDRA TIWARI and DINESH KUMAR SINGH

5 For author affiliations, see end of text.

6 Received December 6, 2006; Accepted June 30, 2007

7 This paper is available online at <http://ijpt.iuims.ac.ir>8 **ABSTRACT**

9 Use of molluscicides in the attractant food pellet (AFP) is one of the effective methods of snail control.
10 attractant food pellets containing starch and agar plus different concentrations of these molluscicides
11 (*Azadirachta indica* bark powder, *Allium sativum* bulb powder, *Polygonum tuberosum* bulb powder, *Annona*
12 *squamosa* seed powder, their active components azadirachtin, allicin, hecogenin, acetogenin; herbal mol-
13 luscicide pestoban and a synthetic molluscicide, Snail Kill, were tested for molluscicidal activity for 144h
14 against the snail, *Lymnaea acuminata*. Active components of all the plant derived molluscicides were
15 highly toxic to *L. acuminata* compared with their crude forms. The stability of bait formulations was stud-
16 ied by storing the pellets up to 4 weeks. Storage of molluscicide baits caused higher reduction in their
17 toxicity in comparison with synthetic molluscicides.

18 **Keywords:** Attractant food pellets, Bait formulation, *Fasciola*, *L. acuminata*, Molluscicides, Starch

19 Liver- flukes *Fasciola hepatica* Linnaeus and
20 *Fasciola gigantica* Cobbold cause endemic fascioliasis
21 in cattle population of eastern Uttar Pradesh [1-2]. The
22 snail *Lymnaea (Radix) acuminata* Lamarck (*Lymnaei-*
23 *idae*) is the vector of these flukes. One way to reduce the
24 incidence of fascioliasis is to de-link the life cycle of
25 fluke by destroying the intermediate hosts [3-8]. The
26 development of a selective and safe molluscicide should
27 always be a realistic goal. It must be effective at low
28 concentrations and exert minimal adverse effect on the
29 other biota sharing the same habitat with snail. Lack of
30 contact between molluscicides and target snail popula-
31 tion due to meshy vegetation, dilution in upwelling
32 sewage water are two main causes of the failure of snail
33 control programme. The snails use chemical signals for
34 locating food sources. These signals are released from
35 the dead and living aquatic organisms into the modular
36 system of the snails [9-13]. Starch is the strongest at-
37 tractant for *L. acuminata* [14]. Bait formulation contain-
38 ing attractant and a molluscicide is an expedient ap-
39 proach in order to lure the target snail population to the
40 molluscicide. In the present study different plant derived
41 molluscicides have been used along with starch in bait
42 formulation against *L. acuminata*.

43 **MATERIALS AND METHODS**

44 Agar, starch, different plant derived molluscicides
45 such as *Allium sativum* and *Polygonum tuberosum* bulb
46 powder, *Annona squamosa* seed powder and
47 *Azadirachta indica* bark powder and their active com-
48 ponents allicin, hecogenin, acetogenin and azadirachtin
49 (Sigma Chemical Co. USA), Snail Kill (metaldehyde-
50 Pesticide India) and herbal molluscicide, Pestoban (Liq-
51 uid concentrate of *Cedrus deodara*, *Azadirachta indica*
52 and seed powder of *Embelia ribes* in 90:2:1 ratio- In-
53 dian Herbs, Research and Supply Co. Pvt. Ltd., India)
54 were used in bait formulation. Adult *L. acuminata*
55 (2.25±0.20 cm in length) were collected locally from
56 lakes and low lying submerged fields in Gorakhpur. The
57 snails were acclimatized for 72 hours in dechlorinated
58 tap water at 25±1° C. The pH of the water was 7.1-7.3
59 and dissolved oxygen, free carbon dioxide and bicar-
60 bonate alkalinity were set to 6.5-7.2 mg/l, 5.2-6.3 mg/l
61 and 102.0-105.0 mg/l, respectively.
62 Attractant food pellets (AFP) were prepared accord-
63 ing to previous method [15] as modified by us [16]. 10
64 grams of starch (10 mM) was added to 2% agar solu-
65 tion. After boiling, each of the selective molluscicides
66 were added to the solution in different concentrations
67 (Table1), the mixture was stirred constantly for 30 min-
68 utes and spread to a uniform thickness (5 mm). After

Table 1. Mean number of snail *L. acuminata* in zone three in contact with the attractant food pellets (AFP) that contain different molluscicides after two hours from beginning of experiment.

Molluscicides	Concentration of molluscicides				
	0.1%*	0.2%	0.5%	0.7%	1.0%
<i>A. squamosa</i> (SP)	1.16±0.16 (58.0) +	0.83±0.16 (43.0)	1.33±0.21 (53.4)	1.67±0.21 (45.6)	0.83±0.16 (50.0)
<i>A. sativum</i> (BP)	3.67±0.2 (78.0) +	3.5±0.96 (53.8)	3.5±0.22 (61.8)	3.0±0.44 (50.0)	2.67±0.42 (47.0)
<i>P. tuberosa</i> (BP)	3.0±0.25 (60.0) +	2.67±0.21 (57.2)	2.33±0.21 (53.8)	1.83±0.40 (52.2)	1.67±0.21 (45.3)
<i>A. indica</i> (Ba P)	3.5±0.34 (60.0) +	1.83±0.16 (37.8)	3.0±0.22 (55.6)	2.33±0.42 (53.8)	0.83±0.30 (31.2)
Acetogenin	3.0±0.36 (50.0) +	2.0±0.25 (50.0)	1.16±0.3 (27.8)	1.5±0.22 (42.8)	0.5±0.22 (20.0)
Allicin	1.4±0.35 (46.7) +	1.33±0.21 (46.9)	1.0±0.63 (33.3)	1.5±0.34 (42.8)	0.83±0.16 (27.6)
Hecogenin	0.67±0.21 (50.3) +	0.33±0.21 (22.0)	0.5±0.22 (33.3)	0.83±0.30 (27.6)	0.33±0.21 (13.2)
Azadirachtin	1.16±0.16 (46.4) +	1.0±0.25 (31.5)	1.0±0.44 (33.3)	0.83±0.30 (33.3)	0.67±0.42 (21.2)
Snail Kill	1.67±0.21 (45.5) +	1.5±0.34 (42.8)	1.33±0.49 (39.9)	1.16±0.40 (36.7)	0.67±0.21 (40.2)
Pestoban	2.5±0.42 (57.7) +	1.5±0.34 (33.3)	1.33±0.5 (39.9)	1.33±0.42 (30.7)	0.83±0.40 (31.2)
Control (Agar)	4.33±0.21 (76.36)	3.83±0.16 (72.47)	4.5±0.34 (81.81)	4.16±0.16 (71.23)	3.5±0.34 (74.84)

Values in parentheses are percentages of snails in zone 3 (in contact with attractant food pellet) compared with snails in zone 1 and 2.

Statistically significant ($p < 0.05$) when two way ANOVA was applied in between different molluscicides (+) and their different concentrations (*).

Abbreviations: SP- Seed powder, BP- Bulb powder, BaP- Bark powder

cooling, the pellets were cut out from the layer with a corer (5 mm diameter).

Assay and Apparatus

The bioassay was performed as reported earlier [14, 16]. The bioassay chamber consists of a clean glass aquarium having a diameter of 30 cm. Each aquarium was divided into four concentric zones; Zone 3 (central zone), zones 2 and 1 (middle zone) and zone 0 (outer zone) had diameters of 13, 18, 24 and 30 cm, respectively. A small annular elevation of 9 mm height and 2.4 cm in diameter was made in the centre of aquarium (Zone 3). Zone 0 had an area of 254 cm² on the periphery of aquarium. The aquaria were then filled with 500 ml of dechlorinated tap water to a height of 8 mm and maintained at 25±1⁰ C. At the start of the assay ten individually marked snails of uniform size were placed on the circumference of zone 0. The distance between two snails was 66 mm. Simultaneously, one of the prepared bait of different molluscicides was added on the small annular elevation in the centre (Zone 3). The location of

each snail was recorded every 15 min for two hours. Six sets of experiments were carried out with ten snails each for every molluscicide used in this study. The mortality of the snails was observed after the test with every 24h up to 144h. Lethal values (LC₅₀), lower and upper confidence limits (LCL and UCL), slope values, t- ratio, 'g' value and heterogeneity factor were calculated using POLO computer programme [17]. One/two-way ANOVA and product moment correlation coefficient was applied between the different data obtained in Tables 1-4 [18]. These experiments were repeated in stored pellets kept for 7, 14, 21 and 28 days under laboratory conditions.

RESULTS

Low attraction (45.5%) of the snails was observed by the Snail Kill compared to plant derived molluscicides in zone 3 at 0.1% concentration in AFP (Table 1). 0.1% AFP containing *A. squamosa* seed powder, *A. sativum*, *P. tuberosa* bulb powder, *A. indica* bark powder,

Table 2. Mean number of snail *L. acuminata* in zone three in contact with the stored attractant food pellets (AFP) containing 0.1% molluscicides.

Molluscicides	TIME OF STORAGE (IN DAYS)				
	0	7	14	21	28
<i>Pestoban</i>	2.5±0.42 (57.7) +	2.16±0.16 (50.0)	1.83±0.16 (61.1)	1.75±0.19 (43.2)	1.16±0.16 (41.2)
<i>Snail Kill</i>	1.67±0.21 (45.5) +	1.67±0.21 (44.2)	1.33±0.21 (36.3)	0.83±0.16 (38.2)	0.67±0.21 (34.5)
<i>A. indica</i> (Ba P)	3.5±0.34 (60.0) +	3.33±0.21 (55.5)	1.83±0.16 (47.7)	1.5±0.22 (40.9)	1.33±0.21 (34.5)
<i>A. sativum</i> (BP)	3.67±0.2 (78.0) +	4.0±0.36 (44.4)	3.16±0.47 (41.2)	2.72±0.51 (40.4)	2.16±0.16 (35.0)
<i>P. tuberosa</i> (BP)	3.0±0.25 (60.0) +	1.83±0.16 (31.3)	1.67±0.21 (35.7)	1.16±0.16 (27.8)	0.83±0.16 (31.2)
<i>A. squamosa</i> (SP)	1.16±0.16 (58.0) +	1.67±0.21 (35.7)	1.33±0.42 (30.7)	1.16±0.16 (27.8)	0.83±0.16 (31.2)
Acetogenin	3.0±0.36 (50.0) +	2.67±0.21 (47.0)	2.0±0.44 (40.0)	1.5±0.34 (33.3)	1.16±0.16 (27.8)
Azadirachtin	1.16±0.16 (46.4) +	0.83±0.16 (31.2)	0.83±0.16 (31.2)	0.67±0.42 (28.6)	0.5±0.22 (25.0)
Allicin	1.4±0.35 (46.7) +	1.33±0.42 (36.3)	1.16±0.16 (34.9)	0.83±0.3 (31.2)	0.83±0.3 (31.2)
Hecogenin	0.67±0.21 (50.3) +	0.67±0.21 (28.6)	0.67±0.21 (28.6)	0.5±0.22 (25.0)	0.33±0.21 (24.0)
Control (Agar)	4.5±0.34 (81.81)	4.66±0.21 (78.23)	5.5±0.16 (74.87)	4.33±0.21 (78.56)	5.33±0.47 (77.68)

Values in parentheses are percentages of snails in zone 3 (in contact with the stored attractant food pellet) compared with snails in zone 1 and 2.

+ Statistically significant ($p < 0.05$) when one way ANOVA was applied in between the number of snails in different storage period of bait formulations.

Abbreviations as in table 1.

Table 3. Toxicity in different bait formulations of molluscicides against the snail *L. acuminata* at different time exposure.

Expo-sure Period	Molluscicides	LC ₅₀ % in AFP	LCL	UCL	Slope value	t-ratio	g-value	Heterogeneity
24h	<i>A. sativum</i>	1.57	1.07	4.38	1.80±0.45	3.92	0.24	0.28
	<i>P. tuberosa</i>	2.07	1.5	5.0	2.09±0.54	3.87	0.25	0.17
	Allicin	1.35	0.84	4.1	1.45±0.33	4.38	0.20	0.32
	Hecogenin	1.54	1.04	6.0	1.9±0.57	3.34	0.34	0.13
	Snail Kill	1.36	1.04	2.56	2.52±0.57	4.38	0.19	0.30
48h	<i>A. sativum</i>	1.53	0.93	11.29	1.12±0.37	3.03	0.41	0.15
	<i>P. tuberosa</i>	1.19	1.00	1.60	2.41±0.49	4.84	0.16	0.15
	Allicin	0.92	0.59	2.59	1.14±0.27	4.15	0.22	0.16
	Hecogenin	0.90	0.74	1.32	2.34±0.51	4.52	0.19	0.16
	Snail Kill	1.30	0.97	2.68	2.07±0.50	4.08	0.23	0.22
72h	<i>A. sativum</i>	0.86	0.61	2.0	1.16±0.34	3.35	0.34	0.22
	<i>P. tuberosa</i>	1.01	0.85	1.33	2.12±0.48	4.36	0.20	0.27
	Allicin	0.32	0.24	0.45	1.34±0.25	5.21	0.14	0.28
	Hecogenin	0.70	0.58	0.93	2.0±0.48	4.33	0.20	0.15
	Snail Kill	0.94	0.76	1.43	2.08±0.46	4.48	0.19	0.16
96h	<i>A. sativum</i>	0.44	0.28	0.61	1.19±0.33	3.55	0.30	0.21
	<i>P. tuberosa</i>	0.76	0.64	0.88	2.61±0.50	5.22	0.14	0.29
	Allicin	0.20	0.14	0.27	1.36±0.25	5.30	0.13	0.21
	Hecogenin	0.55	0.45	0.67	2.15±0.47	4.53	0.18	0.20
	Snail Kill	0.68	0.55	0.91	1.84±0.44	4.19	0.21	0.16
120h	<i>A. sativum</i>	0.28	0.08	0.42	0.99±0.33	2.99	0.42	0.28
	<i>P. tuberosa</i>	0.57	0.46	0.66	3.26±0.56	5.80	0.11	0.36
	Allicin	0.12	0.07	0.15	1.68±0.28	5.99	0.10	0.20
	Hecogenin	0.39	0.27	0.46	2.11±0.47	4.43	0.19	0.20
	Snail Kill	0.41	0.29	0.51	1.89±0.44	4.28	0.20	0.18
144h	<i>A. sativum</i>	0.18	0.07	0.26	1.46±0.35	4.15	0.22	0.34
	<i>P. tuberosa</i>	0.48	0.38	0.55	3.92±0.70	5.55	0.12	0.44
	Allicin	0.09	0.05	0.12	2.03±0.33	6.11	0.10	0.42
	Hecogenin	0.27	0.17	0.34	2.63±0.53	4.95	0.15	0.53
	Snail Kill	0.33	0.25	0.40	2.90±0.50	5.82	0.11	0.31

Product moment correlation showed significant ($p<0.05$); negative correlation in between the exposure period and LC₅₀ of different molluscicides.

Abbreviations as in table 1.

108der, their active components i.e. acetogenin, allicin, 1373 and 4). The crude preparations of plant derived mol-
109hecogenin, azadirachtin and pestoban caused more at-138luscicides and AFP containing pestoban caused signifi-
110traction (58.0%, 78.0%, 60.0%, 60.0% 50.0%, 46.7%, 139cant molluscicidal activity against *L. acuminata* (Table
11150.3%, 46.4% and 57.7%, respectively) than the AFP1403 and 4).

112containing 0.2% to 1.0% of the same molluscicides. The141 The slope values given in Tables 3 and 4 were steep.
113attraction of the snails was significantly ($p<0.05$) re-142Separate estimate of LC₅₀ based on each of the six repli-
114duced with increasing concentration of different mollus-143cates was found to be within 95% confidence limits.
115cicides in AFP. Lowest attraction (13.2%) of snails was144The t- ratio was greater than 1.96 and the heterogeneity
116observed for 1.0% hecogenin containing AFP. There145less than 1.0. The 'g' value was less than 0.5 at all prob-
117was a significant ($p<0.05$) decrease in the number of the146ability levels (90, 95, 99).

118snails attracted by all other 0.1% stored AFP for 7, 14,

11921 and 28 days except *A. sativum* containing AFP (Ta-

120ble2).

121 Molluscicidal activity of different AFP containing148 Higher attraction of the snails towards AFP contain-
122products/compounds against *L. acuminata* followed a149ing 0.1% plant derived molluscicides compared with
123time and dose dependence relationship (Table 3-4).150Snail Kill, appears to be due to the slower release of
124There was a significantly ($p<0.05$) negative correlation151molluscicidal compounds in comparison with synthetic
125between exposure period and LC₅₀ in different mollus-152ones. Higher concentration of plant derived mollus-
126cicides. AFP containing bioactive components of differ-153cicides and their active components in AFP caused less
127ent plants were more toxic (acetogenin 24h LC₅₀-1.02%154attraction than corresponding concentration of Snail
128in AFP; 144h LC₅₀-0.12% in AFP) than synthetic ones.155Kill. It indicates that when higher titer of active compo-
129The molluscicidal activity of garlic crude bulb powder156nents of plant derived molluscicides was used in AFP,
130(144h LC₅₀-0.18% in AFP) was higher than Snail Kill157snails were less attracted. Higher concentration (1.0%)
131(144h LC₅₀-0.33% in AFP). Allicin was more toxic158of allicin, hecogenin and azadirachtin [19-21] attracted
132(144h LC₅₀-0.09% in AFP) than the crude bulb powder159less snails than Snail Kill. There was a significant de-
133of *A. sativum* (144h LC₅₀-0.18% in AFP). The bioactive160crease in the attraction of *L. acuminata* towards AFP
134components hecogenin, acetogenin and azadirachtin161containing molluscicides compared with AFP alone
135were more toxic (24h LC₅₀- 1.54%, 1.35% and 1.10% in162with a significant variation in mean number of snails in
136AFP, respectively) than their crude preparations (Table163zone 3 containing different concentrations of mollus-

DISCUSSION

Table 4. Toxicity in different bait formulations of molluscicides against the snail *L. acuminata* at different time exposure.

Exposure Period	Molluscicides	LC ₅₀ % in AFP	LCL	UCL	Slope value	t-ratio	g-value	Heterogeneity
24h	<i>A. indica</i> BaP	1.48	0.86	6.98	1.24±0.33	3.76	0.27	0.18
	<i>A. squamosa</i>	1.53	1.23	2.57	3.18±0.74	4.27	0.21	0.26
	Azadirachtin	1.10	0.86	1.99	2.26±0.54	4.15	0.22	0.12
	Acetogenin	1.02	0.72	2.16	1.68±0.36	4.62	0.18	0.29
	Pestoban	2.82	2.24	5.71	3.14±0.86	3.62	0.29	0.23
48h	<i>A. indica</i> BaP	1.01	0.62	3.90	1.03±0.28	3.61	0.29	0.17
	<i>A. squamosa</i>	1.25	1.08	1.88	2.75±0.65	4.24	0.21	0.14
	Azadirachtin	1.03	0.78	2.14	1.82±0.50	3.64	0.28	0.13
	Acetogenin	0.78	0.54	1.75	1.23±0.29	4.22	0.21	0.19
	Pestoban	2.46	1.98	4.94	2.43±0.71	3.40	0.33	0.23
72h	<i>A. indica</i> BaP	0.53	0.37	1.09	1.05±0.27	3.88	0.25	0.15
	<i>A. squamosa</i>	0.98	0.83	1.32	2.38±0.60	3.95	0.24	0.11
	Azadirachtin	0.73	0.59	1.11	1.78±0.47	3.75	0.27	0.13
	Acetogenin	0.39	0.29	0.58	1.23±0.27	4.57	0.18	0.21
	Pestoban	1.73	1.48	2.27	2.44±0.66	4.14	0.28	0.21
96h	<i>A. indica</i> BaP	0.25	0.17	0.34	1.23±0.26	4.66	0.17	0.22
	<i>A. squamosa</i>	0.63	0.50	0.73	2.71±0.61	4.45	0.19	0.11
	Azadirachtin	0.41	0.25	0.52	1.60±0.46	3.45	0.32	0.14
	Acetogenin	0.32	0.24	0.42	1.47±0.27	5.42	0.13	0.52
	Pestoban	1.33	0.99	1.60	2.07±0.63	3.26	0.36	0.28
120h	<i>A. indica</i> BaP	0.15	0.70	0.23	1.10±0.26	4.20	0.21	0.23
	<i>A. squamosa</i>	0.53	0.45	0.59	4.75±0.74	6.37	0.09	0.37
	Azadirachtin	-	-	-	-	-	-	-
	Acetogenin	0.19	0.10	0.27	1.09±0.26	4.18	0.22	0.49
	Pestoban	1.06	0.75	1.25	2.58±0.64	4.00	0.24	0.28
144h	<i>A. indica</i> BaP	0.09	0.03	0.14	1.27±0.27	4.57	0.84	0.36
	<i>A. squamosa</i>	0.48	0.40	0.54	5.32±0.88	6.04	0.10	0.37
	Azadirachtin	-	-	-	-	-	-	-
	Acetogenin	0.12	0.06	0.16	1.40±0.27	5.10	0.14	0.55
	Pestoban	0.96	0.75	1.10	3.65±0.68	5.32	0.13	0.86

Product moment correlation showed significant ($p < 0.05$); negative correlation between the exposure period and LC₅₀ of different molluscicides.

Abbreviations as in table 1.

cicides after two hours of exposure. AFP containing from the seeds of *A. squamosa* is higher than other plant acetogenin attracted more snails at lower concentrations derived molluscicides and Snail Kill (24h LC₅₀- 1.36% than *A. squamosa* seed powder. It indicates that *A.* in AFP). Seeds of *A. squamosa* were used to kill human *squamosa* seed powder, instead of acetogenin contains lice [22] and their organic extracts have been reported to some other compounds which reduce the attraction of possess insecticidal activity [23-24]. Molluscicidal ac-snails towards AFP. In contrast AFP containing *A. sati*-tivity of *A. indica* bark powder (24h LC₅₀ 1.48% in *vum* and *P. tuberosa* bulb powder and *A. indica* bark AFP) is lower than the Snail Kill. However, its active powder attracted more snails than their pure compounds component azadirachtin (24h LC₅₀ 1.10% in AFP) is viz. allicin, hecogenin and azadirachtin. It seems that more toxic than Snail Kill. Toxicity of AFP containing these plant derived molluscicides either contain some azadirachtin was effective only up to 96h. It indicates other compounds which attract the snails or the concen- that it is less stable in water or it is metabolized in snail trations of active molluscicidal components are less in body [20]. AFP containing hecogenin in AFP is 1.5 time crude preparations. The storage of attractant food pellets more toxic than the crude bulb powder of *P. tuberosa*. It for up to 28 days caused significant decrease in the at- has been reported that treatment of *P. tuberosa* bulb traction of snails. Thus, it seems logical to assume that powder and hecogenin caused significant reduction in AFPs containing plant derived molluscicides are less the reproduction of the *L. acuminata* [20]. It has been effective in attracting snails, when stored up to 28 days. reported that the allicin caused an uncompetitive inhibi- However, toxicity of these AFP containing mollus- tion of acetylcholinesterase and competitive inhibition cicides was time and dose dependent as evident from 209 of lactic dehydrogenase and alkaline phosphatase activ- the negative correlation between LC₅₀ in different mol- tivity in the nervous tissue of *L. acuminata* [25]. The toxic- luscicides and exposure period.

211ity of molluscicide Pestoban is due to the presence of Treatment of bulb powder of *A. sativum* and *P. tube*- *Cedrus deodara*, *A. indica* and *Embelia ribes* in liquid *rosa*, seed powder of *A. squamosa*, bark powder of *A.* concentrate form [26]. The toxicity of Pestoban to the *indica* and their active components such as allicin, snail is lower (24h LC₅₀ 2.82% in AFP) than the syn- hecogenin, acetogenin and azadirachtin in aquatic envi- thetic molluscicide Snail Kill (24h LC₅₀ 1.36% in AFP). ronment are highly toxic to *L. acuminata* [19-21]. Tox- The steep slope values indicate that a small increase 217 in the concentration in different molluscicides cause a

218 significant mortality in the snail. t- ratio value greater
 219 than 1.96 indicates that the regression is significant.
 220 Values of heterogeneity less than 1.0 denote that in the
 221 replicates the concentration response line would fall
 222 within 95% confidence limit and thus the model fit the
 223 data adequately. The value of 'g' is less than 0.5 indi-
 224 cates the index of significance of potency estimation.
 225 Use of plant derived molluscicides in aquatic envi-
 226 ronments requires large amounts of molluscicides for
 227 effective control of snails. Using attractant food pellets
 228 like this study will be beneficial since it requires small
 229 quantities of molluscicides while killing the target pest
 230 specifically. The present study shows that the use of
 231 AFPs containing plant derived molluscicides is very
 232 effective in killing the snail *L. acuminata*. Use of these
 233 plant derived molluscicides inside the baits (Attractant
 234 food pellets) are ecologically sound, target specific and
 235 economic.

236 REFERENCES

2371. Singh O, Agarwal RA. 1981. Toxicity of certain pesticides to
 238 two economic species of snails in northern India. *J Econ Ento-*
 239 *mol* 74:568-571.
2402. Agarwal RA, Singh DK. 1988. Harmful gastropods and their
 241 control. *Acta Hydrochim Hydrobiol* 16:113-138.
2423. Godan D. 1983. "Pests slugs and snails. Biology and control"
 243 (ed., Dora Godan) translated by Sheila Grouber, Springer Verlag,
 244 Berlin Heidelberg New York .
2454. Marston A, Hosttetmann K. 1985. Plant molluscicides. *Phyto-*
 246 *chemistry* 24:639-652.
2475. Marston A, Hosttetmann K. 1987. Antifungal, molluscicidal and
 248 cytotoxic compounds from plants used in traditional medicines.
 249 In: "Biologically Active Natural Products" (Ed. Hosttetmann, K
 250 and Lea, PJ). Clarendon Press Oxford 65-83.
2516. Ndamba J. 1995. Response of the molluscicidal berry plant
 252 *Phytolacca dodecandra* to different climatic and edaphic condi-
 253 tions. *Trop Agric* 72:135-140.
2547. Singh A, Singh DK, Mishra TN and Agarwal RA. 1996a. Mol-
 255 luscicides of plant origin. *Biol Agri Horti* 13:205-252.
2568. Singh K, Singh DK. 2000. Effect of different combinations of
 257 MGK-264 and piperonyl butoxide with plant derived mollus-
 258 cicides on snail reproduction. *Arch Environ Contam Toxicol*
 259 38:182-190.
2609. MacInnis AJ, Bethal WM, Cornford EM. 1974. Identification of
 261 chemicals of snail origin that attract *Schistosoma mansoni*
 262 miracidia. *Nature* 248:361-363.
26310. Sterry PR, Thomas JD, Patience, RL. 1985. Changes in the con-
 264 centrations of short-chain carboxylic acids and gases during de-
 265 composition of the aquatic macrophytes *Lemna paucicostata* and
 266 *Ceratophyllum demersum*. *Freshw Biol* 15:139-153.
- Thomas JD. 1982. Chemical ecology of the snail hosts of schis-
 33:81-91.
- Thomas JD, Kowalczyk C, Somsundaram B. 1989. The bio-
 chemical ecology of *Biomphalaria glabrata*, a snail host of
Schistosoma mansoni; short chain carboxylic and amino acids as
 phagostimulants. *Comp Biochem Physiol* 93A:899-911.
- Kpikpi JE K, Thomas JD. 1992. A study of sugar chemorecep-
 tion niches of two bulinid snail hosts of schistosomiasis. *Ann*
Trop Med Parasitol 86:181-198.
- Tiwari F, Singh DK. 2004a. Behavioural responses of the snail
Lymnaea acuminata to carbohydrates in snail attractant pellets.
Naturewissenschaften 91:378-380.
- Madsen H. 1992. A comparative study on the food locating
 ability of *Helisoma duryi*, *Biomphalaria camerunensis* and *Bu-*
linus truncatus (Pulmonata:Planorbidae). *J Appl Ecol* 29:70-78.
- Tiwari F, Singh DK. 2004b. Attraction to amino acids by *Lym-*
naea acuminata, the snail host of *Fasciola* species. *Braz J Med*
Biol Res 37:587-590 .
- Russel RM, Robertson JL, Savin NE. 1977. POLO: A new com-
 puter programme for probit log analysis. *Bull Entomol Soc*
Amer 23:209-213.
- Sokal RR, Rohlf FJ. 1973. Introduction to Biostatistics. W H
 Freeman, San Francisco, 368pp.
- Singh VK, Singh DK (1995) Characterization of allacin as a mol-
 luscicidal agent in *Allium sativum*. *Biol Agric Horti* 12:119-
 131.
- Singh K, Singh A, Singh DK. 1996b. Molluscicidal activity of
 neem (*Azadirachta indica* A. Juss). *J of Ethnopharmacol* 52:35-
 40.
- Singh K, Singh VK, Singh DK. 1999. Effect of *Polianthes tube-*
rosa (Amarillidaceae) on the reproduction and biochemical pa-
 rameters in the ovotestis of the snail *Lymnaea acuminata* (Mol-
 lusca:Pulmonata). *Acta Hydrochim Hydrobiol* 27:32-37
- Reyes FR, Santos AC. 1931. Isolation of Anonaine from *Annona*
squamosa Linn. *Philippines J of Sci* 44:409-410.
- Chattoraj AN, Tiwari SC. 1965. A note on the insecticidal prop-
 erty of *Annona squamosa* (Annonaceae). *Natl Acad Sci India*
Proc Sec B (Biol Sci) 35:351-353.
- Mukerja TD, Govind R. 1958. Indigenous insecticidal plants:II
Annona squamosa. *J Indian Res (India)* 17C:9-15.
- Singh VK, Singh DK. 1996. Molluscicidal activity of pre- and
 post-harvest (garlic) *Allium sativum*. *Biol Agric Horti* 12:311-
 318.
- Singh K, Singh A, Singh DK. 1995. Molluscicidal activity of
 different combinations of the plant products used in the mollus-
 cicide Pestoban. *Biol Agri Horti* 12:253-261.

314 CURRENT AUTHOR ADDRESSES

- Farindra Tiwari, D.D.U. Gorakhpur University, Gorakhpur, India.
 Dinesh Kumar Singh, D.D.U. Gorakhpur University, Gorakhpur,
 India. E-mail: dksing_gpu@yahoo.co.in (Corresponding author)