

1735-2657/07/62-159-163 **IRANIAN JOURNAL OF PHARMACOLOGY & THERAPEUTICS** Copyright © 2006 by Razi Institute for Drug Research (RIDR) IJPT 6:159-163, 2007

RESEARCH ARTICLE



# 2Effect of Chromatographic Fractions of Ethanolic Extract of Crotalaria Juncea (L.) Seeds on Ovarian Follicular Kinetics and Estrous Cycle in Albino Rats

5VIJAYKUMAR B. MALASHETTY and SARASWATI B. PATIL

6 For author affiliations, see end of text.

7 Received November 7, 2006; Revised May 5, 2007; Accepted August 14, 2007

This paper is available online at http://ijpt.iums.ac.ir

# **ABSTRACT**

I o In spite of considerable development in contraceptive technology, search for female antifertility agent in uplants continues to be a potential area of investigation. The ethanol extract of Crotalaria juncea seeds 12 which showed promising antiovulatory activity in female albino rats was examined for the isolation of its 13 active fractions. Two fractions were obtained using Thin Layer Chromatography (TLC) of the extract. Both 14 fractions were subjected for testing their anti-ovulation activity and estrous cycle in rats. After preliminary 15 trials, the fraction I (200mg/kg body weights) showed maximum antiovulatory activity when administered 16 orally to the rats for 30 days. Decreased number of healthy follicles (Class I – ClassVI) and corpora lutea 17 and increased number of regressing follicles (Stage IA, Stage IB, Stage IIA, Stage IIB) were observed in 18the ovary after 30 days treatment. The treatment caused an increase in the cholesterol level and 19 acid/alkaline phosphatase activity and a decrease in protein and glycogen contents of the ovary. Estrous 20 cycle was affected as a significant increase in estrus and metaestrus phases and a decrease in diestrus 21 and proestrus phases in the treated groups during experimental period of 30 days were observed. These 22 results suggest that a fraction of ethanolic extract of crotalaria juncea might be used as a contraceptive in 23 the females.

24 Keywords: Crotalaria juncea; Antiovulatory; Estrous cycle; Antifertility; Rat

26hemp, belongs to the family Papilionaceae. The medici- 46Gulbarga University, Gulbarga (HGUG No. 141), India; 27 nal properties have been described in Ayurveda, by Su- 47 where voucher specimens were deposited. 28 shruta as well as in ancient books like Sarangadhara and 29Bhavaprakasha. In Ayurveda, the leaves are used as an 30emetic, laxative, abortifacient and analgesic, and for 31 treating diarrhea, dysentery and bleeding disorders. The seeds are used as abortifacient and in the treatment of 33 impetigo, psoriasis and as an emmenogogue [1-2]. Pre-34 vious works on the ethanol extract of C. juncea demon-35 strated their anti-implantation [3], antiovulatory [4] and 36anti-spermatogenic activities [5-7] in rats and mice. In 53sorbent. The extract was loaded on the preparative plates 37the present work, we have undertaken the investigation 54 and developed with solvent system benzene: methanol 380f the antiovulatory activity of chromatographic frac- 55(80:20). Two major bands were observed by Iodine vapors. 39tions of crude ethanol extract of C. juncea to elucidate 56The compounds having high retention power (Rf) was des-40its active ingredient.

### MATERIALS AND METHODS

### 42 Plant material

44 local source during October and November 2003, and 62 making required doses in Tween-80 (1%).

Crotalaria juncea (L.), commonly called as Sunn 45 authenticated at the Herbarium, Department of Botany,

# 48 Preparation of test material

Fresh dried seeds of C. juncea were powdered and 50 soxhleted with ethanol (95%) for 24 hours. The filtrate 51 was dried under reduced pressure and chromatographed 52by thin layer chromatography over silica gel 'G' as ab-57 ignated as fraction I and the compounds with low Rf value 58 was designated as fraction II. The fraction I of the ethanol 59 extract yielded brownish gummy material and the fraction 60 II yielded yellow gummy material when the silica gel was The fresh seeds of C. juncea were obtained from a 61 washed with methanol. Both the fractions were used after



Fig 1. C. S. of liver of control mice showing normal well organized hepatic chords with parenchymatous hepatocytes and blood vessels (×125).

#### 63 Acute toxicity studies

The acute toxicity study was performed according to 65the method described by Kattan and colleagues [8]. 66 Adult albino mice of either sex were divided into five 67 groups containing ten animals in each. Graded doses 68(100 and 200 mg/kg b.w.) of fraction I and II of ethanol 69 extract of C. juncea seeds in tween-80(1%) were admin-70 istered orally by mean of intragastric catheter to healthy 71 adult mice (30-35g). Following administration of the 72 extracts, the animals were observed continuously for 2 73horrs and then frequently for further 4 hours. Mortality 74 was recorded. After 7<sup>th</sup> day, the detection of hematotox-75 icity, haemoglobin concentration, total RBC and WBC 76 counts were examined. The histological changes in the<sup>1</sup> 77 liver were also studied.

### 78 Animals

Sexually matured, healthy, colony-bred virgin fe-117 ing an electronic balance. somale rats of Wistar strain (Rattus norvegicus), aged 3118 Histopathological studies 81 months and weighing 150-200g were used for the ex-82periments. The rats were housed in polypropylene cages<sup>119</sup> 83 measuring 12"x10"x8", under well-ventilated animal 120 Bouin's fluid, embedded in paraffin wax, sectioned at 84 house conditions (ambient temperature: 28-31°C, pho-1215µm, stained with haematoxylin-eosin for follicular 85 toperiod: 12h natural light and 12h dark; relative humid-122 studies. 86ity: 50-55%). The rats were given pelleted feed (Hindu-87 stan Lever Ltd., India) and tap water ad labium. They 88 were maintained as per the principles of Laboratory 124 89Animal Care [9]. The experimental protocol was ap-125 classify follicles using established methods [10-11] as 90 proved by the Institutional Animal Ethics Committee.

# 91 Experimental design

93 of six animals in each group.

- 94 Group I: Control, received 0.2ml Tween-80 (1%)
- 95Group II: Received 100mg /kg b.w. fraction I in 0.2ml132Class V: Large sized antral follicles (LSAF) (431-96 tween - 80 (1%) 133**490µm**)
- 97Group III: Received 200mg /kg b.w. fraction I in 0.2ml134Class VI: Graafian follicles (GF) (>491µm) Follicles under regression were classified depending 98 tween - 80 (1%)
- 99Group IV: Received 100mg /kg b.w. fraction II in 0.2ml136on the degree of regression as: 100 tween-80 (1%)

137 Stage IA: Pyknosis in some granulose cells.

Fig 2. C. S. of liver of crude ethanol extract of C. juncea treated mice showing normal histological picture similar to that of control ( $\times 125$ ).

101 Group V: Received 200mg /kg b.w. fraction II in 0.2ml tween-80 (1%)

All the above treatments were given orally by using of intragastric catheter for 30 days to cover six regular estrous cycles. The treatment was started from estrous phase (by observing cornified cells in the vaginal smear), as the ovarian activities change markedly from one phase to another phase of estrous cycle. The treatment was given orally everyday between 10.00 and 11.00 am. The stages of estrous cycle were recorded daily by observing vaginal smears.

# Autopsy and organ weight

On 31<sup>st</sup> day, 24 hours after last dosing, all the ani-14mals were weighed and sacrificed by cervical disloca-115tion. The ovaries were dissected out immediately and 116 separated out from the adherent tissue and weighed us-

The ovary from one side of each animal was fixed in

#### 123 Morphometric analysis

Follicular diameter and morphologies were used to 126 follows:

127 Class I: Small preantral follicles (SPAF) (<90µm)

- 128Class II: Large preantral follicles (LPAF) (91-260µm)
- The animals were divided into 5 groups consisting 129 Class III: Small antral follicles (SAF) (261-350µm)
  - 130 Class IV: Medium sized antral follicles (MSAF) (351-131430µm)

# Malashetty and Patil

### Effect of Fractions of Ethanolic Extract of Crotalaria Juncea (L.)

ijpt.iums.ac.ir | 161



Fig 3. Control rat with normal ovarian follicle with antrum, corona radiata, cumulus oophorus and granulosa membrane ( $\times 400$ ).



Fig 4. Rat treated with fraction I of ethanol extract of C juncea seeds showing greater follicular atresia (x 400).

138 Stage IB: Degenerative changes in the entire granulose 173 Changes in estrous cycle 139**layer**.

140Stage IIA: Oocytes with pyknotic nuclei, blocked meio-<sup>174</sup>

143Stage IIB: Characterized by oocytes floating in the an-144 trum with few pyknotic bodies

Morphometric studies of the ovary were calculated 146 by using stage and ocular micrometers.

### 147 Biochemical studies

Ovary from the other side of each animal was used 149 for biochemical estimations like protein [12], glycogen 150[13], cholesterol [14], acid and alkaline phosphatase<sup>184</sup> creased the ovarian weight significantly (p<0.001); 151[15].

#### 152 Data processing

The statistical analysis was done to determine sig-154nificant difference of values between treated and control<sup>188</sup> 155 groups using Students t- test.

# RESULTS

# 157 Acute toxicity studies

159 body weight of animals was observed. The mortality 196 nificant (p < 0.01) in increasing their enzyme activities 160 was nil. Blood variables, i.e., RBC, WBC and haemo-197(Table 5). 161 globin were within the normal range. There was no his-162tological change in the liver of treated mice with frac-163 tion I and II of ethanol extract of C. juncea seeds when 199

164 compared to that of control (Table 1 & 2; Figs 1 & 2).

## 165 Changes in body weight

167 els to female rats revealed no change in the body 204 cles, but it was significant with class III (p < 0.001), 168 weight. The treated rats were healthy and maintained 205 class V (p<0.001) and class VI (p<0.01) follicles (Table 169 normal growth rate throughout the experiment. Though 2066; Figs 3 & 4).

170 the body weight of the animals in treated groups showed 207 171 slight fluctuation when compared to control group, this 208 caused significant (p < 0.001) increase in regressing fol-172was negligible (Table 3).

Administration of fraction I at both doses levels sig-141 sis in metaphase I (pseudomaturation) and degenerating  $\frac{1}{176}(p<0.001)$  and decreased the diestrous and proestrous 177 phases. The administration of fraction II increased the 78estrous phase nonsignificantly but metestrous phase significantly (p < 0.001) and decreased proestrus phase 80nonsignificantly (Table 4).

### Changes in the ovary

## 182 Gravimetric changes

Administration of fraction I at both dose levels de-185 whereas nonsignificant reduction was obtained with 186 both doses of fraction II (Table 5).

### 187 Biochemical changes

Protein content was reduced significantly (p < 0.001)189 with both doses of fraction I and II. Glycogen content 190 was reduced and cholesterol content was increased sig-191 nificantly (p < 0.001) with both low and high dose of 192 fraction I and high dose of fraction II. Acid and alkaline 193 phosphatase activity showed significant (p < 0.001) in-194 crease with both the doses of fraction I and high dose of In acute toxicity tests, no change in the behaviour or 195 fraction II. However, fraction II at low dose is less sig-

#### 98 Changes in follicular kinetics

Healthy follicles: Fraction I and II at both dose lev-200els decreased the number of follicles of class I to class 201 V significantly (p<0.001). The class VI or Graafian fol-202licles were totally absent. Fraction II administration at Administration of fraction I and II at both dose lev-203low dose level reduced all the classes of healthy folli-

> Regressing follicles: Fraction I at low dose level 209licles. Fraction II at low dose level increased the number

2100f stage IB (p < 0.001) and stage IIB (p < 0.05) regressing 267 crude extracts of C. juncea, the rats have not lost the 211 follicles. Fraction II at high dose level caused signifi-268 rhythmic cyclicity. These findings on the estrous cycle 212cant reduction in stage IA (p < 0.001), stage IB<sub>269</sub>clearly corroborate the potent estrogenic nature of the 213(p < 0.001), stage IIA (p < 0.05) and stage IIB (p < 0.001)270extract.

214 regressing follicles (Table 7; Figs. 3 & 4).

# 215 Changes in corpora lutea

217 significantly (p<0.001) with both doses of fraction I and 275 phins due to the treatment with C. juncea seed extracts. 218 high dose of fraction II (Table 6).

DISCUSSION

221 estrous cycle gives a reasonable index of the ovarian 282 ble for follicular regression rather than maturation. The 222 activity and its hormonal synthesis of estrogen and pro-283 reduced number or total absence of corpora lutea in ex-223 gesterone. The levels of these hormones are controlled 284 tracts-treated ovaries of rats indicates the blockade of 224by hypothalamic releasing hormones and pituitary go-285 ovulation which depicts the antiovulatory property of the 225 nadotrophins [16]. A feedback mechanism also operates 286 extracts. 226 where the pituitary gonadotrophins secretion in turn is  $\frac{1}{287}$ 227 controlled by estrogen and progesterone. The cornifica-288 utilization of cholesterol for steroidogenesis in two cell 228tion in the vaginal epithelial cells is mainly due to high 229 levels of estrogens secreted by the ovarian matured fol-230 licles. It is also known that exogenous administration of 231 estrogen consistently stimulates the proliferation of the 292 rats. Protein is considered to be the building material 232vaginal epithelium in adult spayed animals [17-18]. The 233 basic functional unit of reproduction within the ovary is 234 the follicle [19]. Follicles start to grow at all times and 235 as they develop, they produce large number of granu-236 losa and thecal cells. The conversion of follicles to 237 atretic state is functional rather than a degenerative 238 process and is considered to be integral part of ovarian 239 function [20-10]. Most of the follicles undergo atresia 240 and very few mature to ovulate among the new crop of 301 in reproduction. It is involved in providing energy to 241 recruited follicles during every cycle. After the early 302 various processes like ovulation, transportation and sur-242 stage of gonadotrophin independence, the entire process<sub>303</sub> vival of eggs and implantation. All these changes are 243 of follicle growth becomes dependent on the continuous 244 presence of gonadotrophins [21-22]. Evans et al.  $[23]_{305}$  gen content in extracts of *C. juncea* seeds treated rats 245 have shown that the ovarian androgen and inhibin secre-306 may be due to lowered steroidogenesis, which attributed 246 tion by follicles may play an important part in the regu-307 to nonavailability of gonadotrophins.  $^{247}$ lation of FSH secretion and follicular dynamics. The  $^{308}_{308}$ 248 integral role in the control of ovarian function is played 309 healthy follicles and corpora lutea but increased the 249by the hypothalamo-pituitary unit. Functioning in a co-310 number of regressing follicles. This indicates nonavail-250 ordinated manner with appropriate signals provided by 311 ability of gonadotrophins for follicular development and 251 ovary via pituitary gland is responsible for the synthesis  $_{312}$  ovulation. Hence, fraction I of ethanol extract of C. 252 and storage of gonadotrophins LH and FSH. These gly-<sub>313juncea</sub> has strong antiovulatory property. 253 coprotein hormones in turn play a key role as regulators 254 of folliculogenesis.

The data obtained in the present study reveal that the 314 REFERENCES 256 control rats exhibited regular estrous cycle of 4-5 days. 3151. 257Treatment with extracts of Crotalaria juncea seeds 316 258 caused a significant increase in the estrous and 3172. 259 metestrous with concomitant decrease in the duration of  $_{318}$ 260 diestrous and proestrous phases. Similar results have 3193. 261 been obtained with Hibiscus rosa sinensis [24] and 320 262 Momordica charantia [25] in mice and rats respectively.<sup>321</sup> 263 This may be attributed to the fact that the increased es-<sup>322</sup> 264 trogen production at regular intervals which is influ-3234. 265 enced by the crude extracts of C. juncea is responsible  $_{325}^{324}$ 266 for vaginal cornification. Inspite of the influence of 326

In the present study, the decrease in the number of 272 healthy follicles from class I to class VI and increase in 273 the number of regressing follicles and atretic follicles The number of corpora lutea were reduced highly274 attributes to the non-availability of pituitary gonadotro-

276 The recruitment of SPAF (class I) from primary follicles 277 depends on availability of FSH and further follicu-278 logenesis from SPAF (class I) to GF (class VI) requires 279 both FSH and LH [26-29]. The observed estrogenic na-280 ture of the extracts might have brought the inhibition in Cyclic changes in the vaginal smear observed in the 281 the gonadotrophins secretion and release that is responsi-

> Pituitary FSH, LH and prolactin are essential for 289 compartments of theca and granulosa cells in the ovary. The nonavailability of these gonadotrophins increases the cholesterol depot in the ovary of extracts-treated 3 and is involved in the alteration of almost every physiological function. In the present study the low protein content of the ovary indicates the retarded ovarian growth. It is well understood that FSH is essential for protein synthesis in gonads [30]. The blockade of pituitary FSH releases in extracts-treated rats might have resulted in low protein content.

> The presence of glycogen plays very important role 304 hormone-dependent [31]. The decreased ovarian glyco-

In conclusion, the fraction I reduced the number of

Kirtikar KR, Basu AD. In: Indian Medicinal Plants. Vol. I. Lalit Mohan Basu, Allahabad. 1935; p 792.

Wealth of India. Raw Materials.Vol. IV F-G, Published by C. S. I. R., New Delhi. 1952.

Vijaykumar BM, Sharanabasappa A, Saraswati BP. Post-coital antiimplantation and pregnancy interruption potency of the seed extracts of Crotalaria juncea. Orient Pharm Exp Med 2004; 4: 70-6

Vijaykumar BM, Sangamma I, Sharanabasappa A, Saraswati BP. Effect of Crotalaria juncea seed extracts on the estrous cycle and ovarian activity in albino mice. Orient Pharm Exp Med 2004; 4: 77-81.

# FICLE IN

37221

#### Effect of Fractions of Ethanolic Extract of Crotalaria Juncea (L.)

- 3275. Vijaykumar BM, Sangamma I, Sharanabasappa A, Saraswati 37020. BP. Antifertility activity of various extracts of Crotalaria juncea371
- Linn. seeds in male mice. Philip J Sci 2003; 132: 39-46.
- 3306 Vijaykumar BM, Sangamma I, Saharanabasappa A, Saraswati 373
- BP. Antispermatogenic and hormonal effect of Crotalaria juncea<sub>37422</sub>. Linn. seed extracts in male mice. Asian J Androl 2004; 6: 67-71. 375
- Vijaykumar BM, Saraswati BP. Spermatogenic index and hor-37623. 3337 monal profile in the rats received chromatographic fractions of 377ethanol extract of Crotalaria juncea L. seeds. Orient Pharm Exp<sub>378</sub>
- Med 2006: 6: 86-95. Kattan GFE, Goudgaon NM, Ilksvo N, Huang JT, Watanabe380 3378
- KA, Sommadossi JP, Schinazi RF. Synthesis and biological<sub>38124</sub>. 5-O-Carboranyl-1-(2-deoxy-2-fluoro-β-D-382 properties of arabinofuranosy). J Med Chem 1994; 37: 2583.
- Hariharan S. Laboratory animals information service center<sub>38425</sub>. 3419. news ICMR, Hyderabad. 1980.
- 34310. Hirshfield AN. Development of follicles in the mammalian 386 ovary. Int Rev Cytol 1991; 124: 43-100. 38726.
- 34511. Sanjay VS, Joshi BN. Melatonin and exposure to constant388 light/darkness affects ovarian follicular kinetics and estrous cy-389 cle in Indian desert gerbil Meriones hurrianae (Jerdon). Gen<sub>39027</sub>. Comp Endocrinol 1997; 180: 352-7.
- Lowry OHN, Rosenbrough J, Farr, NL, Randall RJ. Protein<sup>392</sup> 34912 measurement with folic phenol reagent. J Biol Chem 1951; 193:393 265-75. 39428.
- Carrol NV, Langelly RW, Row RH. Glycogen determination in 395 35213. liver and muscle by use of anthrone reagent. J Biol Chem 1956;396 20: 583-93.
- Peters JP, Vanslyke DD. Quantitative Clinical Chemistry, Vol. I. 39829. 35514. Williams and Wilkins, Baltimore. 1946.
- Bessey OA, Lowry OH, Brick NJ. A method for rapid determi-35715. nation of acid and alkaline phosphatase with 5 cu. mm. of serum.40130. 402 J Biol Chem 1946: 164: 321.
- Lerner LJ. The biology of non-steroidal antifertility. In: Contra-40331. 36016. ception, chemical control of fertility. ed. by Lednicer D, Marcel404 Derker Inc, New Yark. 1969; p. 161.
- Mandel AM. Cyclical changes in the vaginal smear of adult405 CURRENT AUTHOR ADDRESSES 36317. ovariectomized rats. J Exp Biol 1951; 28: 585-92.
- Boettiger EG. Changes in the glycogen and water contents of the 36518. rat uterus. J Cell Comp Physiol 1946; 27: 9-13.
- Hsuch AJW, Adashi EY, Jones PBC, Welsh THJ. Hormonal<sup>408</sup>mail:viju\_yes@rediffmail.com, (Corresponding author) 367**19**. regulation of the differentiation of cultured ovarian granulosa409
- cells. Endocrine Rev 1984; 5: 76-127.

- Zuckerman L, Weir BJ. The ovary-1. 2nd ed. General aspects. Academic Press. Inc. New York and London. 1977: 277-279.
- Hodgon GD. Neuroendocrinology of the normal menstrual cycle. J Reprod Med 1989; 34: 68-75.
- Scott RT, Anderson GD. The ovarian follicle: life cycle of the pelvic clock. Obstet Gynecol 1990; 33: 551-61.
- Evans ACO, Komar CM, Wandji SA, Fortune JE. Changes in androgen evretion and lutenizing hormone pulse amplitude are associated with the ecruitment and growth of ovarian follicles during the luteal phase of the ovine estrous cycle. Biol Reprod 1997; 57: 394-401.
- Murthy DRK, Reddy CM, Patil SB. Effect of benzene extract of H. rosa inensis on the estrous cycle and ovarian activity in albino mice. Biol harm Bull 1997; 20: 756-8.
- Sharanabasappa A, Vijaykumar B, Patil SB. Effect of Momordica harantia seed extracts on ovarian and uterine activities in albino rats. Harmceutic Biol 2002; 40: 501-7.
- Richards JS. Maturation of ovarian follicles: actions and interactions of ituitary and ovarian hormone on follicular cell differentiation. Physiol Rev 980; 60: 51-89.
- Young EL, Biard DT, Hillier SG. Melatonin gonadotrophin stimulated rowth and differentiation on human granulosa cells by 31-51 -onophosphate: One molecule, two genes. Clinic Endocrinol 1992; 73: 51-5.
- Wang X, Greenwald GS. Hypophysectomy of the cyclic mouse. I. Effects on folliculogenesis, oocyte growth, follicle stimulating hormone and human chorionic gonadotrophin receptors. Biol Reprod 1993; 48: 585-94.
- Greenwald GS, Roy SK. Follicular development and its control. In: The Physiology of Reproduction, second edition, Knobil E, Neill JD (Eds), Raven Press Ltd, New York, 1994; 629-724.
- Means Ar. Biochemical effects of follicle stimulating hormone on the testis. Endocrinology 1975; 5: 203-18.
- Walaas O. Effect of oestrogens on the glycogen contents of the rat uterus. Acta Endocrinol 1952; 10: 175-92.

Vijaykumar B. Malashetty, Dept. of MRDG, Indian 407 Institute of Science, Bangalore- 560 012, India. E-

Saraswati B. Patil, Department of Zoology, Gul-410barga University, Gulbarga-585 106, India