

# Protective Effect of *Ocimum sanctum* on Disposition Kinetics of Sulphadimidine in Lead-Treated Cockerel

A. PRAKASH, S. P. SINGH, R. VARMA, G. K. CHOUDHARY and R. SAJAN

For author affiliations, see end of text.

Received February 6, 2009; Revised August 29, 2009; Accepted September 5, 2009

This paper is available online at <http://ijpt.iuums.ac.ir>

## ABSTRACT

This study was carried out to evaluate the protective effect of *Ocimum sanctum* (tulsi) on lead-induced alteration in disposition kinetics of sulphadimidine (SD) in cockerels. Twenty-one, four-week male white leghorn (WLH) chicks were randomly divided into groups I, II and III (7 in each) and were fed at dietary with 0 ppm lead and no tulsi (group I, Control), 100 ppm lead (group II), or 100 ppm lead + 100 ppm tulsi (group III). After four weeks, disposition kinetics of intravenous single dose administration of SD (50 mg/kg) was determined in each bird. The mean values of elimination half-life, distribution-half, AUC, and volume of distribution were significantly ( $p < 0.01$ ) higher in lead-treated group than those in lead plus tulsi group (III). The clearance (Cl, mL/kg/h) of SD was slower in lead-treated groups than lead plus tulsi and control groups. The priming dose of SD (mg/kg) in chicks following single i.v. administration is proposed as 68.91 and 122.57 for group I (control), 51.42 and 79.21 for group II (lead) and 61.38 and 106.59 for group III (lead plus tulsi), at 8 h and 12 h interval, respectively. The maintenance doses ( $D_m$ , mg/kg) were minimum in group II (39.31 and 57.54) followed by group III (41.03 and 86.25) and control (47.13 and 100.80) at 8 and 12h interval, respectively. It is concluded that feeding of lead at 100 ppm dietary level altered the kinetics and dosage regimen of SD which were significantly ameliorated following simultaneous feeding with powdered tulsi (100ppm) in cockerels.

**Keywords:** Disposition kinetics, Sulphadimidine, Lead, Cockerels, *Ocimum Sanctum*

Sulphadimidine, a commonly-used sulfonamide antibacterial drug in veterinary practice, competes with paraminobenzoic acid (PABA) for the catalytic site of the enzyme dihydropterotic acid synthetase to inhibit the synthesis of folic acid, required for bacterial proliferation [1]. Sulfonamides are metabolized mainly in liver by acetylation followed by conjugation with sulfate or glucuronic acid, deamination and cleavage of heterocyclic ring [2]. *Ocimum sanctum*, commonly known as tulsi, is a plant widely used in Ayurveda. It has been shown to possess anti-inflammatory, antioxidant and cognition-enhancing properties [3]. Aqueous extract feeding also provided significant liver and aortic tissue protection [4]. *Ocimum sanctum* fixed oil produces hypotensive effect due to its peripheral vasodilatory action; it also increases blood-clotting time due to inhibition of platelet aggregation [5].


Lead is one of the most common heavy metals responsible for toxicity in man and animals. It is absorbed rapidly from gastrointestinal tract and lungs. In the bloodstream, lead binds with hemoglobin in erythrocytes and accumulates in bones. It excretes out

very slowly in urine and bile and fractionally by exfoliation of epithelial tissue including hair with a half-life of 20-30 years [6-8]. Thus, persistence of lead impairs the hepato-renal function and subsequently the pharmacokinetic pattern of the drugs that are metabolized by the liver and excreted in the urine. Variation in pharmacokinetics pattern of sulphonamides has also been reported in animals both in normal as well as stressful conditions [9-13]. In view of these facts, sulphadimidine was considered to study its disposition kinetics and to evaluate protective efficacy of *Ocimum sanctum* (tulsi) in chronically lead-intoxicated cockerels.

## MATERIALS AND METHODS

### Chemical

Pure analytical-grade lead acetate,  $Pb(C_2H_3O_2)_2 \cdot 3H_2O$  (CDH) was used in this study. Mature tulsi plants (*Ocimum sanctum*) were collected from Research and Development Farm, Medicinal and



**Fig 1.** Mean + S.E. Plasma concentration vs time plot of sulphadimidine, sulphadimidine + Lead, sulphadimidine + Lead + Tulsi following single dose (50 mg/kg) i.v. administration in cockerels (n=7)

Aromatic Plants, Department of Horticulture, N.D.U.A.T., and was identified taxonomically.

### Experimental animals

Four-week male white leghorn chicks were procured from government poultry farm chak ganjaria, Lucknow and were reared for four weeks for acclimatization before the start of the study. Four weeks chicks were used in this study and kept in deep litter system of housing and maintained on grower ration procured from U.P. state Agro Industries Corporation limited, Lucknow. Food and water were provided *ad libitum* throughout the study.

Medicated ration was prepared for about a week period and kept in close container. Dried tulsi leaves

were pulverized to prepare 100 ppm tulsi-medicated feed by mixing 600 mg of powder of tulsi in 6 kg food. Similarly, Lead (100 ppm)-treated ration was prepared by taking 145.21 mg of lead acetate containing lead equivalent to 100 ppm and mixing thoroughly. All the birds were fasted over night prior to the start of the experiment.

The ‘Animal Ethical Committee’ of Narendra Dev University of Agriculture and Technology, Kumarganj, Faizabad, Uttar Pradesh, India approved all the procedures adopted in the study.

### Animal treatment

Twenty-one, four weeks old male white leghorn chicks, were randomly divided into three groups of I (control), II (lead 100 ppm), and III (lead 100 ppm + tulsi 100 ppm) (seven birds each). The pharmacokinetics pattern of sulphadimidine was estimated after injecting a single dose (50 mg/kg, I.V.) of that in these groups after 12 weeks of feeding trial [11-13]. Blood samples were collected from wing vein in heparinised sterilized tubes before and after 0.08, 0.17, 0.25, 0.50, 1, 2, 4, 8, 12, 16, 20, 24, 28 and 32 h of drug administration.

### Analytical method

The concentration of free and total sulphonamide in plasma (µg/ml) was estimated according to the method of Bratton and Marshal, 1939 and Richterich, 1969. [14, 15] based on what measured by AUTOCHEM 2011 at 580 nm. Dosage regimen was calculated using information of single dose trial [16, 17]. The Pharmacokinetic analysis of plasma concentration time profile of sulphadimidine for each bird was performed with the aid of PHARMAKIT, (M/s Clyde Soft, Glasgow, UK).

### Statistical analysis

Comparison of pharmacokinetic parameters of treated and control groups were done using one way analysis of variance in SPSS software, Version 11. The *p* values ≤ 0.05 were considered significant.

**Table 1.** Mean plasma concentration (Mean ± SEM) (µg/ml) of sulphadimidine in blood following intravenous administration of single iv dose of sulphadimidine at 50 mg/kg in different groups of cockerel

Time (h)	Mean Plasma Concentration (Mean ± SEM)		
	Control (0 ppm)	Lead (100 ppm)	Lead (100 ppm) +Tulsi (100 ppm)
0.08	126.57 ± 1.30	111.28 ± 1.64	125.00 ± 1.70
0.17	94.14 ± 1.01	90.28 ± 2.14	95.42 ± 0.92
0.25	83.00 ± 0.48	79.28 ± 2.20	78.71 ± 1.55
0.5	63.42 ± 1.21	69.00 ± 2.14	64.00 ± 1.13
1	50.14 ± 1.59	54.57 ± 0.92	52.85 ± 0.70
2	38.71 ± 2.14	43.28 ± 1.44	41.00 ± 0.53
4	27.57 ± 1.65	33.28 ± 0.80	33.28 ± 0.99
8	19.14 ± 1.62	25.28 ± 0.96	22.14 ± 0.46
12	12.44 ± 0.88	16.14 ± 0.50	10.77 ± 0.17
16	6.85 ± 0.38	10.74 ± 0.28	6.91 ± 0.24
20	3.4 ± 0.22	7.64 ± 0.31	3.92 ± 0.54
24	1.12 ± 0.16	5.07 ± 0.42	0.85 ± 0.11
28	n. d.	3.37 ± .34	nd
32	n. d.	0.94 ± 0.13	nd

nd = not detected

## RESULTS

Mean plasma concentration ( $\mu\text{g mL}^{-1}$ ) of sulphadimidine following single dose ( $50 \text{ mg kg}^{-1}$ ) i.v. administration in different groups of chicks (control, 100 ppm lead and 100 ppm lead plus 100 ppm tulsi) is given in Table 1 and disposition curve has been depicted in Fig 1. The minimum therapeutic concentration ( $25 \mu\text{g mL}^{-1}$ ) of sulphadimidine was maintained for 4 hours in control and lead plus tulsi groups whereas 8 h in lead-intoxicated group. Sulphadimidine was detected up to 24 hours in control and lead plus tulsi groups whereas lead-intoxicated chicks were found to have sulphadimidine up to 32 hours of post administration. The plasma concentration-time profiles of sulphadimidine in all groups indicated that disposition of the drug followed two-compartmental open models. The values of various pharmacokinetic parameters were computed from plasma levels of free sulphadimidine in control and treated groups following single i.v. administration at the dose of  $50 \text{ mg kg}^{-1}$  (Table 2).

The mean value of extrapolated drug concentration during distribution phase (A,  $\mu\text{g/mL}$ ) was also less in lead-treated groups as compared to other groups whereas the value of extrapolated drug concentration during elimination phase (B,  $\mu\text{g/mL}$ ) did not reveal any variation among control and treated groups. The mean values of distribution rate constant ( $\alpha$ ,  $\text{h}^{-1}$ ) and elimination rate constant ( $\beta$ ,  $\text{h}^{-1}$ ) were significantly ( $p < 0.01$ ) less in lead-intoxicated groups as compared to control and lead plus tulsi group. The values of distribution rate constant ( $\alpha$ ,  $\text{h}^{-1}$ ) were  $5.227 \pm 0.464$ ,  $3.856 \pm 0.356$  and  $6.062 \pm 0.292$ , respectively, in control, lead and lead plus tulsi groups. The value of elimination rate constants ( $\beta$ ,  $\text{h}^{-1}$ ) were in the order as  $0.144 \pm 0.010$ ,  $0.105 \pm 0.002$  and  $0.137 \pm 0.001$ , respectively, in control, lead and lead plus tulsi groups.

The mean value of elimination half-life ( $t_{1/2}\beta$ , h) was significantly ( $p < 0.01$ ) higher in lead treated group as it was  $6.582 \pm 0.119$  in lead treated group as

compared to  $4.918 \pm 0.310$  in control and  $5.018 \pm 0.068$  in lead plus tulsi group. The mean value of distribution half-life ( $t_{1/2}\alpha$ , hours) was significantly ( $p < 0.01$ ) higher in lead-treated group as it was  $0.189 \pm 0.019$  in lead-treated group,  $0.138 \pm 0.012$  in control and  $0.114 \pm 0.004$  in lead plus tulsi group. There was a significantly ( $p < 0.01$ ) lower rate of transfer of the drug from peripheral to central ( $k_{12}$ ,  $\text{h}^{-1}$ ) compartment in lead-treated chicks in comparison to other groups whereas the rate of transfer from central to peripheral compartment ( $k_{21}$ ) did not reveal any variation among treated and control groups. The volume of distribution ( $V_{d\text{area}}$ ,  $\text{mL kg}^{-1}$ ) did not reveal any significant variation among all the groups as the values of  $V_{d\text{area}}$  were  $0.870 \pm 0.040$ ,  $0.866 \pm 0.014$  and  $0.814 \pm 0.008$  in controls, lead and lead plus tulsi group, respectively. The clearance (Cl,  $\text{mL h}^{-1}$ ) of sulphadimidine was slower in lead-treated birds than control and lead plus tulsi as the value of Cl was  $0.090 \pm 0.002$  in lead,  $0.124 \pm 0.008$  in control and  $0.111 \pm 0.001$  in lead plus tulsi group. The retention of sulphadimidine in the body was significantly ( $p < 0.01$ ) higher in lead-treated chicks than that in other groups as the values of mean residential time (MRT, h) was  $9.156 \pm 0.174$  in lead-treated group,  $6.734 \pm 0.429$  in control and  $6.956 \pm 0.095$  in lead plus tulsi treated group.

The dosage regimens for minimum plasma therapeutic concentration of  $25 \mu\text{g mL}^{-1}$  at 8 and 12 h interval for control and treated groups have been given in Table 3. At 8 h interval, the priming dose ( $D$ ,  $\text{mg kg}^{-1}$ ) for the cockerels fed on lead (100 ppm) after 12 weeks feeding trial was less than control and lead plus tulsi fed cockerels as it was computed to be 51.42 for lead-intoxicated group in comparison to 61.38 for lead plus tulsi and 68.91 for control group. Similarly, at 12 h interval, the  $D$  was only 79.21 for the cockerels of lead-treated groups, 106.59 for lead plus tulsi and 122.57 for the control group. Correspondingly, the maintenance doses ( $D_m$ ,  $\text{mg kg}^{-1}$ ) at 8 and 12 h interval were less for lead-treated chicks (39.3 and 57.54) as compared to lead plus tulsi (41.03 and 86.25) and control group (47.13

**Table 2.** Pharmacokinetic parameters after single dose  $50 \text{ mg kg}^{-1}$  iv sulphadimidine in control, lead-intoxicated and lead plus tulsi-medicated cockerels (Mean  $\pm$  S.E.M.)  $n = 7$

Parameter	Control	Lead (100 ppm)	Tulsi + Lead (each 100ppm)
A ( $\mu\text{g mL}^{-1}$ )	$106.750 \pm 4.235$	$72.580 \pm 1.794$	$107.434 \pm 4.053$
B ( $\mu\text{g mL}^{-1}$ )	$55.210 \pm 2.474$	$55.918 \pm 0.998$	$59.136 \pm 0.718$
$\alpha$ ( $\text{h}^{-1}$ )	$5.227 \pm 0.464$	$3.856 \pm 0.356$	$6.062 \pm 0.292$
$\beta$ ( $\text{h}^{-1}$ )	$0.144 \pm 0.010$	$0.105 \pm 0.002$	$0.137 \pm 0.001$
AUC ( $\mu\text{g mL}^{-1}\text{h}$ )	$410.444 \pm 25.023$	$550.811 \pm 13.324$	$445.979 \pm 6.978$
$t_{1/2} \alpha$ (h)	$0.138 \pm 0.012$	$0.189 \pm 0.019$	$0.114 \pm 0.004$
$t_{1/2} \beta$ (h)	$4.918 \pm 0.310$	$6.582 \pm 0.119$	$5.018 \pm 0.068$
$k_{10}$ ( $\text{h}^{-1}$ )	$0.401 \pm 0.030$	$0.233 \pm 0.007$	$0.373 \pm 0.013$
$k_{12}$ ( $\text{h}^{-1}$ )	$3.084 \pm 0.030$	$2.00 \pm 0.216$	$3.585 \pm 0.216$
$k_{21}$ ( $\text{h}^{-1}$ )	$1.882 \pm 0.172$	$1.734 \pm 0.140$	$2.239 \pm 0.083$
MRT (h)	$6.734 \pm 0.429$	$9.165 \pm 0.174$	$6.956 \pm 0.095$
Cl ( $\text{mL h}^{-1}\text{kg}^{-1}$ )	$0.124 \pm 0.008$	$0.090 \pm 0.002$	$0.111 \pm 0.001$
$V_{d\text{area}}$ ( $\text{mL kg}^{-1}$ )	$0.870 \pm 0.040$	$0.866 \pm 0.014$	$0.814 \pm 0.008$

A and B = extrapolated zero time plasma drug concentration during distribution and elimination phases, respectively = theoretical zero time concentration;  $\alpha$  and  $\beta$  = distribution and elimination rate constants, respectively; AUC = area under plasma concentration time curve;  $t_{1/2} \alpha$  and  $t_{1/2} \beta$  = distribution and elimination half lives, respectively;  $K_{10}$  = rate constant of drug elimination from central compartment;  $K_{12}$  and  $K_{21}$  = micro-rate constant of drug transfer from central to peripheral, peripheral to central compartment, respectively; MRT = mean residential time; Cl = total body clearance;  $V_{d\text{area}}$  = volume of distribution from AUC

**Table 3** Intravenous Dosage regimens of sulphadimidine to maintain desired therapeutic concentration of 25 µg mL<sup>-1</sup> at 8 and 12 h intervals in control, lead-intoxicated and lead plus tulsi-medicated cockerels.

Groups	Interval (h)	D (mg kg <sup>-1</sup> )	C <sub>ss</sub> (min) following D (µg mL <sup>-1</sup> )	C <sub>ss</sub> (max) following D (µg mL <sup>-1</sup> )	Dm (mg kg <sup>-1</sup> )	C <sub>ss</sub> (min) following Dm (µg mL <sup>-1</sup> )	C <sub>ss</sub> (max) following Dm (µg mL <sup>-1</sup> )
Control	8	51.42	43.21	102.52	39.31	25	39.31
	12	122.57	30.40	171.13	100.80	25	140.73
Lead (100ppm)	8	51.42	43.21	102.52	39.31	25	39.31
	12	79.21	34.42	125.78	57.54	25	91.37
Lead (100 ppm) + tulsi (100 ppm)	8	61.38	37.39	112.80	41.03	25	75.41
	12	106.59	30.89	161.85	86.25	25	130.96

D= Priming dose

Dm= Maintenance dose

C<sub>ss</sub> (min) = Minimum steady state concentrationC<sub>ss</sub> (max) = Maximum steady state concentration

and 100.80) respectively, as shown in Table 3.

## DISCUSSION

The disposition kinetics of sulphadimidine revealed that its minimum therapeutic concentration (25 µg mL<sup>-1</sup>) was maintained for 8 h in lead-treated cockerels which was significantly higher as compared to 4 h in lead plus tulsi and control groups. The drug was detected for 32 h in lead-treated cockerels as compared to other groups. Drug disposition suggests slow elimination and retention of drug for longer duration as well as maintenance of therapeutic concentration in lead-treated groups. Thus, dosage regimen varies among various treated and control groups.

The distribution half life ( $t_{1/2\alpha}$ , h) explains the distribution of sulphadimidine. The slow distribution of drug in lead-treated cockerels might be attributed to the alteration in transportation of drug as a result of toxic effect of lead on capillaries [18] and other tissues causing disruption in the transport of drug across membrane. The mean value of sum of  $K_{12}$  and  $K_{21}$  was also higher than  $K_{10}$  (elimination rate constant). Thus, the elimination of the drug is slower than its distribution in the body. This might be due to renal insufficiency caused by lead as previously reported [19, 20]. The mean value of elimination rate constant was significantly lower in lead-treated cockerels than other groups suggesting slow rate of elimination of sulphadimidine from body. Lead has been reported to produce nephrotoxic effect in poultry [19, 20]. The value of drug transfer rate constant  $k_{12}$  and  $k_{21}$  are significantly less in lead-treated cockerels which also suggests slower rate of the movement of the drug across these compartments in lead intoxicated cockerels [21]. The elimination half life ( $t_{1/2\beta}$ , h) was higher in lead-treated group than lead plus tulsi, which indicates slow elimination of drug from lead-treated cockerels. The slow rate of elimination of sulphadimidine is suggestive of slow hepatic metabolism and slow excretion through in lead treated group. As reported earlier, lead have hepatotoxic and nephrotoxic properties [14, 15] but tulsi nullifies these effects [4].

The volume of distribution ( $V_{d\text{area}}$ ) was considered as indicator of extravascular distribution of the drug. However, it was comparable in control and other groups indicating equal distribution of the drug in body tissues.

Area under curve (AUC) of concentration-time profile explains the relative availability of the drug in the body. The values of AUC of sulphadimidine were the highest in lead-treated cockerels followed by lead plus tulsi and control groups. The effectiveness of drug depends not only on the plasma concentration of drug achieved but also duration for which these concentrations exist in the body [22]. That may be correlated with availability of therapeutic concentration for longer duration i.e. 8 h in lead-treated group in comparison to only 4 h in other groups.

The value of MRT was higher in lead-treated cockerels which are suggestive of retention of the drug for longer duration in lead-intoxicated cockerels. The clearance ( $Cl$  mL Kg<sup>-1</sup> h<sup>-1</sup>) of sulphadimidine was slower in lead-treated cockerels than that in other groups. The pharmacokinetic pattern of drug in lead-treated cockerels is attributed to impaired hepatic metabolism of sulphadimidine and renal excretion of the drug as the sulphonamides are metabolized by acetylation in the liver and excreted by kidneys. Altered pharmacokinetic pattern of sulphonamides with longer half-life in comparison to untreated control was also observed in cockerels treated with pesticide [23] and carbon tetrachloride (CTC) [24]. Similarly, allyl alcohol-induced hepatotoxic sheep also revealed prolonged retention of the sulphonamides [25].

The main objective of pharmacokinetic study of a drug is to calculate its dosage regimen. The cockerels fed on lead (100 ppm) were computed to have least priming and maintenance doses in comparison to lead plus tulsi and control groups at both 8 and 12 h interval to maintain minimum therapeutic concentration of 25 µg mL<sup>-1</sup>. It is concluded from this study that lead-induced toxic effects altered the disposition kinetic of sulphadimidine in cockerels as reported earlier that lead have hepatotoxic and nephrotoxic effects which might alter the kinetic of drug metabolized by body enzyme system. When lead-treated animals were simultaneously treated with tulsi (100 ppm), the tulsi nullified the toxic effect of lead. These findings may be correlated with other species of animals and other drugs as well.

## ACKNOWLEDGEMENT

The author would like to thank the Dean, CV Sc & AH and Director of Research, NDU&T, Kumarganj,

Faizabad, UP for providing financial and facilities to carry out this research.

## REFERENCES

- Mandell GL, Sonde MA. Goodman and Gillman's. The Pharmacological basis of therapeutics. 8th ed. Mc Millan Publ, New York, USA 1990; pp:1095-2015.
- Bevill RF. Sulphonamide. In: Booth NH, MacDonald LE. eds. Veterinary pharmacology and therapeutics. 1988 Panima Publ Corp, New Delhi, pp: 785-95.
- Yanpallewar SU, Rai S, Kumar M, Acharya SB. Evaluation of antioxidant and neuroprotective effect of *Ocimum sanctum* on transient cerebral ischemia and long term cerebral hypoperfusion. *Pharmacol Biochem Behav* 2004; 79:155-64.
- Geetha R, Vasudevan DM. Inhibition of lipid peroxidation by botanical extracts of *Ocimum sanctum*: in vivo and in vitro studies. *Life Sci* 2004; 76:21-8.
- Singh S, Rehan HMS, Majumdar DK. Effect of *Ocimum sanctum* fixed oil on blood pressure, blood clotting time and pentobarbitone-induced sleeping time. *J Ethnopharmacol* 2001; 78:139-43.
- Gross SB, Pfister EA, Yeager DW, Kehoe RA. Lead in human tissues. *Toxicol App Pharmacol* 1975; 32:638-51.
- Somlay I, Lanzlo V, Marto S. Lead concentration of vegetation bordering the roads in the neighbourhood of town Keszthely Magy. *Allatorr Lapja* 1986; 41:432-4.
- Paul BH, Robert PB. In: Casarett and Doull's Toxicology. The Basic Science of Poisons. 1980, 2nd ed, Macmillon Publishing Co Inc, New York, USA, pp: 417-8.
- Kumar S, Sharma LD, Ahmad AH. Pharmacokinetics of sulfadimidine in hepatically-impaired buffalo calves. *Ind J Vet Med* 1997; 17:39-44.
- Kumar S, Ahmad AH, Sharma LD. Pharmacokinetics of sulfamethoxypyridazine in normal and renal impaired goats following single dose intravenous administration. *J appl Animal Res* 1999; 16:147-52.
- Ram S, Singh SP, Varma R, Choudhary G. Effect of arsenic on the disposition kinetics of sulfadiazine and protective efficacy of pipili (*Piper longum*) in cockerels. *IJPT* 2008; 7:175-9.
- Singh SP, Sharma LD, Ahmad AH. Alteration in pharmacokinetic pattern of SMPZ in cockerels fed on alphamethrin. *Proceeding of National Symposium on Indigenous medicinal plants and STOX seminar on Emerging Environmental challenges: Production strategies*. IVRI, July 3-5 1998, P.26.
- Singh SP, Sharma LD, Ahmad AH. Pharmacokinetics of sulphomethoxypyridazine (SMPZ) in monocrotophos feed cockerels. *Proc XVII Ann Conf STOX* Oct, 22-24. 1998, Tirupati: p.50.
- Bratton AC, Marshall EK. New coupling component for sulphanilamide determination. *J Biol Chem* 1939; 128:537-50.
- Richterich R. Clinical chemistry: Theory and practice. Translated from second German ed, 1969, Academic Press, New York, USA.
- Johu D B. Clinical Pharmacokinetics. *J Vet Pharmacol Therap* 1980;3:
- Snedecor GW, Cochran WG. Statistical methods. 6th Edition 1967; Oxford and IBH Publishing Co. Calcutt, P.258.
- Goodman IS, Gilman AG, Rau TW, Nies AS, Taylor P. The Pharmacological Basis of Therapeutics. 9th Ed, Mc Millan Publ, New York, USA, 2001:1862-5.
- Goyer RA. Mechanism of lead and cadmium nephrotoxicity. *Toxicology Lett* 1989; 46:153-62.
- Jeng SL, Lee SJ, Liu YF, Yang SC, Liou PP. Effect of lead ingestion on concentration of lead in tissues and eggs of laying Tsaiya ducks in Taiwan. *Poultry Sci* 1997; 76:13-6.
- Notari RE. Biopharmaceutics and clinical pharmacokinetics. 3rd Ed. Marcel Dekker. Inc. New York, USA, 1980.
- Niazi S. Text book of Biopharmaceutics. Appleton Century Coofts New York, USA, 1979.
- Singh SP. Immunotoxicological and teratological studies of Alphamethrin, Butachlor and Monocrotophos in experimental animals. PhD Thesis Submitted to GB Pant Uni Of Agric and Tech, Pantnagar, India, 1997.
- Kumar S. Pharmacokinetics of sulphadimidine in liver damaged buffalo calves (*Bubalus bubalis*). PhD Thesis GB Pant University of Agriculture and Technology, Pantnagar, India, 1991.
- Argaw T, Brugere H. Changes in disposition kinetics of sulphadimidine in response to acute regional liver damage in sheep. *J Vet Med Series A* 1990; 37:787-94.

## CURRENT AUTHOR ADDRESSES

- Prakash, Department of Pharmacology & Toxicology, College of Veterinary Science & Animal Husbandry, N.D.U.A. & T., Kumarganj, Faizabad, (U.P.), India.
- P. Singh, Department of Pharmacology & Toxicology, College of Veterinary Science & Animal Husbandry, N.D.U.A. & T., Kumarganj, Faizabad, (U.P.), India.
- Varma, Department of Pharmacology & Toxicology, College of Veterinary Science & Animal Husbandry, N.D.U.A. & T., Kumarganj, Faizabad, (U.P.), India. E-mail: drgovindvet@yahoo.co.in (Corresponding author)
- K. Choudhary, Department of Pharmacology & Toxicology, College of Veterinary Science & Animal Husbandry, N.D.U.A. & T., Kumarganj, Faizabad, (U.P.), India.
- Sajan, Department of Pharmacology & Toxicology, College of Veterinary Science & Animal Husbandry, N.D.U.A. & T., Kumarganj, Faizabad, (U.P.), India.