

# Antioxidant Activity of *Eclipta alba* L. in Normal Rat Liver

A. SANDEEP MAJUMDAR, M. NATVARLAL SARAF and R. YADAV KAMBLE

For author affiliations, see end of text.

Received October 11, 2009; Revised March 3, 2010; Accepted May 27, 2010

This paper is available online at <http://ijpt.iuims.ac.ir>

## ABSTRACT

The present study investigated the antioxidant potential of the ethyl acetate fraction of the aerial parts *Eclipta alba* L. Hassk (EA) which was widely used as hepatoprotective plant. EA was orally administered at doses of 50, 100 and 200 mg/kg (n=6) for 7 days in male Charles Foster rats. The extent of hepatoprotective potential of *E. alba* was studied by assessing the biochemical parameters like lipid peroxides (LPO), superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), glutathione reductase (GR), ascorbic acid and  $\alpha$ -tocopherol in the liver of rats. Oral administration of the EA significantly decreased LPO and elevated the activity of antioxidant enzymes SOD, CAT, GPx and GR as well as endogenous levels of ascorbic acid and  $\alpha$ -tocopherol in the liver of male Charles Foster rats. This study has revealed the significant antioxidant potential of *E. alba* in rat liver.

**Keywords:** comma separated keywords

*Eclipta alba* L. Hassk. (Asteraceae), commonly known as Bhingaraja (Sanskrit), Maka (Marathi), and Bhangra (Hindi) has been reported to show protective effect on experimental liver damage in rats and mice [1]. The plant has been reported for the treatment of liver cirrhosis and infective hepatitis [2]. *E. alba* (EA) powder has been found to counteract carbon tetrachloride (CCl<sub>4</sub>)-induced increase in liver weight, hepatic lipid peroxidation, liver  $\gamma$ -glutamyl transpeptidase, serum alanine transferase, serum alkaline phosphatase and serum albumin to globulin ratio in rats [3]. According to Ayurvedic text, EA is the main herb for hair and serosis; maintaining and rejuvenating hair, teeth, memory, sight and healing [4]. It also possesses an anti-aging property.

The plant has been reported to exhibit protective effect on CCl<sub>4</sub>-induced acute liver damage by reducing centrilobular necrosis, hydropic degeneration and fatty change of the hepatic parenchymal cells [5]. Wagner *et al.* (1986) confirmed that the coumestan constituents of the plant, wedelolactone and demethylwedelolactone, are responsible for the potent antihepatotoxic activities in CCl<sub>4</sub>-, galactosamine- and phalloidin-induced liver damage in rats. However, there are no reports on the effect of the plant on endogenous liver antioxidant system. On the basis of earlier reports [6], it was considered worthwhile to take the ethyl acetate fraction

of *Eclipta alba* (EA) for evaluating the antioxidant activity in liver. Hence, the present study aimed to assess the antioxidant potential of EA by estimating the activity of antioxidants (enzymatic and non-enzymatic) and levels of lipid peroxides on oral administration.

## MATERIAL AND METHODS

### Plant material

Aerial parts of *E. alba* L. Hassk. were procured and authenticated by Dr. Ashish Phadke, Head, Department of Draavyaguna (Herbal Pharmacology), Y.M.T. Ayurvedic Medical College, Khargar, Navi Mumbai, Mumbai, India. Specimen sample of *Eclipta alba* was preserved in the Pharmacognosy department of Bombay College of Pharmacy, Mumbai, India with the voucher no. 14/E. alba/ Majumdar/ Feb' 2004.

### Animals

Inbred male Charles Foster rats (150-200 g) were used. Ethical clearance for the study was obtained from the Institutional Animal Ethics Committee of the Bombay College of Pharmacy. They were housed in standard environmental conditions and allowed free access to standard laboratory rat pellet diet and water *ad libitum*.

**Table 1.** Effect of EA administration on the antioxidant defense system in rat liver

Parameters	Control	EA 50 mg/kg	EA 100 mg/kg	EA 200 mg/kg
SOD activity <sup>a</sup>	1.35 ± 0.05	1.27 ± 0.04	2.36 ± 0.27*	3.17 ± 0.29**
CAT activity <sup>a</sup>	10.28 ± 0.66	12.89 ± 1.45	23.40 ± 1.40**	25.05 ± 2.91**
GPx activity <sup>a</sup>	0.35 ± 0.08	0.31 ± 0.05	0.61 ± 0.04*	4.69 ± 0.28**
GR activity <sup>a</sup>	9.51 ± 0.98	13.34 ± 1.84	25.91 ± 2.81**	29.72 ± 4.59**
Lipid peroxidation <sup>b</sup>	4.14 ± 0.44	1.44 ± 0.16**	1.11 ± 0.23**	0.44 ± 0.22**
Vitamin C <sup>c</sup>	0.72 ± 0.01	0.98 ± 0.10*	1.09 ± 0.01**	1.23 ± 0.02**
Vitamin E <sup>c</sup>	0.19 ± 0.02	0.27 ± 0.00*	0.39 ± 0.02**	0.41 ± 0.01**

All values are expressed as the mean ± SD, for 6 rats in each group. <sup>a</sup> Units (mg protein)<sup>-1</sup>, <sup>b</sup> nmol of MDA (mg protein)<sup>-1</sup> and <sup>c</sup> µg (mg protein)<sup>-1</sup>. \**p* < 0.05, \*\**p* < 0.01, different from control-vehicle treated group evaluated by one-way analysis of variance test followed by Mann-Whitney *U*-test, or Kruskal Wallis's test, if necessary.

### Preparation of the Fraction

The aerial parts of the plant *E. alba* were shade dried and the coarse powder (1 kg) was exhaustively extracted with methanol in a soxhlet extractor (yield: 15.7% w/w). The solvent was evaporated and the extract was suspended in water and heated on a steam bath (30 min, < 80° C). This methanolic extract was filtered under vacuum using Whatmann paper and air dried. The aqueous phase was partitioned with ethyl acetate five to six times. The organic fraction was dried with sodium sulphate and the solvent evaporated to yield ethyl acetate fraction of *Eclipta alba* (EA). The yield of the fraction was 2.26% (w/w).

### Determination of total phenolic content of EA

The assay of total phenolic content of EA was determined as per the procedure stated by Randolph (1995) in Association of Agricultural Chemist [7].

### Acute toxicity evaluation of the fraction

Acute toxicity studies were performed according to Organization for Economic Co-Operation and Development (OECD)-423 guidelines [8].

### Experimental procedure

Young adult Charles Foster rats were divided into four groups. Group 1 served as control and groups 2, 3 and 4 were administered a dose of 50, 100 and 200 mg/kg 0.3% carboxymethyl cellulose suspension of EA in distilled water respectively by oral gavage once daily between 10.00 am and 11.00 am for 7 days. Animals were sacrificed 1 h after the last drug or vehicle administration on day 7. A 10% liver homogenates prepared in ice-cold 20 mM phosphate buffer (pH 7.0) was incubated with 0.1% tritonX-100 for 1h at 5°C and centrifuged (20000×g, 20 min, 4°C). The supernatants were used for measuring activity of antioxidant enzymes such as CAT [9], SOD [10], GPx [11] and GR [12]. The results were expressed as units/mg protein. Further, 10% liver homogenates prepared in KCl (150 mM) was used for determination of LPO [13], ascorbic acid [14] and α-tocopherol [15]. The protein content of the rat liver homogenate was determined by the method of Biuret reaction using Bovine serum albumin (BSA) as a standard [16].

### Statistical analysis

All results were evaluated by one-way analysis of variance test followed by Mann-Whitney *U*-test or Kruskal Wallis's test, if necessary. Significance was set at *p* < 0.05.

## RESULTS AND DISCUSSION

The results are shown in Table 1.

### Effects of EA on enzymatic antioxidant system

The present study revealed that oral administration of EA for 7 days at a dose of 100 mg/kg and 200 mg/kg in rats significantly elevated the activity of the enzymes involved in scavenging reactive oxygen species such as superoxide dismutase and catalase. This indicated that EA improved the enzymatic antioxidant status in rat liver since it is known that a marked increase in SOD and CAT activity can offer first line protection against the damaging effects of superoxide radicals in the liver [17]. EA administration for 7 days at a dose of 100 mg/kg and 200 mg/kg resulted in improved activity of enzymes GPx and GR reflecting elevated levels of GSH in rat liver, which may be one of the key mechanisms mediating the potent hepatoprotective action of *E. alba*. However no significant change in the enzyme levels were observed in the groups treated with 50 mg/kg.

### Effects of EA on non-enzymatic antioxidant system

EA treatment significantly inhibited lipid peroxides in rat liver in a dose-dependent manner. This proves the potent antioxidant potential of EA, since it is known that oxygen radicals produce peroxidation of lipids leading to disruption of cell membrane and structural damage to the tissue [18]. EA was also found to significantly improve the levels of cellular antioxidants vitamin E and vitamin C in rat liver. Vitamin E is the first antioxidant present in the cell membrane that counteracts the entry of free radicals in the cells and ultimately gets transformed into tocopheroxyl radicals [19]. Vitamin C acts as a co-antioxidant by regenerating α-tocopherol from α-tocopheroxyl radical produced during scavenging of reactive oxygen species (ROS) [20]. Although vitamin E levels were elevated distinctly only at 100 and 200 mg/kg dose, vitamin C levels in the

liver homogenates were significantly elevated in all the three groups treated with 50, 100 and 200 mg/kg. Thus, EA imparted antioxidant activity by elevating the levels of endogenous antioxidants vitamin E and vitamin C.

These observations are consistent with the conclusion that the ethyl acetate fraction of *E. alba* improves both enzymatic and non enzymatic antioxidant status in rat liver. Further acute toxicity studies of EA at a dose of 2 g/kg p.o. revealed no evident toxicity or mortality, thereby confirming the traditional safety profile of *E. alba* as per the citation [1].

Increased oxidative stress has been suggested to play a major role in liver damage due to cirrhosis of liver [21]. Hence, when the normal levels of antioxidants are not enough for the eradication of free-radical-mediated injury, then administration of antioxidant compounds has a potential role to play in reducing liver damage. Several phenolic components in plants like flavonoids [22], coumarins [23] and anthraquinones [24] are reported to exhibit antioxidant activity. Thus, the potent antioxidant activity expressed by the fraction of *E. alba* might be due to the presence of 40.95 % of phenolic compounds as per our finding, predominantly the coumestan derivatives, wedelolactone and demethylwedelolactone and also triterpenes and flavonoids [6].

Earlier studies have reflected that *E. alba* expressed hepatoprotective activity by regulating hepatic lysosomal enzymes [25]. Our study suggests that hepatoprotective action of *E. alba* may be also due to its antioxidant potential, thus justifying its use in liver tonics as a prophylactic supplement for combating oxidative-stress-induced liver damage. The result of this investigation is of importance, since *E. alba* is widely used in India for the traditional treatment of liver disorders.

#### ACKNOWLEDGEMENT

The authors wish to acknowledge the University Grants Commission (UGC), India for the minor research grant.

#### REFERENCES

- Singh B, Saxena AK, Chandan BK, Agarwal SG, Bhatia MS, Anand KK. Hepatoprotective effect of ethanolic extract of *Eclipta alba* on experimental liver damage in rats and mice. *Phytother Res* 1993; 7:154-8.
- Chopra RN, Nayar SL, Chopra IC. Glossary of Indian Medicinal Plants. New Delhi: Council of Scientific and Industrial research, 1966; pp.104.
- Chandra T, Sidique J, Somasundaram S. Effect of *Eclipta alba* on inflammation and liver injury. *Fitoterapia* 1987; 58:23-32.
- Tirtha SSS, In: The Ayurveda Encyclopedia Natural secrets to Healing, Prevention, and longevity. Indian Medical Science series No.68. Sri Satguru Publications, Delhi, India, 1998; pp. 77.
- Khin MM, Nyunt N, Khin MT. The protective effects of *Eclipta alba* on carbon tetrachloride induced acute liver damage. *Toxicol Appl Pharmacol* 1978; 45:723-8.
- Wagner H, Geyer B, Kiso Y, Hikino H, Rao GS. Coumestans as the main active principles of the liver drugs *Eclipta alba* and *Wedelia claudulacea*. *Planta Med* 1986; 52:370-4.
- Ecobichon DJ. The Basis of Toxicology Testing. CRC Press, New York. 1997; pp. 43-86.
- Randolph HD. In: Food consumption; Additives; Natural contaminants. AOAC (16<sup>th</sup> Editions). 952.03
- Ming S, Zigman S. An improved spectrophotometric assay for superoxide dismutase based on epinephrine autoxidation. *Anal Biochem* 1978; 90: 81-9.
- Beers RF, Sizer IW. A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. *J Biol Chem* 1952; 195:133-40.
- Carrillo MC, Kanal S, Nokuibo M, Kitani K. (-) Deprenyl induces activities of both superoxide dismutase and catalase but not glutathione peroxidase in the striatum of young male rats. *Life Sci* 1991; 48:517-21.
- Acedo A, Mezzetti A, Dillio C, Calafiore AM, De Cesare D, Bosco G, Acciaai N, Cappelletti L, Federici G, Cuccurullo F. Effect of ischaemia-reperfusion on glutathione peroxidase, glutathione reductase and glutathione transferase activities in human heart protected by hypothermic cardioplegia. *Free Radic Res Commun* 1990; 8: 85-91.
- Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 1979; 95:351-8.
- Omaye ST, Turnbull JD, Sauberlich HE. Selected methods for the determination of ascorbic acid in animal cells, tissues and fluids. *Methods Enzymol* 1979; 62:1-11.
- Desai ID. Vitamin E analysis method for animal tissues. *Methods Enzymol* 1984; 105: 138-43.
- Gornall AG, Bardawill CJ, Davis M. Determination of serum proteins by means of the biuret reaction. *J Biol Chem* 1949; 177:751-6.
- Deneke SM, Fanburg BL. Regulation of cellular glutathione. *Am J Physiol* 1989; 257:L163-73.
- Sevanian A, Hochstein P. Mechanisms and consequences of lipid peroxidation in biological system. *Ann Rev Nutr* 1985; 5:365-75.
- Mccay PB. Vitamin E: interactions with free radicals and ascorbate. *Annu Rev Nutr* 1985; 5:323-40.
- Packer JE, Slater TF, Willson RL. Reaction of the carbon tetrachloride related peroxy free radical (CCl<sub>3</sub>O<sub>2</sub>.) with amino acid: pulse radiolysis evidence. *Life Sci* 1978; 23:2617-20.
- Cabre M, Camps J, Ferre N, Paternain JL, Joven J. The antioxidant and hepatoprotective effects of zinc are related to hepatic cytochrome P450 depression and metallothionein induction in rats with experimental cirrhosis. *Int J Nutr Res* 2001; 71:229-36.
- Robak J, Shirdi F, Wolbis M, Krolkowska M. Screening of the influence of flavonoids on lipoxygenase and cyclooxygenase activities as well as on nonenzymic lipid peroxidation. *Pol J Pharmacol Pharm* 1988; 40:451-8.
- Paya M, Halliwell B, Hoult JR. Interaction of a series of coumarins with reactive oxygen species: scavenging of superoxide, hypochlorous acid and hydroxyl radicals. *Biochem Pharmacol* 1992; 44:205-14.
- Malterud KE, Farbroth TG, Huse AE, Sund RB. Antioxidant and radical scavenging effects of anthraquinones and anthrones. *Pharmacol* 1993; 47:77-85.
- Saxena AK, Singh B, Anand KK. Hepatoprotective effects of *Eclipta alba* on subcellular levels in rats. *J Ethnopharmacol* 1993; 40:155-61.

#### CURRENT AUTHOR ADDRESSES

- Sandeep Majumdar, Department of Pharmacology, Bombay College of Pharmacy, Mumbai, India. E-mail: anuradha.majumdar@gmail.com (Corresponding author)
- Natvarlal Saraf, Department of Pharmacology, Bombay College of Pharmacy, Mumbai, India.
- Yadav Kamble, Department of Pharmacology, Bombay College of Pharmacy, Mumbai, India.