



## Drug-drug interactions: Influence of verapamil on the pharmacokinetics of sitagliptin in rats and *Ex vivo* models

Siddhartha Nuthakki<sup>1</sup>, Sivaprasad Pendyala<sup>1</sup>, Sivaramakrishna Kondru<sup>1</sup>, Ravindrababu Pingili<sup>1</sup>,  
Naveenbabu Kilaru<sup>2\*</sup>

<sup>1</sup> Department of Pharmacology, KVSR Siddhartha College of Pharmaceutical Sciences, Vijayawada, Andhra Pradesh, India

<sup>2</sup> Department of Pharmaceutics and Pharmaceutical Biotechnology, KVSR Siddhartha College of Pharmaceutical Sciences, Vijayawada, Andhra Pradesh, India

### Please cite this article as:

Nuthakki S, Pendyala S, Kondru S, Pingili R, Kilaru N. Drug-drug interactions: Influence of verapamil on the pharmacokinetics of sitagliptin in rats and *Ex vivo* models. Iranian J Pharmacol Ther. 2018 (May);16: 1-11.

### ABSTRACT

P-glycoprotein (P-gp) and cytochrome P450 3A4 (CYP3A4) play a significant role in the disposition and elimination of drugs. The objective of this study was to investigate the mechanism underlying the interaction between sitagliptin (substrate of P-gp and CYP3A4) and verapamil (known modulator of P-gp and CYP3A4) using *in vivo*, *ex vivo* and *in situ* models. Rats were treated with sitagliptin (10 mg/kg, oral and/or 5 mg/kg, intravenous) alone and in combination with verapamil (40 mg/kg, oral) for 15 consecutive days. Blood samples were collected from the tail vein on 1st day in single dose pharmacokinetic study (SDS) and on 15th day in multiple dose pharmacokinetic study (MDS). The plasma concentrations of sitagliptin were significantly higher in the verapamil pretreated group when compared to sitagliptin control group. Verapamil pretreatment significantly increased the mean area under the plasma concentration-time curve from 0 to 24h (AUC<sub>0-24h</sub>), peak plasma concentration (C<sub>max</sub>), percent absolute bioavailability (AB%), elimination half-life (t<sub>1/2</sub>) and decreased the volume of distribution (V<sub>z</sub>/F), clearance (CL/F) and apparent volume of distribution at steady state (V<sub>ss</sub>/F) of sitagliptin in both SDS and MDS (oral and intravenous). *Ex vivo* study results showed that the apparent permeability coefficient (P<sub>app</sub>), net efflux and efflux ratio values were significantly increased by the known P-gp and CYP3A4 inhibitors (itraconazole and ketoconazole) and verapamil. In single pass intestinal perfusion (*In situ*) study, the effective permeability coefficient (P<sub>eff</sub>) and intestinal absorption rate constant (K<sub>a</sub>) were increased in the presence of verapamil (p<0.05). The present study results revealed that verapamil enhanced the bioavailability of sitagliptin probably by inhibiting its absorption via P-gp and/or the CYP3A4-mediated biotransformation in rats. Verapamil can be co-administered with sitagliptin without dose adjustment due to high safety margin of sitagliptin.

**Conflicts of Interest:** Declared None

**Funding:** Siddhartha Academy of General and Technical Education (SAGTE)

### Keywords

P-glycoprotein,  
CYP3A4,  
Drug-drug interactions,  
Everted gut sacs,  
Sitagliptin

### Corresponding to:

Naveenbabu Kilaru,  
KVSR Siddhartha College of  
Pharmaceutical Sciences,  
Polyclinic road,  
Siddhartha Nagar,  
Vijayawada,  
Andhra Pradesh,  
India-520010

### Email:

[naveenbabukvsr@gmail.com](mailto:naveenbabukvsr@gmail.com)

Received: 14 Nov 2017

Revised: 19 Dec 2017,

Accepted: 6 Mar 2018

### INTRODUCTION

Drug efflux transporters (especially P-glycoprotein, P-gp) and drug-metabolizing enzymes, particularly cytochrome

P450 3A4 (CYP3A4) play an important role in the absorption, distribution and elimination of a wide variety of drugs [1, 2]. In the marketed drugs, upto 50% have been identified to be P-gp substrates and/or inhibitors and 60% identified to be CYP3A4 substrates and/or inhibitors [3, 4]. The broad and overlapping substrate specificity of P-gp and CYP3A4 renders dual substrates prone to numerous potential drug-drug interactions [5, 6]. P-gp and CYP3A4 may act synergistically in limiting the oral bioavailability of drugs thereby the pharmacokinetics of a drugs may be altered when co-administered with compounds which inhibit or induce P-gp and CYP3A4 [7].

Diabetes and hypertension are intertwined clinical conditions that share a significant overlap in underlying risk factors and complications [8]. These complications can be divided into macrovascular (coronary artery disease, myocardial infarction, congestive heart failure, stroke, and peripheral vascular disease) and microvascular (retinopathy, nephropathy, neuropathy and cardiomyopathy) are conventionally linked to hyperglycemia, studies have shown that hypertension constitutes an important risk factor [9]. Sitagliptin is a first selective dipeptidyl peptidase-4 (DPP-4) inhibitor approved for the treatment of type 2 diabetes [10]. In addition to antihyperglycemic activity [11], sitagliptin also has cardioprotective [12], antioxidant [13], renoprotective [14] and neuroprotective [15] activities.

Verapamil, the prototype of the Calcium channel blockers (CCBs) has been extensively used in various cardiovascular conditions including hypertension [16], angina, supraventricular arrhythmias [17], heart failure [18], and congestive heart failure [19]. It has been reported to have cardioprotective [20], nephroprotective [21], neuroprotective [22], renoprotective [23], hepatoprotective [24], antioxidant and anti-inflammatory [25] activities. It is used as second line add-on drug and has beneficial effects in managing hypertension in patients with diabetes [26]. It is a substrate and inhibitor of P-glycoprotein [27] and also inhibits CYP3A4 isoenzymes [28].

Many clinically significant interactions have been reported with verapamil. Previous studies reported that verapamil inhibited the P-gp and CYP3A4-mediated metabolism of levobupivacaine, lidocaine [29], imipramine [30], atorvastatin [31], simvastatin, nilotinib [32] cerivastatin [33], darifenacin, ethosuximide [34], fentanyl, pimozide, sildenafil, terfenadine and zolpidem [35]. Pharmacokinetically, sitagliptin is a substrate of P-gp and CYP3A4 thereby it is considered likely to cause interactions with P-gp and CYP3A4 substrates and/or inhibitors [36]. Cyclosporine A significantly inhibited P-gp mediated transport of sitagliptin [37]. However, data (either clinical or preclinical) are lacking on the pharmacokinetic interactions between sitagliptin and verapamil. Therefore, the study was planned to evaluate the effect of verapamil on the pharmacokinetics of sitagliptin in rats and to explore the roles of P-gp and CYP3A4 in the absorption of sitagliptin using rat intestinal sacs *ex vivo* and *in situ* models.

## MATERIALS AND METHODS

### Drugs and chemicals

Sitagliptin and verapamil were obtained as gift samples from Actis Pharma, Hyderabad, India and Torrent Pharmaceuticals, Secunderabad, India, respectively. Ketoconazole was obtained from Mylan Pharmaceuticals, Hyderabad. Methanol, acetonitrile and water of high-performance liquid chromatography (HPLC) grade were purchased from Finar Chemicals Ltd, Ahmadabad, India. All other chemicals and reagents used were of analytical grade.

### Experimental animals

Animal experiments were performed according to the institutional guidelines for the care and use of laboratory animals, and approved by the animal ethics committee of KVSRR Siddhartha College of Pharmaceutical Sciences (SCOPS), Vijayawada, Andhra Pradesh, India (Protocol No: KVSRRSCOPS/11-03-04-007). The college was recognized by the Govt. of India (993/PO/E/S/06/CPCSEA). Male wistar rats (180-220 g) were procured from Mahaveer Enterprises, Hyderabad, and Andhra Pradesh, India. Animals were housed six per cage and given free access to food (Hindustan Lever, Mumbai, India) and water *ad libitum* in animal house at the KVSRR SCOPS. Before starting the experiments, animals were kept under standard laboratory conditions (12/12 h light/dark cycle, 22±2°C and 50-60% humidity) for at least a week.

### Experimental design

The entire study consists of four experiments as it has been described previously by Pingili *et al.* in our laboratory [38]. First two experiments are single dose pharmacokinetic study (SDS) and multiple dose pharmacokinetic study (MDS) *in vivo*. In these studies, animals were treated with sitagliptin alone and in combination with verapamil once daily for 15 consecutive days [39]. The doses of sitagliptin and verapamil were calculated and selected based upon the observations from the earlier experiments. Third and fourth experiments are *ex vivo* absorption studies using rat gut sacs [40] and *in situ* drug permeability (Single pass intestinal perfusion) studies to determine the role of P-gp on the absorption of sitagliptin using rat small intestine [41, 42]. Finally, the concentrations of sitagliptin were estimated in the plasma and other (*in vitro* & *in situ*) samples by using reversed phase-high pressure liquid chromatography (RP-HPLC).

### *In vivo* oral pharmacokinetic (SDS and MDS) studies

Sitagliptin and verapamil were suspended in sodium carboxymethylcellulose (0.5%) for oral administration. Male Wistar rats were randomly divided into two groups of six animals in each group. Group I (sitagliptin control group), treated orally with sitagliptin (10 mg/kg) alone and Group II (treatment group): pre-treated (30 min) with verapamil (40 mg/kg, oral) followed by sitagliptin (10 mg/kg, oral). The treatment was given once daily for 15 consecutive days. After treatment, 200 µL blood samples were collected from

tail vein at intervals of 0.5, 1, 2, 3, 4, 6, 8, 12 and 24 h on the 1<sup>st</sup> day in SDS and on the 15<sup>th</sup> day in MDS. After collection of each blood sample, an equal volume of heparinized 0.9% NaCl (20 IU/mL) was administered to the animals. Blood samples were collected in heparinized eppendorf tubes and kept on ice during the experiment. The plasma was separated by centrifugation (Remi, R4C Compact model, Mumbai, India) at 5000 rpm for 6 min and stored at -20°C until analysis.

#### *In vivo intravenous pharmacokinetic (SDS and MDS) studies*

For intravenous administration, sitagliptin phosphate was directly dissolved in phosphate buffer saline (pH 7.2) and verapamil was suspended in sodium carboxymethylcellulose (0.5%) for oral administration. Wistar rats were randomly divided into two groups of six animals in each group and their lateral tail veins were cannulated for blood withdrawal and sitagliptin administration. Group I (sitagliptin control group), treated intravenously with sitagliptin (5 mg/kg) alone and Group II (treatment group), pre-treated (30 min) with verapamil (40 mg/kg, oral) followed by sitagliptin (5 mg/kg) intravenously. The treatment was given once daily for 15 consecutive days.

After treatment, 200 µL blood samples were collected from tail vein at intervals of 5, 15, 60, 120, 180, 240 and 720 min on the 1<sup>st</sup> day in SDS and on the 15<sup>th</sup> day in MDS. After collection of each blood sample, an equal volume of heparinized 0.9% NaCl (20 IU/mL) was administered to the animals. Blood samples were collected in heparinized eppendorf tubes and kept on ice during the experiment. The plasma was separated by centrifugation (Remi, R4C Compact model, Mumbai, India) at 5000 rpm for 6 min and stored at -20°C until analysis.

#### *Extraction of sitagliptin from plasma*

Liquid-liquid extraction method was used to extract the


sitagliptin from the rat plasma. To an aliquot of 50 µL plasma, 50 µL of 1M sodium hydroxide and 1.5 mL ethyl acetate were added. Vortex mixed for 5 min on a Remi vortex mixer (Remi CM-101 Cyclomixer, Mumbai, India) and centrifuged (Remi, R4C Compact model, Mumbai, India) at 3000 rpm for 5 min. The supernatant (1.4 mL) was evaporated to dryness at 40°C under a stream of nitrogen. The dry residue was reconstituted in 200 µL of the mobile phase and a 20 µL aliquot was injected into HPLC system for analysis.

#### *Determination of sitagliptin by HPLC*

Sitagliptin concentrations in rat plasma were estimated by RP-HPLC as it has been described by Xiang et al. with minor modifications [43]. A Shimadzu HPLC system consisted of a pump (LC-20AT VP), C18 column (ODS Thermo Hypersil, 150 mm × 4.6 mm, 5.0 µm, Thermo Electron Corp, Beverly, MA) and a dual wavelength UV-visible detector (SPD-10A VP). LC solution software was used to collect and process the data. The mobile phase was composed of methanol: water (60:40, v/v) containing 10 mM tris and 10 mM triethylamine and titrated to pH 9.0 with 1M hydrochloric acid. The mobile phase was vacuum-filtered through 0.45 µm nylon Millipore membranes (Millipore, Billerica, MA), and degassed by ultrasonication (Remi, Mumbai, India) for 20 min before use. The flow of mobile phase was set at 1.0 mL/min and the injection volume was 20 µL. After equilibration with the solvent to obtain a stable baseline, aliquots of samples were injected. The absorbance of the eluent was monitored at 267nm. All the analyses were performed at 25.0 ± 0.5°C. The retention times of blank plasma and sitagliptin were obtained at 2.883 and 6.181 min, respectively (Fig. 1).

#### *Preparation of standard and stock solutions*

Sitagliptin stock solution was prepared by dissolving 1 mg in 10 mL of methanol (100 µg/mL) and stored in amber



**Figure 1.** Representative chromatograms of (A) blank plasma (2.883); (B) sitagliptin 2 µg/mL (6.181); (C) plasma (2.904) + sitagliptin 6 µg/mL (6.224); (D) plasma sample obtained from rats treated with 10 mg/kg of sitagliptin monitored at 267 nm.

glass containers at  $-20^{\circ}\text{C}$ . Calibration standards were prepared by spiking appropriate amounts of sitagliptin in 100  $\mu\text{L}$  of rat plasma. The final concentrations of calibration standards (0.1, 0.25, 0.5, 0.75, 1, 2, 4, 6, 8 and 10  $\mu\text{g}/\text{mL}$ ) were stored at  $2-8^{\circ}\text{C}$ . Calibration curves were constructed between the concentration and peak area. The lower limit of quantification was 0.75  $\mu\text{g}/\text{mL}$ .

#### Calculation of pharmacokinetic parameters

The plasma concentrations versus time data obtained from each individual rat were submitted to a non-compartmental pharmacokinetic analysis using Thermo Kinetics, Version 5.1 (Thermo Electron Corporation, Beverly, MA). The maximum plasma concentrations ( $C_{\text{max}}$ ) and times to achieve maximum plasma concentrations ( $t_{\text{max}}$ ) were obtained directly from the individual plasma concentration-time curves. Area under the plasma concentration-time curve from time zero to 24 h ( $\text{AUC}_{0-24}$ ) was calculated by the linear trapezoidal rule and  $\text{AUC}_{0-\infty}$  was determined by the following formula:

$$\text{AUC}_{0-\infty} = C_{\text{last}}/K_{\text{el}} + \text{AUC}_{0-24}$$

Where  $C_{\text{last}}$  is the last quantifiable concentration. The apparent total body clearance or oral clearance ( $\text{CL}/F$ ) was calculated as follows:

$$\text{CL}/F = \text{Dose}/\text{AUC}_{0-24}$$

The elimination half-life ( $t_{1/2}$ ) was calculated as  $0.693/K_{\text{el}}$  and the mean residence time (MRT) was calculated as follows:

$$\text{MRT} = \text{AUMC}_{0-\infty}/\text{AUC}_{0-\infty}$$

The area under the first moment curve ( $\text{AUMC}_{0-\infty}$ ) was calculated from the plasma concentration-time curve. The volume of distribution ( $V_z/F$ ) was estimated by means of  $V_z/F = \text{dose}/\text{AUC}_{0-\infty}/K_{\text{el}}$ . The apparent volume of distribution at steady state ( $V_{\text{ss}}/F$ ) was estimated by means of  $V_{\text{ss}}/F = \text{CL} \times \text{MRT}$ . The absolute oral bioavailability was determined by using following equation:

$$\text{Absolute bioavailability (AB\%)} = \frac{\text{AUC}_{0-24, \text{ oral}} \times \text{Dose}_{\text{ i.v.}}}{\text{AUC}_{0-24, \text{ i.v.}} \times \text{Dose}_{\text{ oral}}} \times 100$$

#### Ex vivo drug absorption studies using rat gut sacs

Rat everted and non-everted gut sacs (NEGS) are simple and useful *ex vivo* models to investigate the role of P-gp and CYP3A4 in drug disposition [44]. Both sacs were prepared using a method described previously by Pingili et al. in our laboratory with some modifications [45]. Healthy male wistar rats (weighing about 180-220 g,  $n = 3$ ) were deprived of food for 1 day and provided with only double-distilled water before the experiments. The rats were anesthetized with pentobarbital sodium (40 mg/kg, i. v.) and the small intestine was dissected out. The intestine was immediately rinsed with ice-cold Krebs-Ringer-Hensleit bicarbonate (KRH) buffer (7.8 g NaCl, 0.35 g KCl, 0.37 g  $\text{CaCl}_2$ , 1.37 g  $\text{NaHCO}_3$ , 0.32 g  $\text{NaH}_2\text{PO}_4$ , 0.02 g  $\text{MgCl}_2$ , 1.4 g glucose, pH 6.8) in order to discard the intestinal digesta and the distal

ileums (approximately 10 cm each) were taken for the study.

#### Effect of verapamil on the intestinal transport of sitagliptin

NEGS were used to evaluate the transport of sitagliptin from mucosal to serosal (M-to-S) side of the intestine. NEGS were loaded with 1mL of modified Krebs-Ringer (128.5 mM NaCl, 4.7 mM KCl, 3.3 mM  $\text{CaCl}_2$ , 2.3 mM  $\text{MgCl}_2$ , 16.3 mM  $\text{NaHCO}_3$ , 1.87 mM  $\text{NaH}_2\text{PO}_4$ , 7.8 mM glucose, pH 7.4) bicarbonate (KRB) buffer containing sitagliptin (50 $\mu\text{g}/\text{mL}$ ) in the presence or absence of known P-gp inhibitors (itraconazole and ketoconazole) and verapamil (50 $\mu\text{g}/\text{mL}$ ). Each sac was placed in individual 50 mL Erlenmeyer flasks contain 30 mL of oxygenated ( $\text{O}_2/\text{CO}_2$ ; 95:5) KRB and incubated at  $37^{\circ}\text{C}$  for 60 min in a shaker bath. Aliquots (150  $\mu\text{L}$ ) of serosal fluid were collected at 10, 20, 30, 40, 50 and 60 min and then replaced by the same volume of buffer. At the end of incubation, all samples were centrifuged (Remi, R4C Compact model, Mumbai, India) at 3500 rpm for 10 min and supernatants were analyzed for sitagliptin by RP-HPLC. Each experiment was triplicate. The same experiment was repeated with everted gut sacs (EGS) to study the transport of sitagliptin from serosal to mucosal (S-to-M) side of the intestine. The intestinal segments were everted using a stain less steel rod. The apparent permeability coefficient ( $P_{\text{app}}$ ), efflux ratio and net efflux were determined by the following formula:

$$\text{Apparent permeability coefficient (cm/s)} = \frac{dQ}{dt} \times \frac{1}{A \times C_0}$$

$$\text{Efflux ratio} = \frac{P_{\text{app}}(\text{serosal to mucosal})}{P_{\text{app}}(\text{mucosal to serosal})}$$

$$\text{Net efflux} = P_{\text{app}}(\text{serosal to mucosal}) - P_{\text{app}}(\text{mucosal to serosal})$$

Where  $dQ/dt$  ( $\mu\text{g}/\text{min}$ ) is the permeation rate of drug;  $A$  ( $\text{cm}^2$ ) is the surface area of the intestinal sac and  $C_0$  ( $\mu\text{g}/\text{mL}$ ) is the initial loading concentration.

#### In situ drug permeability (Single pass intestinal perfusion) studies

To further confirm the involvement of P-gp and CYP3A4 in the intestinal permeability of sitagliptin, single pass intestinal perfusion (*in situ*) experiment was performed as described previously by Yang et al. [46] and Neerati et al. [47]. In brief, healthy male wistar rats (weighing about 180-220 g,  $n = 6$ ) were maintained on a 12h light-dark cycle and fasted 16-20 h before each experiment. Animals were anaesthetized with an intravenous injection of pentobarbital sodium (40 mg/kg) and placed on a heated surface to maintain a normal body temperature. A midline incision was made on the abdomen and an ileal segment of approximately 13-15 cm was isolated using the ileo-caecal junction as a distal marker. The ileal segment was cannulated at both ends with flexible plastic tubing and rinsed with normal saline ( $37^{\circ}\text{C}$ ). The segment was infused with KRB buffer for 10 min at a flow rate of 1.0 mL/min using a syringe micro infusion pump (Aspire, Plenum tech Pvt. Ltd, Nagpur,

India). Subsequently, KRB buffer containing sitagliptin (50  $\mu\text{M}$ ) with or without 100  $\mu\text{M}$  verapamil was perfused through the intestinal segment at a constant flow rate of 0.25 mL/min for 90 min. Perfusate samples obtained from the outlet of the ileum were collected every 10 min. During the entire course of the experiment, care was taken to avoid disturbing the circulatory system, and the exposed segment was kept moist after cannulation with isotonic saline-moistened gauze (37°C). At the end of the experiment (i.e., following the last sample collection), the radius and length of the ileal segment were measured. The samples were stored at -20°C until HPLC analysis. The effective permeability coefficient ( $P_{\text{eff}}$ ) is a quantitative estimate of the rate of drug permeability across a membrane and is calculated based on the steady state concentration of drug in the collected perfusate as follows:

$$\text{Effective permeability coefficient} = \frac{-Q \times \ln(C_{\text{out}}/C_{\text{in}})}{2\pi rL}$$

Where  $Q$  is the perfusion flow rate through the ileal segment;  $C_{\text{out}}$  and  $C_{\text{in}}$  are the outlet and inlet concentrations of the drug, respectively;  $r$  is the radius of the perfused rat small intestine segment (0.19 cm); and  $L$  is the length of the perfused segment (15 cm). Intestinal absorption rate constant ( $K_a$ ) was calculated by using the following relationship:


$$\text{Intestinal absorption rate constant (K}_a) = \frac{Q_{\text{in}}(1 - C_{\text{out}}/C_{\text{in}})}{V_{\text{ps}}}$$

Where  $V_{\text{ps}}$  is the volume of perfused ileal segment ( $=r^2\pi L$ ).

## RESULTS

### Effect of verapamil on the oral pharmacokinetics of sitagliptin in vivo

The mean plasma concentration-time profiles of sitagliptin after oral administration of sitagliptin (10 mg/kg) alone and with verapamil (40 mg/kg) pretreatment are



**Figure 2.** Mean plasma concentration–time profiles of sitagliptin following an oral administration of sitagliptin (10 mg/kg) to rats with or without verapamil (40 mg/kg, oral) pretreatment (A) on 1<sup>st</sup> day; (B) on 15<sup>th</sup> day ( $n=6$ ). (●) SP (10 mg/kg); (▲) SP with verapamil (40 mg/kg) pretreatment. All values are Mean  $\pm$  SD. Bars represent the standard deviation. \* $p < 0.01$  when compared to the SP control group. SP, Sitagliptin Phosphate; VER, Verapamil. (Two-way ANOVA followed by Bonferroni post-tests to compare to each column to column).

**Table 1.** Pharmacokinetic parameters of sitagliptin phosphate (10 mg/kg) following an oral administration of sitagliptin to rats with or without verapamil (40 mg/kg) in SDS and MDS

Parameter	SDS ( $n=6$ )		MDS ( $n=6$ )	
	SP (10 mg/kg)	SP + VER (40 mg/kg)	SP (10 mg/kg)	SP + VER (40 mg/kg)
$C_{\text{max}}$ ( $\mu\text{g/mL}$ )	3.651 $\pm$ 0.855	6.120 $\pm$ 1.524***	3.953 $\pm$ 1.210	7.525 $\pm$ 1.550***
$\text{AUC}_{0-24}$ ( $\mu\text{g h/mL}$ )	25.092 $\pm$ 5.320	40.175 $\pm$ 6.625***	30.404 $\pm$ 6.235	45.266 $\pm$ 4.632***
$\text{AUC}_{0-\infty}$ ( $\mu\text{g h/mL}$ )	41.121 $\pm$ 7.740	69.830 $\pm$ 10.023***	61.146 $\pm$ 8.255	82.068 $\pm$ 9.624***
$t_{\text{max}}$ (h)	0.500 $\pm$ 0.000	0.500 $\pm$ 0.000 <sup>NS</sup>	0.500 $\pm$ 0.000	0.500 $\pm$ 0.000 <sup>NS</sup>
$t_{1/2}$ (h)	15.863 $\pm$ 4.261	20.935 $\pm$ 3.145**	25.569 $\pm$ 5.241	39.280 $\pm$ 2.547***
MRT (h)	23.954 $\pm$ 5.620	28.182 $\pm$ 3.251*	54.918 $\pm$ 7.158	66.055 $\pm$ 4.325***
CL/F (mL/h/kg)	0.103 $\pm$ 0.052	0.048 $\pm$ 0.016*	0.124 $\pm$ 0.055	0.024 $\pm$ 0.008*
Vz/F (mL/kg)	3.114 $\pm$ 0.563	1.113 $\pm$ 0.245**	1.668 $\pm$ 0.854	1.602 $\pm$ 0.472 <sup>NS</sup>
$V_{\text{ss}}$ /F (mL/kg)	2.906 $\pm$ 0.400	1.165 $\pm$ 0.366***	2.000 $\pm$ 0.700	1.460 $\pm$ 0.551 <sup>NS</sup>
AB (%)	1.161 $\pm$ 0.285	1.530 $\pm$ 0.186*	1.654 $\pm$ 0.166	2.136 $\pm$ 0.269*


SP, sitagliptin phosphate; VER, verapamil; SDS, single dose pharmacokinetic study; MDS, multiple dose pharmacokinetic study;  $C_{\text{max}}$ , peak plasma concentration;  $\text{AUC}_{0-24}$ , area under the plasma concentration–time curve from 0 h to 24 h;  $\text{AUC}_{0-\infty}$ , area under the plasma concentration–time curve from 0 h to infinity;  $t_{\text{max}}$ , time to reach peak plasma concentration;  $t_{1/2}$ , terminal half-life; MRT, mean residence time; CL/F, apparent total body clearance or oral clearance; Vz/F, apparent volume of distribution;  $V_{\text{ss}}$ , apparent volume of distribution at steady state; AB, absolute bioavailability. All values are mean  $\pm$  SD. \*\*\* $p < 0.001$ , \*\* $p < 0.01$ , \* $p < 0.05$ , <sup>NS</sup> $p > 0.05$  when compared to sitagliptin alone group (two-way ANOVA followed by Bonferroni post-tests to compare to each column to column).

depicted in Figure 2 (A, SDS and B, MDS). The pharmacokinetic parameters of sitagliptin are summarized in Table 1. Verapamil pretreatment significantly affected the pharmacokinetics of sitagliptin in both SDS and MDS. The sitagliptin plasma concentrations were significantly higher in the verapamil pretreated rats than rats treated with sitagliptin alone. The  $AUC_{0-24h}$ ,  $C_{max}$ , AB (%), MRT and  $t_{1/2}$  of sitagliptin were significantly increased by the verapamil pretreatment. The mean  $\pm$  SD  $AUC_{0-24h}$  of sitagliptin was significantly increased from  $25.092 \pm 5.320$  to  $40.175 \pm 6.625$   $\mu\text{g h/mL}$  (in SDS,  $p < 0.001$ ) and  $30.404 \pm 6.235$  to  $45.266 \pm 4.632$   $\mu\text{g h/mL}$  (in MDS,  $p < 0.001$ ),  $C_{max}$  increased from  $3.651 \pm 0.855$  to  $6.120 \pm 1.524$   $\mu\text{g/mL}$  (in SDS,  $p < 0.001$ ) and  $3.953 \pm 1.210$  to  $7.525 \pm 1.550$   $\mu\text{g/mL}$  (in MDS,  $p < 0.001$ ), AB (%) increased from  $1.161 \pm 0.285$  to  $1.530 \pm 0.186$  (in SDS,  $p < 0.05$ ) and  $1.654 \pm 0.166$  to  $2.136 \pm 0.269$  (in MDS,  $p < 0.05$ ), MRT increased from  $23.954 \pm 5.620$  to  $28.182 \pm 3.251$  h (in SDS,  $p < 0.05$ ) and  $54.918 \pm 7.158$  to  $66.055 \pm 4.325$  h (in MDS,  $p < 0.001$ ),  $t_{1/2}$  increased

from  $15.863 \pm 4.261$  to  $20.935 \pm 3.145$  h (in SDS,  $p < 0.01$ ) and  $25.569 \pm 5.241$  to  $39.280 \pm 2.547$  h (in MDS,  $p < 0.001$ ). The verapamil effect was not statistically significant on the  $t_{max}$  of sitagliptin. The  $V_z/F$ ,  $CL/F$  and  $V_{ss}/F$  of sitagliptin were significantly decreased in verapamil pretreated rats when compared to sitagliptin control group. The mean  $\pm$  SD  $V_z/F$  of sitagliptin was decreased from  $3.114 \pm 0.563$  to  $1.113 \pm 0.245$  mL/kg (in SDS,  $p < 0.01$ ) and  $1.668 \pm 0.854$  to  $1.602 \pm 0.472$  mL/kg (in MDS,  $p > 0.05$ ),  $CL/F$  decreased from  $0.103 \pm 0.052$  to  $0.048 \pm 0.016$  mL/kg (in SDS,  $p < 0.05$ ) and  $0.124 \pm 0.055$  to  $0.024 \pm 0.008$  mL/kg (in MDS,  $p < 0.05$ ),  $V_{ss}$  decreased from  $2.906 \pm 0.400$  to  $1.165 \pm 0.366$  mL/kg (in SDS,  $p < 0.05$ ) and  $2.000 \pm 0.700$  to  $1.460 \pm 0.551$  mL/kg (in MDS,  $p > 0.05$ ).

#### Effect of verapamil on the intravenous pharmacokinetics of sitagliptin in vivo

The mean plasma concentration versus time curves of sitagliptin after intravenous administration of sitagliptin (5



**Figure 3.** Mean plasma concentration–time profiles of sitagliptin following an intravenous administration of sitagliptin (10 mg/kg) to rats with or without verapamil (40 mg/kg, oral) pretreatment (A) on 1<sup>st</sup> day; (B) on 15<sup>th</sup> day (n=6). (●) SP (10 mg/kg); (■) SP with verapamil (40 mg/kg) pretreatment. All values are Mean  $\pm$  SD. Bars represent the standard deviation. \* $p < 0.01$  when compared to the SP control group. SP, Sitagliptin Phosphate; VER, Verapamil. (Two-way ANOVA followed by Bonferroni post-tests to compare to each column to column).

**Table 2.** Pharmacokinetic parameters of sitagliptin (5 mg/kg) following an intravenous administration of sitagliptin to rats with or without verapamil (40 mg/kg) in SDS and MDS

Parameter	SDS		MDS	
	SP (5 mg/kg)	SP + VER (40 mg/kg)	SP (10 mg/kg)	SP + VER (40 mg/kg)
$C_{max}$ ( $\mu\text{g/mL}$ )	$8.657 \pm 2.632$	$11.524 \pm 3.415^*$	$10.771 \pm 2.400$	$15.141 \pm 3.482^{**}$
$AUC_{0-24}$ ( $\mu\text{g h/mL}$ )	$1318.990 \pm 120.451$	$2079.980 \pm 251.362^{***}$	$1521.115 \pm 100.356$	$3613.370 \pm 256.400^{***}$
$AUC_{0-\infty}$ ( $\mu\text{g h/mL}$ )	$1509.766 \pm 146.844$	$2281.145 \pm 235.478^{***}$	$1706.612 \pm 154.262$	$2178.000 \pm 325.162^{***}$
$t_{max}$ (h)	$5.000 \pm 0.000$	$5.000 \pm 0.000^{NS}$	$5.000 \pm 0.000$	$5.000 \pm 0.000^{NS}$
$t_{1/2}$ (min)	$183.225 \pm 20.415$	$201.473 \pm 30.487^*$	$189.379 \pm 17.255$	$377.183 \pm 40.236^{**}$
MRT (h)	$263.294 \pm 40.855$	$307.790 \pm 50.325^*$	$279.672 \pm 30.480$	$475.332 \pm 62.354^{***}$
$CL/F$ (mL/h/kg)	$0.003 \pm 0.0004$	$0.001 \pm 0.002^{NS}$	$0.0015 \pm 0.0001$	$0.0011 \pm 0.0001^{NS}$
$V_z/F$ (mL/kg)	$0.712 \pm 0.250$	$0.364 \pm 0.052^{***}$	$0.952 \pm 0.240$	$0.436 \pm 0.040^{***}$
$V_{ss}/F$ (mL/kg)	$0.651 \pm 0.324$	$0.276 \pm 0.063^{**}$	$0.865 \pm 0.300$	$0.315 \pm 0.050^{**}$


SP, sitagliptin phosphate; VER, verapamil; SDS, single dose pharmacokinetic study; MDS, multiple dose pharmacokinetic study;  $C_{max}$ , peak plasma concentration;  $AUC_{0-24}$ , area under the plasma concentration–time curve from 0 h to 24 h;  $AUC_{0-\infty}$ , area under the plasma concentration–time curve from 0 h to infinity;  $t_{max}$ , time to reach peak plasma concentration;  $t_{1/2}$ , terminal half-life; MRT, mean residence time;  $CL/F$ , apparent total body clearance or oral clearance;  $V_z/F$ , apparent volume of distribution;  $V_{ss}$ , apparent volume of distribution at steady state. All values are mean  $\pm$  SD. \*\*\* $p < 0.001$ , \*\* $p < 0.01$ , \* $p < 0.05$ , <sup>NS</sup> $p > 0.05$  when compared to sitagliptin alone group (two-way ANOVA followed by Bonferroni post-tests to compare to each column to column).

mg/kg) alone and with verapamil (40 mg/kg) pretreatment are shown in Figure 3 (A, SDS and B, MDS). The pharmacokinetic parameters of sitagliptin are presented in Table 2. Verapamil pretreatment was significantly affected the pharmacokinetics of sitagliptin (intravenous) in both SDS and MDS. When sitagliptin was administered with verapamil pretreatment, the plasma concentrations of sitagliptin were significantly higher compared to the sitagliptin control (Fig. 3). The mean  $\pm$  SD  $AUC_{0-24h}$  of sitagliptin was significantly increased from  $1318.990 \pm 120.451$  to  $2079.980 \pm 251.362$   $\mu\text{g h/mL}$  (in SDS,  $p < 0.001$ ) and  $1521.115 \pm 100.356$  to  $3613.370 \pm 256.400$   $\mu\text{g h/mL}$  (in MDS,  $p < 0.001$ ),  $C_{max}$  increased from  $8.657 \pm 2.632$  to  $11.524 \pm 3.415$   $\mu\text{g/mL}$  (in SDS,  $p < 0.05$ ) and  $10.771 \pm 2.400$  to  $15.141 \pm 3.482$   $\mu\text{g/mL}$  (in MDS,  $p < 0.01$ ), MRT increased from  $263.294 \pm 40.855$  to  $307.790 \pm 50.325$  h (in SDS,  $p < 0.05$ ) and  $279.672 \pm 30.480$  to  $475.332 \pm 62.354$  h (in MDS,  $p < 0.001$ ),  $t_{1/2}$  increased from  $183.225 \pm 20.415$


$201.473 \pm 30.487$  h (in SDS,  $p < 0.05$ ) and  $189.379 \pm 17.255$  to  $377.183 \pm 40.236$  h (in MDS,  $p < 0.01$ ). Verapamil had no significant effect on the  $t_{max}$  of sitagliptin in both SDS and MDS. The  $V_z/F$ ,  $CL/F$  and  $V_{ss}/F$  of sitagliptin were significantly decreased in verapamil pretreated rats when compared to sitagliptin control group. The mean  $\pm$  SD  $CL/F$  of sitagliptin was decreased from  $0.003 \pm 0.0004$  to  $0.001 \pm 0.002$  mL/kg (in SDS,  $p > 0.05$ ) and  $0.0015 \pm 0.0001$  to  $0.0011 \pm 0.0001$  mL/kg (in MDS,  $p > 0.05$ ),  $V_z/F$  decreased from  $0.712 \pm 0.250$  to  $0.364 \pm 0.052$  mL/kg (in SDS,  $p < 0.001$ ) and  $0.952 \pm 0.240$  to  $0.436 \pm 0.040$  mL/kg (in MDS,  $p < 0.001$ ),  $V_{ss}/F$  decreased from  $0.651 \pm 0.324$  to  $0.276 \pm 0.063$  mL/kg (in SDS,  $p < 0.01$ ) and  $0.865 \pm 0.300$  to  $0.315 \pm 0.050$  mL/kg (in MDS,  $p < 0.01$ ).

#### Effect of verapamil on the intestinal transport of sitagliptin *ex vivo*

The apparent permeability coefficient ( $P_{app}$ ), net efflux



**Figure 4.** The effect of verapamil on the apparent permeability coefficient ( $P_{app}$ ) of sitagliptin phosphate from (A) mucosal to serosal side of non-everted gut sacs; (B) serosal to mucosal side of everted gut sacs *ex vivo* ( $n=3$ ). All values are mean  $\pm$  SD.  $**p < 0.01$ ,  $*p < 0.05$ ,  $^{NS}P > 0.05$  when compared to sitagliptin alone group (two-way ANOVA followed by Bonferroni post-tests to compare to each column to column). SP, Sitagliptin Phosphate; VER, Verapamil; ITR, Itraconazole; KTZ, Ketoconazole.



**Figure 5.** The effect of verapamil on the (A) Net efflux and (B) efflux ratios of sitagliptin using rat gut sacs *ex vivo* ( $n=3$ ). All values are mean  $\pm$  SD.  $***p < 0.001$ ,  $*p < 0.05$ ,  $^{NS}P > 0.05$  when compared to sitagliptin alone group (two-way ANOVA followed by Bonferroni post-tests to compare to each column to column). SP, Sitagliptin Phosphate; VER, Verapamil; ITR, Itraconazole; KTZ, Ketoconazole.

and efflux ratio values were calculated to evaluate the effect of verapamil on the P-gp mediated intestinal absorption of sitagliptin using *non-everted rat gut sac* absorption models are shown in Figures 4 & 5. The results indicated that known P-gp inhibitors (itraconazole and ketoconazole) and verapamil significantly affected the  $P_{app}$  of sitagliptin from mucosal to serosal (M-to-S), net efflux and efflux ratio. But the effect was statistically not significant on the  $P_{app}$  of sitagliptin from serosal to mucosal (S-to-M) when compared to sitagliptin alone. The sitagliptin mean  $\pm$  SD  $P_{app}$  from M-to-S increased significantly from  $4.479 \pm 0.989 (\times 10^{-4} \text{ cm/s})$  to  $7.519 \pm 1.122 (\times 10^{-4} \text{ cm/s})$  and  $6.675 \pm 1.567 (\times 10^{-4} \text{ cm/s})$  to  $9.381 \pm 1.445 (\times 10^{-4} \text{ cm/s})$  with verapamil at 60 and 90 min, respectively. The mean  $\pm$  SD net efflux of sitagliptin was decreased from  $1.623 \pm 0.690$  to  $0.318 \pm 0.056$  (at 60 min,  $P < 0.05$ ) and  $2.938 \pm 0.893$  to  $1.509 \pm 0.682$  (at 90 min,  $p < 0.05$ ) in the presence of verapamil. Verapamil was also significantly decreased the mean  $\pm$  SD efflux ratio of sitagliptin from  $1.415 \pm 0.483$  to  $1.038 \pm 0.452$  ( $p < 0.05$ ) and  $1.529 \pm 0.409$  to  $1.213 \pm 0.218$  ( $p < 0.05$ ) at 60 and 90 min, respectively.

#### Effect of verapamil on the intestinal permeability of sitagliptin *in situ*


The effective permeability coefficient ( $P_{eff}$ ) and intestinal absorption rate constant ( $K_a$ ) values were determined to further evaluate the influence of verapamil on the P-gp mediated intestinal permeation of sitagliptin using *single pass intestinal perfusion* models are depicted in Figure 6. The results showed that the P-gp inhibitors (itraconazole and ketoconazole) and verapamil significantly increased the  $P_{eff}$  and  $K_a$  of sitagliptin when compared to sitagliptin alone. The mean  $\pm$  SD  $P_{eff}$  of sitagliptin was significantly increased from  $0.810 \pm 0.050 (\times 10^{-3} \text{ cm/s})$  to  $1.986 \pm 0.514 (\times 10^{-3} \text{ cm/s})$  and  $1.163 \pm 0.150 (\times 10^{-3} \text{ cm/s})$  to  $2.739 \pm 0.452 (\times 10^{-3} \text{ cm/s})$  in the presence of verapamil at 60 and 90 min, respectively. The sitagliptin mean  $\pm$  SD  $K_a$  was increased

from  $0.102 \pm 0.025$  to  $0.187 \pm 0.051$  (at 60 min,  $p < 0.05$ ) and  $0.113 \pm 0.032$  to  $0.196 \pm 0.045$  (at 90 min,  $p < 0.05$ ) with verapamil.

#### DISCUSSION

Oral administration is the most common route for drug administration. Orally-administered drugs have to cross the gastrointestinal epithelial cell membrane and are then transported into the port vein to the liver and eventually into the systemic circulation to exhibit their pharmacological effects [48]. P-gp and CYP3A4 are constitutively expressed in intestinal epithelial cells and form a significant barrier to the absorption of some orally administered drugs like sitagliptin and verapamil. Moreover, P-gp may act with CYP3A4 to increase the metabolism of sitagliptin in the intestine [49, 50]. Previous studies reported that verapamil inhibited the P-gp and CYP3A4-mediated metabolism of levobupivacaine, lidocaine [29], imipramine [30], doxorubicin [51], atorvastatin [31], carbamazepine [52], simvastatin, nilotinib [32], cerivastatin [33], darifenacin, ethosuximide [34], cyclosporine A [53], prazosin [54], digoxin [55], quinine, quinidine and zolpidem [35]; thereby increased their plasma concentrations. In the present study also, the plasma concentration of sitagliptin was significantly increased in SDS and MDS may be due to inhibition of P-gp and CYP3A4.

Risperidone is a substrate of P-gp. The  $C_{max}$  of risperidone was significantly increased by 1.69-fold with verapamil, a strong P-gp inhibitor in male healthy volunteers [56]. Another study revealed that the  $C_{max}$  of aliskiren (P-gp substrate) was increased by 1.92-fold when co-administered with verapamil (P-gp inhibitor) in healthy participants [57]. In the present study, verapamil pretreatment significantly increased the  $C_{max}$  of sitagliptin by 1.67 and 1.90-folds in SDS and MDS, respectively. The  $C_{max}$  of sitagliptin was increased by 1.33-fold (SDS) and 1.40-folds (MDS) after intravenous administration of sitagliptin with or without oral



**Figure 6.** The effect of verapamil on the (A) effective permeability coefficient ( $P_{eff}$ ) and (B) intestinal absorption rate constant ( $K_a$ ) of sitagliptin in the presence or absence of verapamil using rat gut sacs *in situ* ( $n=3$ ). All values are mean  $\pm$  SD.  $**p < 0.01$ ,  $*p < 0.05$ ,  $^{NS}P > 0.05$  when compared to sitagliptin alone group (two-way ANOVA followed by Bonferroni post-tests to compare to each column to column). SP, Sitagliptin Phosphate; VER, Verapamil; ITR, Itraconazole; KTZ, Ketoconazole.



pretreatment of verapamil. Verapamil is a P-gp inhibitor may increase the exposure of afatinib (P-gp substrate) when co-administered with afatinib. Aprepitant is a CYP3A4 substrate. Coadministration of daily oral aprepitant (230 mg, or 1.8 times the recommended single dose) with verapamil (moderate CYP3A4 inhibitor) increased the aprepitant AUC 2-fold [58]. The total AUC of risperidone was increased 1.59-fold by verapamil (strong P-gp inhibitor) in male healthy volunteers [56]. In the present study also the AUC<sub>0-24</sub> of sitagliptin was significantly increased by 1.60-fold (SDS) and 1.48-fold (MDS) with verapamil pretreatment. The AUC<sub>0-24</sub> of sitagliptin was also increased by 1.57-fold (SDS) and 2.37-fold (MDS) after intravenous administration of sitagliptin with verapamil pretreatment.

Aliskiren is a substrate for P-gp transporter: The  $t_{max}$  of aliskiren was not affected significantly when co-administer with verapamil (P-gp inhibitor) in healthy participants [57]. Erik et al. (2014) reported that the  $t_{max}$  of fexofenadine (P-gp substrate) was not significantly affected by the concomitant use of verapamil in pigs [59]. Verapamil pretreatment had no significant effect on the  $t_{max}$  of sitagliptin in the present study. Colchicine is a P-gp and CYP3A4 substrate and verapamil is a P-gp and CYP3A4 inhibitor. Half-life of colchicine in the serum was found to be 8-fold higher than the normal when used concomitantly with verapamil [60]. In the present study, the sitagliptin half life was 1.31-fold (SDS) and 1.53-fold (MDS) higher in verapamil pretreatment group compared to sitagliptin control group. The half life of sitagliptin was increased significantly after intravenous administration in both SDS and MDS. Verapamil increased the MRT of moxidectin from  $17.7 \pm 2.56$  to  $19.7 \pm 4.24$  days in sheep [61]. In another study, the MRT of irinotecan was increased 1.42-fold by verapamil in rats [62]. The MRT of sitagliptin was significantly increased by 1.17-fold (SDS) and 1.20-fold (MDS) in the present study. The same results were observed in the intravenous pharmacokinetic study also.

Verapamil inhibits the CYP3A4 metabolism and decreases clearance of imipramine [30], Doxorubicin [51], Theophylline [63] and Caffeine [64]. Paclitaxel is a substrate of P-gp. The clearance of paclitaxel was decreased 0.5-fold by verapamil (P-gp inhibitor) in women with breast cancer [65]. The clearance of Polyphyllin I was significantly decreased 0.17-fold with verapamil treatment [66]. Daunomycin clearance was decreased 0.10-fold when co-administered with verapamil in rats [67]. In the present study also, verapamil significantly decreased the clearance of sitagliptin by 0.33-fold (SDS) and 0.19-fold (MDS). The clearance of sitagliptin was significantly decreased by 0.33-fold (SDS) and 0.73-fold (MDS) after intravenous administration of sitagliptin in verapamil pretreated rats. The Vz/F of ivermectin (P-gp substrate) was significantly decreased by 0.58-fold with verapamil in sheep [68]. The Vz/F of sitagliptin was significantly decreased in the present both (oral and intravenous) studies.

The Vss/F of paclitaxel (P-gp substrate) was significantly decreased by 0.49-fold when co-administered with verapamil

(P-gp inhibitor) in women with breast cancer [65]. Irinotecan Vss/F was also decreased by 0.23-fold when concomitantly administered with verapamil in rats [62]. In the present study also, the Vss/F of sitagliptin was significantly decreased by 0.40-fold (SDS) after oral administration of sitagliptin to verapamil pretreated rats. In intravenous study also, sitagliptin Vss/F was significantly decreased. Polyphyllin I (PPI), one of the steroidal saponins in Paris polyphylla, is a promising natural anticancer candidate but its oral bioavailability is poor due to P-gp. Verapamil was greatly enhances the bioavailability of PPI through the inhibition of P-gp [66]. The absolute bioavailability of paclitaxel was increased 1.77-fold with verapamil (P-gp inhibitor) in rats [69]. Another study reported that the irinotecan bioavailability was increased by 4.33-fold with verapamil co-administration in rats [62]. The oral bioavailability of sitagliptin was significantly increased by 1.31-fold (SDS) and 1.29-fold (MDS) in rats pretreated with verapamil.

FB2 is a promising Abl/Src dual tyrosine kinase inhibitor. Its bioavailability is poor due to the P-gp mediated active efflux and first-pass metabolism in the rat intestine. The  $P_{app}$  (A-B) of FB2 was increased 28.39-fold and the  $P_{app}$  (B-A) of FB2 was decreased 0.86-fold with 100  $\mu$ M verapamil in MDCK-MDR1 cell model. The efflux ratio was decreased from 111.71 to 3.37 and the net efflux was decreased from 46.50 to 2.64 with 100  $\mu$ M verapamil in the same model [70]. Verapamil also decreased the efflux ratio of etoposide from 4.01 to 1.33 in everted rat gut sac model [71]. In the presence of 20  $\mu$ M cyclosporin A, the efflux ratio value of aconitine was reduced from 8.17 to 0.99 by increasing the AP-BL and reducing the BL-AP AC fluxes. Verapamil at 100  $\mu$ M exhibited an effect similar to that of cyclosporin A; however, the decrease in the BL-AP direction was greater, and the increase in the reverse direction was lower. The efflux ratio value of AC at 2 h decreased to 1.31 upon co-incubation with verapamil [72]. In the present study, the  $P_{app}$  (M-to-S) of sitagliptin was significantly increased and the  $P_{app}$  (S-to-M) of sitagliptin was slightly increased (statistically insignificant) with 50  $\mu$ g/mL verapamil in rat gut sac model. The efflux ratio was decreased from 1.529 to 1.213 and the net efflux was decreased from 2.938 to 1.509 with 50  $\mu$ g/mL verapamil in the same model.

The coperfusion of verapamil (100  $\mu$ M) and aconitine (5  $\mu$ M) resulted in a significant increase in intestinal permeability (12.9-fold) of aconitine in situ single-pass perfusion experiment in the rat ileum [72]. The effective intestinal permeability ( $P_{eff}$ ) of fexofenadine (P-gp substrate) was increased 5-fold with verapamil, a P-gp inhibitor in porcine model [73]. In the present study, the  $P_{eff}$  of sitagliptin was significantly increased by 2.35-fold with verapamil in situ single-pass perfusion experiment in the rat ileum. The intestinal absorption rate constant ( $K_a$ ) of fexofenadine (P-gp substrate) was increased by 1.76-fold with verapamil, a P-gp inhibitor in porcine model [73]. The  $K_a$  of paclitaxel was increased 1.17-fold with verapamil (P-gp inhibitor) pretreatment in rats [69]. In the present study also, the  $K_a$  of

sitagliptin was increased by 1.73-fold with verapamil in the same model.

### CONCLUSION

Verapamil pretreatment significantly increased the bioavailability and affected the pharmacokinetics of sitagliptin in rats might be through the inhibition of P-gp and CYP3A4. Further studies are needed to confirm this interaction at cellular level using P-gp and CYP3A4 over expressed cell lines and in human subjects.

### ACKNOWLEDGEMENTS

The results described in this paper were part of student thesis. This study was supported by Siddhartha Academy of General and Technical Education (SAGTE). The authors are grateful to N. Venkateswarlu, President and P. Lakshmana Rao, Secretary of SAGTE for providing necessary facilities.

### CONFLICT OF INTEREST

The authors declare that this research does not have any conflict of interest with anyone or any Institute.

### REFERENCES

- Gang Z, Jiangeng H, Kewen X, Luqin S, Gao L. Enhanced intestinal absorption of etoposide by self-microemulsifying drug delivery systems: Roles of P-glycoprotein and cytochrome P450 3A inhibition. *Eur J Pharmac Sci* 2013;50:429-439.
- Jorg K, Fabian M, Martin FF. Transporters and Drug-Drug Interactions: Important Determinants of Drug Disposition and Effects. *Pharmacol Rev* 2013;65:944-966.
- Fenner KS, Troutman MD, Kempshall S, Cook JA, Ware JA, Smith DA, et al. Drug-Drug Interactions Mediated Through P-Glycoprotein: Clinical Relevance and In Vitro-In Vivo Correlation Using Digoxin as a Probe Drug. *Clin Pharmacol Therapeut* 2009;85:173-181.
- Potschka H. Targeting regulation of ABC efflux transporters in brain diseases: a novel therapeutic approach. *Pharmacol Ther* 2010;125:118-127.
- Andrew F, Peter P. P-glycoprotein and its role in drug-drug interactions. *Aust Prescr* 2014;37:137-139
- Jef S, Pieter A, Patrick A. Site dependent intestinal absorption of darunavir and its interaction with ketoconazole. *Eur J Pharmac Sci* 2013;49:51-55.
- Giacomini KM, Huang SM, Tweedie DJ, Benet LZ, Brouwer KL. Membrane transporters in drug development. *Nat Rev Drug Discov* 2010;9:215-236.
- Amanda NL, Samuel DJ. The Comorbidities of Diabetes and Hypertension: Mechanisms and Approach to Target Organ Protection. *J Clin Hypertens (Greenwich)* 2011;13:244-251.
- George LB, James RS. ASH Position Paper: Treatment of Hypertension in Patients With Diabetes-An Update. *J Clin Hypertens (Greenwich)* 2008;10:707-713.
- Verspohl EJ. Novel therapeutics for type 2 diabetes: incretin hormone mimetics (glucagon-like peptide-1 receptor agonists) and dipeptidyl peptidase-4 inhibitors. *Pharmacol Ther* 2009;124:113-138.
- Yukiko M et al. Voglibose in patients with type 2 diabetes inadequately controlled via diet and/or a single oral antihyperglycemic agent: a multicenter, randomized trial. *BMJ Open Diabetes Res Care* 2016;4:e000190.
- El-Agamy DS, Abo-Haded HM, Elkablawy MA. Cardioprotective effects of sitagliptin against doxorubicin-induced cardiotoxicity in rats. *Exp Biol Med* 2016;1535370216643418.
- Andre JS. Cardiovascular effects of gliptins. *Nature Rev Cardiol* 2013;10:73-84.
- Vaghasiya J, Sheth N, Bhalodia Y, Manek R. Sitagliptin protects renal ischemia reperfusion induced renal damage in diabetes. *Regul Pept* 2011;166:48-54.
- Tsai TH, Sun CK, Su CH, Sung PH, Chua S, Zhen YY, et al. Sitagliptin attenuated brain damage and cognitive impairment in mice with chronic cerebral hypo-perfusion through suppressing oxidative stress and inflammatory reaction. *J Hypertens* 2015;33:1001-1013.
- Domenic AS. Pharmacotherapy Review: Calcium Channel Blockers. *J Clin Hyperten* 2006;8:53-56.
- Wing LMH. Calcium channel antagonists. *Aust Prescr* 1997;20:5-8.
- Elkayam U, Shotan A, Mehra A, Ostrzega E. Calcium channel blockers in heart failure. *J Am Coll Cardiol* 1993;22:139A-144A.
- Colucci WS, Fifer MA, Lorell BH, Wynne J. Calcium channel blockers in congestive heart failure: theoretic considerations and clinical experience. *Am J Med* 1985;78(2B):9-17.
- France P, Gaetan J, Louis D. Cardioprotective Efficacy of Verapamil and Mibefradil in Young UM-X7.1 Cardiomyopathic Hamsters. *Cardiovasc Drugs Ther* 1999;13:525-530.
- Kedziora KK, Szram S, Kornatowski T, Szadujkis SL, Kedziora J, Bartosz G. The effect of verapamil on the antioxidant defence system in diabetic kidney. *Clin Chim Acta* 2002;322:105-112.
- Yuxin L, Yi CL, Li Q, Fulton TC, Belinda W, Hui LC, et al. Verapamil Protects Dopaminergic Neuron Damage through a Novel Anti-inflammatory Mechanism by Inhibition of Microglial Activation. *Neuropharmacology* 2011;60:373-380.
- Zou Z, Xu FY, Wang L, An MM, Zhang H, Shi XY. Antihypertensive and renoprotective effects of trandolapril/verapamil combination: a meta-analysis of randomized controlled trials. *J Hum Hypertens* 2011;25: 203-210.
- Basim ASM, Amira MAY. Protective effects of fish oil, allopurinol, and verapamil on hepatic ischemia-reperfusion injury in rats. *J Nat Sci Biol Med* 2015;6:351-355.
- Skalskii SV. Effect of verapamil on production of anti-inflammatory cytokines by peritoneal mononuclears. *Vestn Ross Akad Med Nauk* 2010;3:33-35.
- Ganesh J, Viswanathan V. Management of diabetic hypertensives. *Indian J Endocr Metab* 2011;15:374-379.
- Wessler JD, Grip LT, Mendell J et al. The P-glycoprotein transport system and cardiovascular drugs. *J Am Coll Cardiol* 2013;61:2495-2502.
- Bent HH, Odd GN. In vitro Inhibition of CYP3A4 Metabolism and P-Glycoprotein-Mediated Transport by Trade Herbal Products. *Basic Clin Pharmacol Toxicol* 2008;102:466-475.
- Naguib M, Magboul MMA, Samarkandi AH, Attia M. Adverse effects and drug interactions associated with local and regional anesthesia. *Drug Safety* 1998;18:221-250.
- Hermann DJ, Krol TF, Dukes GE, Hussey EK, Danis M, Han YH, et al. Comparison of verapamil, diltiazem, and labetalol on the bioavailability and metabolism of imipramine. *J Clin Pharmacol* 1992;32:176-183.
- Tobert JA, Vega JM, Dobrinska M. Calcium channel blocker-simvastatin interaction. *Clin Pharmacol Ther* 1999;65:583-585.
- Mousa O, Brater DC, Sunblad KJ, Hall SD. The interaction of diltiazem with simvastatin. *Clin Pharmacol Ther* 2000;67:267-274.
- Williams D, Feely J. Pharmacokinetic-pharmacodynamic drug interactions with HMG-CoA reductase inhibitors. *Clin Pharmacokinet* 2002; 41:343-370.
- Bachmann K, He Y, Sarver JG, Peng N. Characterization of the cytochrome P450 enzymes involved in the in vitro metabolism of ethosuximide by human hepatic microsomal enzymes. *Xenobiotica* 2003;33:265-276.
- Hansten P, Horn J. The Top 100 Drug Interactions: A Guide to Patient Management. includes table of CYP450 and drug transporter substrates and modifiers (appendices). H & H Publications, LLP 2014 edition.
- Chu XY, Bleasby K, Yabut J, Cai X, Chan GH, Hafey MJ, et al. Transport of the dipeptidyl peptidase-4 inhibitor sitagliptin by human organic anion transporter 3, organic anion transporting polypeptide 4C1, and multidrug resistance P-glycoprotein. *J Pharmacol Exp Ther* 2007;321:673-683.

37. Krishna R, Bergman A, Larson P, Cote J, Lasseter K, Dilzer S, et al. Effect of a single cyclosporine dose on the single-dose pharmacokinetics of sitagliptin (MK-0431), a dipeptidyl peptidase-4 inhibitor, in healthy male subjects. *J Clin Pharmacol* 2007;47:165-174.
38. Pingili RB, Pawar AK, Challa SR. Systemic exposure of Paracetamol (acetaminophen) was enhanced by quercetin and chrysin co-administration in Wistar rats and in vitro model: risk of liver toxicity. *Drug Dev Ind Pharm* 2015;41:1793-1800.
39. Surya Sandeep M, Sridhar V, Puneeth Y, Ravindra Babu P, Naveen Babu K. Enhanced oral bioavailability of felodipine by naringenin in Wistar rats and inhibition of P-glycoprotein in everted rat gut sacs in vitro. *Drug Dev Ind Pharm* 2014;40:1371-1377.
40. Sridhar V, Surya Sandeep M, Ravindra Babu P, Naveen Babu K. Evaluation of First-Pass Cytochrome P4503A (CYP3A) and P-glycoprotein Activities Using Felodipine and Hesperetin in Combination in Wistar Rats and Everted Rat Gut Sacs in Vitro. *Phytother Res* 2014;28:699-705.
41. Zhiqiang L, Yang L, Baosheng Z, Mingmin T, Honghuan D, Lei Z, et al. Ex vivo and in situ approaches used to study intestinal absorption. *J Pharmacol Toxicol Methods* 2013;68:208-216.
42. Jef S, Pieter A, Patrick A. Site dependent intestinal absorption of darunavir and its interaction with ketoconazole. *Eur J Pharmac Sci* 2013; 49:51-55.
43. Jiy XF, De Zhou T, Lu W. A high performance liquid chromatography method for the quantitative determination assay of sitagliptin in rat plasma and its application in pharmacokinetics study. *J Chin Pharmac Sci* 2011;20:63-69.
44. Kumar KK, Priyanka L, Gnananath K, Babu PR, Sujatha S. Pharmacokinetic drug interactions between apigenin, rutin and paclitaxel mediated by P-glycoprotein in rats. *Eur J Drug Metab Pharmacokinet* 2015; 40: 267-276.
45. Pingili R, Vemulapalli S, Mullapudi SS, Nuthakki S, Pendyala S, Kilaru N. Pharmacokinetic interaction study between flavanones (hesperetin, naringenin) and rasagiline mesylate in wistar rats. *Drug Dev Ind Pharm* 2016;42:1110-1117.
46. Cuiping Y, Tianhong Z, Zheng L, Liang X, Fei L, Jinxiu R, et al. P-glycoprotein is responsible for the poor intestinal absorption and low toxicity of oral aconitine: In vitro, in situ, in vivo and in silico studies. *Toxicol Appl Pharmacol* 2013;273:561-568.
47. Neerati P, Ganji D, Bedada SK. Study on in situ and in vivo absorption kinetics of phenytoin bismodulating P-glycoprotein with verapamil in rats. *Eur J Pharm Sci* 2011;44:27-31.
48. Hai MF, Jian MX, Qiao MEI, Lei DIAO, Mo LC, Juan JIN, et al. Involvement of cytochrome P450 3A4 and P-glycoprotein in first-pass intestinal extraction of omeprazole in rabbits. *Acta Pharmacol Sin* 2009;30: 566-572.
49. Kunta JR, Sinko PJ. Intestinal drug transporters: in vivo function and clinical importance. *Curr Drug Metab* 2004;5:109-124.
50. Chan LM, Lowes S, Hirst BH. The ABCs of drug transport in intestine and liver: efflux proteins limiting drug absorption and bioavailability. *Eur J Pharm Sci* 2004;21:25-51.
51. Kerr DJ, Graham J, Cummings J. The effect of verapamil on the pharmacokinetics of adriamycin. *Cancer Chemother Pharmacol* 1986;18: 239-242.
52. Macphee GJ, McInnes GT, Thompson GG et al. Verapamil potentiates carbamazepine neurotoxicity: a clinically important inhibitory interaction. *Lancet* 1986;1:7003.
53. Martin JE, Daoud AJ, Schroeder TJ, First MR. The clinical and economic potential of cyclosporin drug interactions. *Pharmacoeconomics* 1999; 15:317-337.
54. Elliott HL, Meredith PA, Campbell L, Reid JL. The combination of prazosin and verapamil in the treatment of essential hypertension. *Clin Pharmacol Ther* 1988;43:554-560.
55. Jochen K, Stefan M. Effects of verapamil on pharmacokinetics and pharmacodynamics of digitoxin in patients. *Am Heart J* 1985;110: 1245.
56. Taku N, Norio YF, Manabu S, Tomonori T, Sunao K. Effect of verapamil on pharmacokinetics and pharmacodynamics of risperidone: In vivo evidence of involvement of P-glycoprotein in risperidone disposition. *Clin Pharmacol Ther* 2005;78:43-51.
57. Sam R, Selene L, Sam H, Marion D, Venkateswar J. Effect of Verapamil on the Pharmacokinetics of Aliskiren in Healthy Participants. *J Clin Pharmacol* 2011;51:218-228.
58. Wessler JD, Grip LT, Mendell J, Giugliano RP. The P-glycoprotein transport system and cardiovascular drugs. *J Am Coll Cardiol* 2013;61:2495-2502.
59. Erik S, Mikael H, Ulf B, Hans L. Effects of verapamil on the pharmacokinetics and hepatobiliary disposition of fexofenadine in pigs. *Eur J Pharmac Sci* 2014;57:214-223.
60. Wang RB, Kuo KL, Lien LL, Lien EJ. Structure-activity relationship: analyses of p-glycoprotein substrates and inhibitors. *J Clin Pharm Ther* 2003;28:203-228.
61. Molento MB, Lifschitz A, Sallovitz J, Lanusse C, Prichard R. Influence of verapamil on the pharmacokinetics of the antiparasitic drugs ivermectin and moxidectin in sheep. *Parasitol Res* 2004;92:121-127.
62. Tripta B, Gautam M, Manu J, Roop KK, Sushama T. Effect of P-glycoprotein inhibitor, verapamil, on oral bioavailability and pharmacokinetics of irinotecan in rats. *Eur J Pharmac Sci* 2009;36:580-590.
63. Stringer KA, Mallet J, Clarke M, Lindenfeld JA. The effect of three different oral doses of verapamil on the disposition of theophylline. *Eur J Clin Pharmacol* 1992;43:35-38.
64. Carrillo JA, Benitez J. Clinically significant pharmacokinetic interactions between dietary caffeine and medications. *Clin Pharmacokinet* 2000;39:127-153.
65. Stacey LB, Anthony T, Joyce AO, Andrea MD, Marianne N, Frederick PO, et al. Effect of R-Verapamil on the Pharmacokinetics of Paclitaxel in Women with Breast Cancer. *J Clin Oncol* 1995;13:2039-2042.
66. He Z, Si CZ, Shailendra S, Qian M, Chuan HD, Min HL, et al. Study on the pharmacokinetics profiles of Polyphyllin I and its bioavailability enhancement through co-administration with P-glycoprotein inhibitors by LC-MS/MS method. *J Pharmac Biomed Anal* 2015;107:119-124.
67. Kees N, Robert O, Jan D. Effects of verapamil on the pharmacokinetics of daunomycin in the rat. *Cancer Chemother Pharmacol* 1987;20:176-178.
68. Ruben P, Cristina P, Maria JN, Pedro U, Aurelio S, Luis M, et al. Influence of verapamil on pharmacokinetics and transplacental transfer of ivermectin in sheep. *Small Ruminant Res* 2010;93:103-109.
69. Jun SC, Xiuguo L. The effect of verapamil on the pharmacokinetics of paclitaxel in rats. *Eur J Pharmac Sci* 2005;24:95-100.
70. Kai H, Jinping H, Xue L, Yan L. Role of P-Glycoprotein in Intestinal Absorption of FB2, a Promising Abl/Src Dual Tyrosine Kinase Inhibitor. *Drug Metab. Pharmacokinet* 2012; 27: 486-494.
71. Naseem A, Sushama T, Abdul A, Rook KK, Manu J. Potential of a novel self nanoemulsifying carrier system to overcome P-glycoprotein mediated efflux of etoposide: In vitro and ex vivo investigations. *J Drug Deliv Sci Technol* 2015;28:18e27.
72. Cuiping Y, Tianhong Z, Zheng L, Liang X, Fei L, Jin XR, et al. P-glycoprotein is responsible for the poor intestinal absorption and low toxicity of oral aconitine: In vitro, in situ, in vivo and in silico studies. *Toxicol Appl Pharmacol* 2013;273:561-568.
73. Niclas P, Ebba B, Patrik F, Mikael H, Ulf B, Lars K, et al. First-Pass Effects of Verapamil on the Intestinal Absorption and Liver Disposition of Fexofenadine in the Porcine Model. *DMD* 2006;34:1182-1189.