



## Bone morphogenic protein receptor type 1a (BMPR1A) and Caveolin-1 associated with trastuzumab resistance of breast cancer cells

Zohreh Rezaei<sup>1</sup>, Kazem Dastjerdi<sup>2</sup>, Dor Mohammad Kordi-Tamandani<sup>3\*</sup>

<sup>1</sup> Department of Biology, University of Sistan and Baluchestan, Zahedan, Iran

<sup>2</sup> Department of Medical Biotechnology, Faculty of medicine, Birjand University of Medical Sciences, Birjand, Iran

<sup>3</sup> Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

### Please cite this article as:

Rezaei Z, Dastjerdi K, Kordi-Tamandani DM. Bone morphogenic protein receptor type 1a (BMPR1A) and Caveolin-1 associated with trastuzumab resistance of breast cancer cells. *Iranian J Pharmacol Ther.* 2019 (October);17:1-5.

### ABSTRACT

Trastuzumab is a specific monoclonal antibody used for therapeutic of the human epidermal growth factor receptor 2 (HER-2) -positive metastatic breast cancer. But, resistance to trastuzumab is a major obstacle in clinical efficiency. During the past years, several studies have been done to find the mechanisms contributing to trastuzumab resistance. Previous studies have highlighted that bone morphogenic protein (BMP) signaling can indicate a pathway in cancer for sensitizing cells to chemotherapy. Also, it was suggested that Caveolin-1 is essential for the formation of caveolae and endocytic membrane transport and has a critical role in drug resistance and metastasis in cancer. The purpose of this study was to assess the expression of BMP receptor type1A (BMPR1A) and Caveolin-1 genes in compare with trastuzumab-sensitive and resistance BT-474 cells. Trastuzumab-resistant BT-474 cells were established by continuous subjection to trastuzumab for six months. Then, an MTT assay was done for determining the resistance. After that, the Expression of BMPR1A and Caveolin-1 levels were assessed through real-time PCR. Caveolin-1 expression levels increased significantly (2.4 fold,  $p < 0.05$ ) whereas BMPR1A levels down-regulated significantly (8.26 fold,  $p < 0.05$ ) in BT-474-R compared to the parental cells. Our results proposed that BMPR1A and CAV1 regulation take part in BT-474 trastuzumab resistance breast cancer. Therefore, further experiments are required to confirm the role/s of BMPR1A and CAV1 in trastuzumab resistance breast cancer.

**Conflicts of Interest:** Declared None

**Funding:** None

### Keywords

Breast cancer,  
Drug resistance,  
Trastuzumab,  
CAV1,  
BMPR1A

### Corresponding to:

Dor Mohammad Kordi-Tamandani,  
Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Email:

[dor\\_kordi@yahoo.com](mailto:dor_kordi@yahoo.com)

Received: 29 Apr 2019

Published: 21 Oct 2019

### INTRODUCTION

Amplification or overexpression of HER2 occurs in about 20–30% of metastatic breast cancers. HER2 protein overexpression has come to be identified as important markers for metastatic HER2- overexpressing breast cancer and the target of specific therapies [1]. Trastuzumab is a recombinant humanized monoclonal anti-HER2 antibody (Herceptin) and was the first approved monoclonal antibody by the FDA used for the solid tumor therapy. Most patients with progressive HER-2 positive breast cancer responding initially to trastuzumab acquired resistance over the first year of

treatment [2]. several mechanisms have been studied to identify the cause of resistance to trastuzumab in breast cancer, including reduced antibody affinity or HER2 expression, using alternative receptor tyrosine kinases signaling pathways, and changed intracellular signaling including the loss of PTEN expression, increased Akt activity, reduced activity of p27kip1 ( a cell cycle regulator), leading to the over-proliferation of cells [3].

Caveolin-1 (Cav-1), a 21 kDa protein is in “cave-like” invaginations of the cell membrane known as caveolae. Cav-

1 is a main essential component of organelles with a size range of 50–100 nm and is a necessity for their formation. Cav-1/caveolae are associated with several cellular pathways such as endocytosis, signal transduction, and lipid homeostasis [4, 5].

Studies have been displayed that Cav-1 is a molecular hub that integrates the activity of a multitude of signaling molecules including epidermal growth factor receptor (EGFR), HER2, Src and the mitogen-activated protein kinase (MAPK) cascade. Most of these signals are involved in cancer development [6]. Also, recent studies show that Cav-1 can be a factor in mediating stress-associated and drug resistance. caveolin-1 was involved in up-regulation of adriamycin-resistant breast cancer cells [7], Taxol- and gemcitabine-resistant lung cancer cells [8] and multidrug resistant colon cancer cells [9].

BMP is a special extracellular multifunctional cytokine that is a member of the large transforming growth factor-beta (TGF- $\beta$ ) superfamily. For binding to BMP ligands (BMP2, 4, and 7) the BMP type 1 receptor (BMPR1A) that is a transmembrane receptor on the cell surface, forming a heterodimer with the type 2 receptor (BMPR2). Several investigators have revealed that these morphogens play critical roles in proliferation, development, differentiation, and apoptosis [10].

Further experiments showed BMP signaling act as a tumor suppressor. Howe et al. displayed the formation of Juvenile Polyposis Syndrome through the loss of BMPR1a [11]. Also, the loss of BMPR1a form hamartomas that usually are benign tumors. Recent works have offered that low expression of BMPR1A may be related to poor prognosis in tumors [12, 13]. However, the expression levels of cav1 and BMPR1A and their effect on drug resistance of HER2 positive breast cancer have not been thoroughly evaluated.

## MATERIALS AND METHODS

### Cell culture and establishment of trastuzumab-resistant cells

The human BT-474 breast cancer cells were taken from the Iranian Biological Research Center. The cells were cultured in media (DMEM-f12) with 10% fetal bovine serum (FBS) and incubated under humidified chamber containing 5% CO<sub>2</sub> at 37°C. As reported previously, to obtain Trastuzumab-resistant cells, The BT-474 parental cells were cultured continuously in the presence of low-dose trastuzumab (5  $\mu$ g/ml) for 6 months [14]. Then, parental and trastuzumab-resistant BT-474 breast cancer cells were cultured in the ab-

sence and presence of trastuzumab, respectively.

### RNA extraction and quantitative real-time PCR (qPCR)

Total RNA was extracted from each samples (sensitive and resistant cultured BT-474 cells) by using the Trizol Reagent (Invitrogen, Carlsbad, CA, USA). MRNAs were reverse transcribed to cDNAs using the miScript II RT Kit (Qiagen). qPCR was done through the StepOne™ Real-Time PCR System (Applied Biosystems Inc., Hercules, CA, USA). Then cDNAs were detected and amplified by Thermo Fisher Master Mix of SYBR Green PCR (Thermo Fisher, England). The sequence information used in this quantitative Real-Time PCR was listed in Table 1. GAPDH was as internal controls for mRNAs. Quantitation of gene expression evaluated by  $\Delta\Delta$ Ct calculation, where Ct is the threshold cycle. QRT-PCR was performed in triplicate.

### Cell survival assay

Cell survival was estimated by a colorimetric MTT assay [21]. 3-(4,5-Dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) was obtained from Atocel (Graz, Austria). Briefly, BT-474 cells were cultured for 24 h into the 96-well plate at a density of  $1 \cdot 10^4$  for each well. Then, cells were incubated with medium containing various concentrations of trastuzumab (0.21–2100  $\mu$ g/mL) for another 72 hours. After the incubation period, 100  $\mu$ L of 10mM MTT working solution was added into every well of the plate and incubated for 4 h at 37°C. After addition of DMSO, the plate was placed on an orbital shaker for 45 min at RT for dissolving the formazan crystals. The absorbance intensity of cells was evaluated at 570 nm with the Epoch Microplate Reader. The relative cell viability (%) was calculated through the following formula: survival percentage =  $(A_{\text{drug-treated cells}} - A_{\text{blank cells}}) / (A_{\text{untreated cells}} - A_{\text{blank cells}}) \cdot 100$ .

### Statistical analysis


Statistical analyses were done using SPSS version 16.0 for Windows. The significance of differences in the quantitative PCR results was evaluated by Mann-Whitney test. The P-values less than 0.05 were assumed as statistically significant.

## RESULTS

To define the status of HER2 signaling in sensitive and resistance BT-474 cells, these cells were treated with trastuzumab for 72 hours, and the survival of cells was as-

Table 1. Primer sequences for quantitative real-time polymerase chain reaction

Primer	Sequence
CAV1 Forward	GCGACCCTAAACACCTCAAC
CAV1 Reverse	ATGCCGTCAAAACTGTGTGTC
BMPR1A Forward	TAGTTCGCTGAACCAATAAAGG
BMPR1A Reverse	GTCAGAAAATGGAGTAACCTTA
GAPDH Forward	TGGACTCCACGACGTACTCAG
GAPDH Reverse	CGGGAAGCTTGTTCATCAATGGAA



**Figure 1.** Analysis of Sensitivity to trastuzumab by MTT cell survival assay. Cells were plated in 96-well plates and trastuzumab (0.21–2100 mg/mL) containing medium was added 24 h later. MTT assay was performed after 72 h incubation. \* $p < 0.05$  or \*\* $p < 0.01$  statistically significant when compared between two groups using t-test.

essed by MTT assay. As revealed in Figure 1, BT-474-R cells represent significantly higher resistant to trastuzumab in compare with the sensitive cells ( $P < 0.01$ , Fig. 1). At the maximum concentration of trastuzumab (2100 mg/mL), the survival cells were about 80% , but just about 44% of the sensitive cells be viable [15]. These results proposed that these resistant cells show in vitro proliferation advantages over non-resistant BT-474 cells.

**Upregulation of cav1 and downregulation of BMPR1A contributing to trastuzumab-chemotherapy resistance in bt-474 cells**

To assess the function of BMPR1a and CAV1 in trastuzumab resistance of breast cancers, the expression of these genes were surveyed in parental and trastuzumab-resistant BT-474 cells by Quantitative RT-PCR. The expression level of CAV1 was meaningfully upregulated ( $P < 0.05$ , fold change 2.4); but BMPR1a was downregulated significantly ( $P < 0.05$ , fold change -8.26) (Table 2 and Fig. 2).


**DISCUSSION**

In the present study, we showed that enhanced level of CAV1 or downregulation of BMPR1A could be independent predictors of drug resistance in HER2 positive breast cancer. Caveolin-1 was shown that interact with different intracellular signaling pathways. Agelaki et al displayed that Cav-1 contribute EGFR signaling and promote proliferation and migration [16]. Cav-1 inhibits EGFR pathway through ERK1/2 and Grb2-Sos-Ras, but activates PI3K pathway [17]. Interaction of Cav-1 with Akt causes EGFR activation which causes an increase in cell survival [18].

The expression of was mainly associated with EGFR positivity, for instance, CAV1 was expressed in 68% of HER2 positive breast cancers, however in EGFR-negative

**Table 2.** Relative expression levels of CAV1 and BMPR1A in BT474 sensitive and resistance cells

	Fold change
CAV1	2.4
BMPR1A	-8.26



**Figure 2.** The fold change in expression of BMPR1A and CAV1 in BT-474 and BT-474-R. The RNA was extracted using TRIzol reagent, converted into cDNA by RT-PCR and submitted to qRT-PCR using specific primers, as described in methods. Each bar indicates the mean  $\pm$  SD of triplicate assays.

breast carcinomas just exhibit 3.6% expression of CAV1 [19]. Primary experiments proposed that EGFR could be localized within caveolae and CAV1 expression would adjust the EGFR signaling pathway by receptor sequestration and controlling receptor trafficking [20].

Gang et al in 2016 showed that Cav-1 overexpression increases HER-2 signaling and enhances proliferation and migration in breast cancer cells [21]. Upregulation of Cav-1 associated with histological differentiation, intrahepatic metastasis, expression of VEGF and venous invasion [22]. In inflammatory breast cancer cells and also tissues expression of Cav-1 was increased [23].

Several studies showed that Cav-1 expression was meaningfully enhanced in some drug-resistant cancer cells. In A549-T12 (taxol-resistant lung cancer cell line) that displayed 9-fold resistance taxol compared with the parental cell line, the expression of Cav-1 upregulated 3.4 fold. Increasing resistance taxol to 17-fold, enhanced Cav-1 expression 9.5-fold [24]. However, upregulation of Cav-1 level has been described in another drug-resistant cancer cells including MCF-7 cells resistant to adriamycin, SKVLB1 cells resistant to vinblastine and HT-29 cells resistant to colchicine [25].

In 1965 BMPs primary presented as an osteogenic factor. BMPs are critical extracellular multifunctional signaling cytokine that demonstrates as a member of TGF- $\beta$  superfamily [26]. BMPRs are transmembrane serine/threonine kinase receptors. when BMPs interacted with type I and II transmembrane serine/threonine kinase receptor, their signals were transduced [27]. There are four different types I receptors and three distinct types II receptors including —Bmpr1a (Alk3), Bmpr1b (Alk6), Acvr11 (Alk1), Acvr1 (Alk2), Bmpr2 (BMPRII), Acvr2a (ActRIIA) and Acvr2b (ActRIIB) [28]. In this study, we showed that BMPRI1A level was lower in BT-474 resistant cells compared with parental BT-474 cells. Due to the metastatic potential of resistance cells, downregulation of this tumor suppressor can have a critical role in the induction of metastasis. Our results in this study showed that downregulation of BMPRI1a decreased tumor burden and metastatic potential. Several surveys have displayed that BMPRI1A has critical roles in the suppression of tumor progression [29] BMPRI1A has been demonstrated to be a tumor suppressor in tumorigenesis of skin .whereas findings of the lower lip squamous cell carcinoma showed that upregulation of BMPRI1A displayed a very strong connection with high malignancy score and advanced clinical staging [30].

## CONCLUSION

In conclusion, we evaluated the levels of Caveolin-1 and BMPRI1A in trastuzumab-resistant and sensitive BT-474 cells. These results show that CAV1 and BMPRI1A may be independent tumor markers reflecting the resistance in BT-474 HER2 positive breast cancer. More researches are essential for defining the function of CAV1 and BMPRI1A in the HER2 positive breast cancer.

## CONFLICTS OF INTEREST

The author(s) declare(s) that there is no conflict of interest regarding the publication of this article

## REFERENCES

- Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science* 1987;235(4785):77-82.
- Slamon DJ, Leyland-Jones B, Shak S, Fuchs H, Paton V, Bajamonde A, et al. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *New Engl J Med* 2001;344(7):83-92.
- Arteaga CL, Sliwkowski MX, Osborne CK, Perez EA, Puglisi F, Gianni L. Treatment of HER2-positive breast cancer: current status and future perspectives. *Nat Rev Clin Oncol* 2011;9(1):16-32.
- Williams TM, Cheung MWC, Park DS, Razani B, Cohen AW, Muller WJ, et al. Loss of caveolin-1 gene expression accelerates the development of dysplastic mammary lesions in tumorprone transgenic mice. *Mol Biol Cell* 2003;14: 1027-42.
- Williams TM, Medina F, Badano I, Hazan RB, Hutchinson J, Muller WJ, et al. Caveolin-1 gene disruption promotes mammary tumorigenesis and dramatically enhances lung metastasis in vivo. Role of Cav-1 in cell invasiveness and matrix metalloproteinase (MMP-2/9) secretion. *J Biol Chem* 2004;279:51630-46.
- Burgermeister ELM, Röcken C, Schmid RM, Ebert MP. Caveats of caveolin-1 in cancer progression. *Cancer Lett* 2008;268:187-201.
- Cai C, Chen J. Overexpression of caveolin-1 induces alteration of multidrug resistance in Hs578T breast adenocarcinoma cells. *Int J Cancer* 2004;111:522-9.
- Selga E, Morales C, Noé V, Peinado MA, Ciudad CJ. Role of caveolin 1, E-cadherin, Enolase 2 and PKC $\alpha$  on resistance to methotrexate in human HT29 colon cancer cells. *BMC Med Genom* 2008;1(35).
- Ho CC, Kuo SH, Houg PH, Houg HY, Yang CHH, Yang PCH. Caveolin-1 expression is significantly associated with drug resistance and poor prognosis in advanced non-small cell lung cancer patients treated with gemcitabine-based chemotherapy. *Lung Cancer* 2008;59:105-10.
- Xiao YT, Xiang LX, Shao JZ. Bone morphogenetic protein. *Biochem Biophys Res Commun* 2007;362(3):550-3.
- Howe JR, Bair JL, Sayed MG, Anderson ME, Mitros FA, Petersen GM, et al. Germline mutations of the gene encoding bone morphogenetic protein receptor 1A in juvenile polyposis. *Nat Genet* 2001;28(2):184-7.
- Kim IY, Ahn HJ, Tokunaqa H, Song W, Devereaux LM, Jin D, et al. Expression of bone morphogenetic protein receptors type-IA, -IB and -II correlates with tumor grade in human prostate cancer tissues. *Cancer Res* 2000;60(11):2840-4.
- Kwan KM, Li AG, Wang XJ, Wurst W, Behringer RR. Essential roles of BMPRIA signaling in differentiation and growth of hair follicles and in skin tumorigenesis. *Genesis* 2004;39(1):10-25.
- Gong CYY, Wang Y, Liu B, Wu W, Chen J, Su F, et al. Up-regulation of miR-21 mediates resistance to trastuzumab therapy for breast cancer. *J Biol Chem* 2011;286(21):19127-37.
- Rezaei Z, Sebzari A, Kordi-Tamandani DM, Dastjerdi K. Involvement of the Dysregulation of miR-23b-3p, miR-195-5p, miR-656-5p, and miR-340-5p in Trastuzumab Resistance of HER2-Positive Breast Cancer Cells and System Biology Approach to Predict Their Targets Involved in Resistance. *DNA and Cell Biol* 2019;38(2):184-92.
- Agelaki S, Spiliotaki M, Markomanolaki H, Kallergi G, Mavroudis D, Georgoulas V, et al. Caveolin-1 regulates EGFR signaling in MCF-7 breast cancer cells and enhances gefitinib-induced tumor cell inhibition. *Cancer Biol Ther* 2009;8(14):70-7.
- Park WY, Park JS, Cho KA, Kim DI, Ko YG, Seo JS, et al. Up-regulation of caveolin attenuates epidermal growth factor signaling in senescent cells. *J Biol Chem* 2000;275(208):47-52.
- Park JH, Lee MY, Han HJ. A potential role for caveolin-1 in estradiol-17 $\beta$ -induced proliferation of mouse embryonic stem cells: involvement of Src, PI3K/Akt, and MAPKs pathways. *Int J Biochem*

- Cell Biol 2009;41(6):59-65.
19. Pike LJ. Growth factor receptors, lipid rafts, and caveolae: an evolving story. *Biochim Biophys Acta* 2005;1746(2):60-73.
  20. Khan EM, Jill M, Heidinger LM, Lisanti MP, Ravid T, Goldkorn T. Epidermal growth factor receptor exposed to oxidative stress undergoes Src- and caveolin-1-dependent perinuclear trafficking. *J Biol Chem* 2006;281(144):86-93.
  21. Gang Zhang TZ, Cui Y, Mishra R, Luan T, Xie P, Zhao R. Caveolin-1 regulates proliferation and metastasis of human breast cancer cells by activating Her-2. *Int J Clin Exp Med* 2016;9(5):7700-9.
  22. Zhang ZB, Cai L, Zheng SG, Xiong Y, Dong JH. Overexpression of caveolin-1 in hepatocellular carcinoma with metastasis and worse prognosis: correlation with vascular endothelial growth factor, microvessel density and unpaired artery. *Pathol Oncol Res* 2009;15:495-502.
  23. Van den Eynden GG VLS, Van der Auwera I, Merajver SD, Van Marck EA, Van Dam P, Vermeulen PB, et al. Overexpression of caveolin-1 and -2 in cell lines and in human samples of inflammatory breast cancer. *Breast Cancer Res Treat* 2006;95(3):21-8.
  24. Yang CP, Galbiati F, Volonte D, Horwitz SB, Lisanti MP. Upregulation of caveolin-1 and caveolae organelles in Taxol-resistant A549 cells. *FEBS Lett* 1998;439(3):68-72.
  25. Lavie Y, Fiucci G, Liscovitch M. Up-regulation of caveolae and caveolar constituents in multidrug-resistant cancer cells. *J Biol Chem* 1998;273(49):32-8.
  26. Urist MR. Bone morphogenetic protein: the molecularization of skeletal system development. *J Bone Miner Res* 1997;12:343-6.
  27. Kawabata M, Imamura T, Miyazono K. Signal transduction by bone morphogenetic proteins. *Cytokine Growth Factor Rev* 1998;9(1):49-61.
  28. Feng XH, Derynck R. Specificity and versatility in tgf-beta signaling through Smads. *Cell Dev Biol* 2005;21:659-93.
  29. Bleuming SA, He XC, Kodach LL, Hardwick JC, Koopman FA, Ten Kate FJ, et al. Bone morphogenetic protein signaling suppresses tumorigenesis at gastric epithelial transition zones in mice. *Cancer Res* 2007;67(17):8149-55.
  30. de Carvalho CHP, Nonaka CFW, de Araujo CRF, de Souza LB, Pinto LP. Immunoeexpression of bone morphogenetic protein-2 (BMP-2), BMP receptor type IA, and BMP receptor type II in metastatic and nonmetastatic lower lip squamous cell carcinoma. *J Oral Pathol Med* 2011;40(2):181-6.