

RESEARCH ARTICLE

Effect of Honey on CYP3A4 Enzyme and P-Glycoprotein Activity in Healthy Human Volunteers

VINOD THOMAS, KESAVAN RAMASAMY, RAJAN SUNDARAM and ADITHAN

CHANDRASEKARAN

For author affiliations, see end of text.

Received May 25, 2006; Revised August 6, 2007; Accepted August 14, 2007

This paper is available online at <http://ijpt.iums.ac.ir>

ABSTRACT

The activity of cytochrome p450 isozyme 3A4 (CYP3A4) enzyme and P-glycoprotein (P-gp) is modulated by grapefruit juice and herbal drugs. CYP3A4 is the major phase I drug metabolizing enzyme and P-gp is an ATP-dependent drug efflux pump that regulates the intestinal absorption of orally administered drugs. Honey is commonly consumed as a dietary supplement. However, its influence on human CYP3A4 and P-gp activity is not yet well documented. Therefore, we investigated the influence of a 10-day honey administration on CYP3A4 and P-gp activity in healthy volunteers using carbamazepine and digoxin as their probe drugs respectively. A within-group pharmacokinetic study was done in 12 healthy volunteers. They were administered single oral dose of carbamazepine (200 mg) and digoxin (0.5 mg) before and after 10 days of honey (10 ml twice daily) intake. Blood samples (5ml) were collected at 0, 0.25, 0.5, 0.75, 1.0, 1.5, 2, 4, 8, 12, 24, 48 and 72 h after drug administration. Concentration of carbamazepine and digoxin in plasma was measured by HPLC and RIA method respectively. Ten days of honey administration did not significantly alter the C_{max} , T_{max} and $AUC_{(0-t)}$ of carbamazepine (probe drug for CYP3A4) and digoxin (probe drug for P-gp). Our results suggest that honey may not significantly modulate the CYP3A4 enzyme and P-glycoprotein activity. The coadministration of honey with drugs may not result in significant drug interactions.

Keywords: Honey, CYP3A4, P-glycoprotein, carbamazepine, digoxin

Honey is a natural saccharine product made by honeybees from the nectar of flowers [1]. Being a natural source of fructose and glucose with some oligosaccharides, proteins, vitamins and minerals, honey has become a dietary supplement for healthy individuals [2]. Honey is also consumed by many patients with diabetes, hypertension and epilepsy who receive drugs for their ailments. This increases the possibility of honey-drug interaction. Most of the herb-drug interactions occur at the level of metabolism and drug transport mediated by CYP 450 group of drug metabolizing enzymes and P-glycoprotein (P-gp) respectively [3].

Among the CYP group of drug metabolizing enzymes, CYP3A4 is the major phase I drug metabolizing enzyme. It is present in the liver, jejunum, colon and pancreas. It has broad substrate specificity and is responsible for metabolism of more than 50% of administered drugs [4]. There are few studies showing the effect of honey on CYP3A4. Animal studies have shown that multiple doses of honey induced CYP3A4 activity [5,6]. In a study done in humans, single oral dose of honey failed to show any significant effect on CYP3A4 [7]. The effect of multiple doses of honey on CYP3A4 in humans has not been reported to date. It has been well documented that the CYP3A4 enzyme is involved in the metabolism and elimination of carbamazepine [8]. The pharmacokinetics of carbamazepine is influenced by alterations in the catalytic activity of CYP3A4 [9]. Hence, carbamazepine is used as a probe drug for assessing the CYP3A4 enzyme activity in our study.

P-glycoprotein (P-gp) is an ATP dependent drug efflux pump. It plays an important role as a secretory system in the intestinal barrier and regulates the intestinal absorption of orally administered drugs [10]. Many clinically important drugs viz., digoxin, losartan, erythromycin and rifampin are substrates for P-gp. Some of them besides being a substrate also induce or inhibit the P-gp activity. Drugs like fexofenadine, digoxin and loperamide are used as probe drugs to assess P-gp activity [11]. Among them, digoxin is most commonly used [12]. The effect of various dietary derivatives and herbal

Table 1. Pharmacokinetic parameters of carbamazepine (200 mg single oral dose) before and after 10 days of honey administration

Pharmacokinetic parameters	Before honey	After honey
C_{\max} ($\mu\text{g.ml}^{-1}$)	4.1 ± 0.28	4.2 ± 0.31
T_{\max} (h)	10.1 ± 1.60	9.0 ± 0.90
$AUC_{(0-72)}$ ($\mu\text{g.h.ml}^{-1}$)	203.1 ± 15.30	208.2 ± 17.20

Values are shown as mean \pm SEM. (n=12)

products on the P-gp activity has also been studied. In an *in vitro* study using various fruit extracts, it was found that extracts of strawberry, orange, apricot and mint inhibited the intestinal P-gp [13]. In another *in vitro* study using rat small intestine, extracts of grapefruit juice and orange juice inhibited the transport activity of P-gp [14]. In a study done in humans, grapefruit juice had no effect on P-gp activity [15]. Another human study revealed that St. John's Wort, an herbal product induced P-gp activity [16]. This shows that P-gp is a potential target for drug interactions exhibited by herbal compounds. The effect of honey on P-gp activity has not been studied so far.

Since we wanted to know whether honey, a natural dietary supplement, will interact with concomitantly administered drugs, we investigated the effect of multiple dose administration of honey on CYP3A4 and P-gp activity in humans using carbamazepine and digoxin as the probe drugs respectively. Carbamazepine is a CYP3A4 substrate but it is not a substrate for P-gp [17]. On the other hand, digoxin is a substrate for P-gp only and not a substrate for CYP3A4 [18]. Hence any change in the pharmacokinetic profile of carbamazepine and digoxin due to honey administration may reflect the change in the activity of CYP3A4 and P-gp respectively.

MATERIALS AND METHODS

A within group pharmacokinetic study was done in 12 healthy male volunteers (Age 20-45 years). The mean age of the volunteers was 27.4 ± 1.96 yrs (mean \pm SEM) and their mean body mass index was 23.2 ± 0.94 Kg/m^2 (mean \pm SEM). The study was approved by institutional ethics committee. A written informed consent was taken from all the volunteers. The health of the volunteers was assessed by doing a thorough physical examination and by performing ECG, liver and kidney function tests. Volunteers suffering from chronic diseases or taking concomitant medications were excluded from the study. Similarly, regular users of alcohol and/or tobacco, those with history of vomiting after

goxin intake, seizures and drug allergy were also excluded.

Study design

On day 1, single oral dose of 200 mg carbamazepine (Tegrital, Novartis [India] Limited) and 0.5 mg digoxin (Lanoxin, Burrough's Wellcome, [India] Limited) were administered to the volunteers at 7 AM who were fasted overnight. They were not allowed to take food for further 2 h. Blood samples were collected from indwelling venous catheter using heparinised disposable syringes just before and at 0.25, 0.5, 0.75, 1.0, 1.5, 2, 4, 8, 12, 24, 48, 72 h after administration of drugs. A standardized breakfast and lunch were given to all the volunteers. From day 5 to day 14, the volunteers were administered 10 ml of honey (Periyakulam Sarwodaya Sangh, Khadi Vastralaya, Theni District, Tamilnadu, South India; Lot No.4/2002) twice daily in empty stomach with 200 ml of water. On day 15, the volunteers were given single oral dose of 200 mg carbamazepine and 0.5 mg digoxin. The blood samples were collected as mentioned before. After separation of the plasma, the samples were stored at -20°C till the drug assays were done. The study protocol is shown as a flow chart in Figure 1.

The honey used in the present study was tested for its purity in Public Health Laboratory, Pondicherry, India. It was found to be within PFA (Prevention of food adulteration act-1955, India) values. It was composed of reducing sugar 71.6%, moisture 24%, sucrose 2.4% and ash 0.3%. The fructose/glucose ratio was 0.97%.

Drug assays

Serum carbamazepine concentration was estimated using a HPLC method [19]. The plasma sample (900 μl) and internal standard (900 μl) were taken in a 2 ml micro centrifuge tube. After vortex mixing, 600 μl was transferred to a conical flask, into which 4:1 mixture of chloroform: methanol was added. After mixing in an orbital shaker, the contents of conical flask were transferred to centrifuging tubes. After centrifugation at 2500 rpm for 10 min, the upper protein layer was transferred into evaporating tubes for evaporation at 50°C . The dried evaporated samples were reconstituted in 400 μl

Table 2. Pharmacokinetic parameters of digoxin (0.5 mg single oral dose) before and after 10 days of honey administration

Pharmacokinetic parameters	Before honey	After honey
C_{\max} ($\mu\text{g.ml}^{-1}$)	2.6 ± 0.22	2.5 ± 0.18
T_{\max} (h)	1.5 ± 0.26	1.2 ± 0.14
$AUC_{(0-4)}$ (ng.h.ml^{-1})	6.1 ± 0.44	6.2 ± 0.24
$AUC_{(0-72)}$ ($\mu\text{g.h.ml}^{-1}$)	28.9 ± 8.80	27.6 ± 2.20

Values are shown as mean \pm SEM. (n=12)




Fig 1. The study plan described as a flow chart.

of mobile phase composed of acetonitrile: methanol: (T_{max}) were read directly from the actual plasma concentration data. The area under the plasma concentration HPLC. The inter-day coefficient of variation for carbamazepine HPLC assay was less than 7%.

The digoxin concentration in plasma was measured

according to the manufacturer's directions, in duplicate

using RIA kits (Orion diagnostics, Finland; Lot No.

1588501). Into the appropriate labeled test tubes, 25 µl

of calibrators, plasma samples (unknown concentration

of digoxin) and 100 µl of antiserum solution were

added. All the tubes were mixed on a vortex mixer and

then incubated for 1 h at room temperature. One ml of

separation reagent was added to all the test tubes and

mixed on a vortex mixer. They were centrifuged for 15-

20 min at 2000 g. After centrifugation, the supernatant

part was decanted and the head of each tube was tapped

firmly against absorbent paper. Radioactivity in each

tube was counted using gamma counter for 1 min. The

measurement range of the kit was 0.5-8.0 nmol/l. The

detection limit of the kit was 0.1 nmol/l.

Calculation of pharmacokinetic parameters:

The pharmacokinetic analysis was done using model independent formulae. The peak plasma concentration (C_{max}) and the time to reach peak plasma concentration

Statistical analysis

Pharmacokinetic data was expressed as mean ± SEM. The normality of the data was assessed by the Kolmogorov-Smirnov test. The C_{max}, T_{max} and AUC₍₀₋₇₂₎ were analysed by paired Student's 't' test. All the statistical analyses were carried out by using GraphPad Instat (version 3.05, 2000, San Diego, USA) software system. *p* < 0.05 was considered statistically significant.

RESULTS

Effect of honey on carbamazepine pharmacokinetics

The plasma carbamazepine concentration measured up to 72 h was not significantly altered by honey administration (Figure 2). After ten days of honey administration, there was no statistically significant change in the mean values of C_{max}, T_{max} or AUC₍₀₋₇₂₎ (Table 1).




Fig 2. Concentration versus time profile of plasma carbamazepine (AUC_{0-72}) before and after honey. Values are shown as mean \pm SEM.

Effect of honey on digoxin pharmacokinetics

The plasma digoxin concentrations measured up to 72 h were not significantly altered by honey administration (Figure 3). There was no statistically significant change in the mean values of C_{max} , T_{max} , $AUC_{(0-4)}$ or $AUC_{(0-72)}$ (Table 2).

DISCUSSION

Herbal extracts of garlic [20], grapefruit juice [21], St. John's Wort [22] and milk thistle [23] modulate the activity of CYP3A4 resulting in drug interactions. The

extracts of certain herbs used in traditional Chinese medicine like Angelica dahurica [24], Angelica sinensis [25] and Glycyrrhiza glabra [26] modulate the CYP3A4 activity. Herbal extracts of Curcumin [27], hawthorn [28], ginseng [29], green tea [30], milk thistle [31], piperine [32], and grapefruit juice [14], orange juice [14] and St. John's Wort [22] modulate P-gp activity.

Flavonoids present in herbs have been found to interact with CYP3A4 and P-gp [3]. Honey is a natural saccharine product rich in sugars and phytochemicals. The flavonoids present in honey are pinocembrine, pinobanskin, chrysin, galangin, quercetin, luteolin and




Fig 3. Concentration versus time profile of plasma digoxin (AUC_{0-72}) before and after honey. Values are shown as mean \pm SEM.

- kaempferol [2]. Studies in rabbits have shown that honey induced the metabolism of diltiazem [5] and carbamazepine [6]. In a human study, where the effect of single dose of honey on CYP3A4 was investigated using carbamazepine as a probe drug, honey failed to show statistically significant effect on carbamazepine pharmacokinetic parameters like C_{max} , T_{max} and $AUC_{(0-72)}$ [7]. Hence, we studied the effect of multiple doses of honey on carbamazepine pharmacokinetics. In our study, multiple doses of honey failed to significantly alter the pharmacokinetics of carbamazepine. Hence we assume that flavonoids present in honey may not have any significant effect on human CYP3A4 activity.
- Since honey did not change the pharmacokinetics of digoxin, it is assumed that the flavonoids present in honey may not have any significant effect on P-gp also. Becquemont *et al* investigated the effect of grapefruit juice on P-gp activity in 12 healthy volunteers using digoxin as a probe drug. It was found that grapefruit juice did not significantly inhibit the intestinal P-gp activity [15]. Although the C_{max} , T_{max} and $AUC_{(0-48)}$ of digoxin did not change significantly, there was a statistically significant increase in $AUC_{(0-4)}$ of digoxin (i.e. in first 4 h) following co-administration with grapefruit juice. This correlates with observations made by Westphal *et al* that P-gp inhibitors alter the early digoxin pharmacokinetics by interfering with the absorption of digoxin [33]. In our study, 10 days of honey administration did not alter even the early absorption pharmacokinetics (AUC_{0-4}) of digoxin.
- Honey and its various derivatives are natural dietary supplements consumed commonly all over the world. Healthy individuals prefer honey to maintain their health and patients with chronic illness take honey along with other medications. Hence the possibility of honey drug interactions cannot be ruled out. Apart from consuming honey as a single dose along with drugs, some patients take honey daily as a nutritional and healthy dietary supplement.
- Since, *in vitro* and *in vivo* studies have reported that herbal extracts may modulate CYP3A4 and P-gp activity resulting in various types of herb drug interactions; the safety of coadministration of honey with drugs needs to be studied. This study is an attempt to investigate the same. To the best of our knowledge, this is the first study in humans where the effect of multi dose honey administration on CYP3A4 and P-gp activity has been investigated. Based upon the present study, it can be concluded that honey does not affect the CYP3A4 mediated metabolism and P-gp mediated transport of concomitantly orally administered drugs. The coadministration of multiple doses of honey with drugs may not produce significant drug interactions.
- ## REFERENCES
- White JW Jr. Honey. *Adv Food Res* 1978; 24: 287-374.
 - Honey Stix [homepage on the internet]: Nutrition Info.: Therapeutic studies of honey: Honey health and therapeutic qualities (article provided by National Honey Board). Available from: <http://www.honeystix.com/honeystix/compendium.pdf>. Accessed on December 18, 2006.
 - Evans AM. Influence of dietary components on the gastrointestinal metabolism and transport of drugs. *Ther Drug Monit* 2000; 22:131-6.
 - Zhang Y, Benet LZ. The gut as a barrier to drug absorption: Combined role of cytochrome P450 3A and P-glycoprotein. *Clin Pharmacokinet* 2001; 40:159-68.
 - Koumaravelou K, Adithan C, Shashindran CH, Asad M, Abraham BK. Influence of honey on orally and intravenously administered diltiazem kinetics in rabbits. *Indian J Exp Biol* 2002; 40:1164-8.
 - Koumaravelou K, Adithan C, Shashindran CH, Asad M, Abraham BK. Effect of honey on carbamazepine kinetics in rabbits. *Indian J Exp Biol* 2002; 40: 560-3.
 - Malhotra S, Garg SK, Dixit RK. Effect of concomitantly administered honey on the pharmacokinetics of carbamazepine in healthy volunteers. *Methods Find Exp Clin Pharmacol* 2003; 25:537-40.
 - Kerr BM, Thummel KE, Wurden CJ, Klein SM, Kroetz DL, Gonzalez FJ *et al*. Human liver carbamazepine metabolism. Role of CYP3A4 and CYP2C8 in 10, 11-epoxide formation. *Biochem Pharmacol* 1994; 47:1969-79.
 - Spina E, Arena D, Scordo MG, Fazio A, Pisani F, Perucca E. Elevation of plasma carbamazepine concentrations by ketoconazole in patients with epilepsy. *Ther Drug Monit* 1997; 19:535-8.
 - Pan G, Wang G, Fawcett P. The role of P-glycoprotein in the intestinal barrier. *Asian J Drug Metab Pharmacokinet* 2002; 2:69-76.
 - Marzolini C, Paus E, Buclin T, Kim RB. Polymorphisms in human MDR1(P-glycoprotein): Recent advances and clinical relevance. *Clin Pharmacol Ther* 2004; 75 :13-33.
 - de Lannoy AI, Silverman M. The MDR1 gene product, P-glycoprotein, mediates the transport of the cardiac glycoside, digoxin. *Biochem Biophys Res Commun* 1992; 189: 551-7.
 - Deferme S, Van GJ, Augustijns P. Inhibitory effect of fruit extracts on P-glycoprotein related efflux carriers: An *in vitro* screening. *J Pharm Pharmacol* 2002; 54:1213-9.
 - Tian R, Koyabu N, Takanaga H, Matsuo H, Ohtani H, Sawada Y. Effects of grapefruit juice and orange juice on the intestinal efflux of P-glycoprotein substrates. *Pharm Res* 2002; 19: 802-9.
 - Becquemont L, Verstuyft C, Kerb R, Brinkmann U, Lebot, M, Jaillon P *et al*. Effect of grapefruit juice on digoxin pharmacokinetics in humans. *Clin Pharmacol Ther* 2001; 70: 311-6.
 - John A, Brockmoller J, Bauer S, Maurer A, Langheinrich M, Roots I. Pharmacokinetic interaction of digoxin with an herbal extract from St John's wort (*Hypericum perforatum*). *Clin Pharmacol Ther* 1999; 66: 338-45.
 - Owen A, Pirmohamed M, Tetley JN, Morgan P, Chadwick D, Park BK. Carbamazepine is not a substrate for P-glycoprotein. *Br J Clin Pharmacol* 2001; 51:345-9.
 - Greiner B, Eichelbaum M, Fritz P, Kreichgauer HP, von Richter O, Zundler J *et al*. The role of intestinal P-glycoprotein in the interaction of digoxin and rifampin. *J Clin Invest* 1999; 104:147-53.
 - Gerson B, Bell F, Chan S. Antiepileptic agents - primidone, phenobarbital, phenytoin and carbamazepine by reversed-phase liquid chromatography. *Clin Chem* 1984; 30:105-8.
 - Piscitelli SC, Burstein AH, Welden N, Gallicano KD, Falloon J. The effect of garlic supplements on the pharmacokinetics of saquinavir. *Clin Infect Dis* 2002; 34: 234-8.
 - Lown KS, Bailey DG, Fontana RJ, Janardan SK, Adair CH, Fortlage LA *et al*. Grapefruit juice increases felodipine oral availability in humans by decreasing intestinal CYP3A protein expression. *J Clin Invest* 1997; 99:2545-53.
 - Durr D, Stieger B, Kullak-Ublick GA, Rentsch KM, Steinert HC, Meier PJ *et al*. St John's Wort induces intestinal P-glycoprotein / MDR1 and intestinal and hepatic CYP3A4. *Clin Pharmacol Ther* 2000; 68:598-604.
 - Venkataramanan R, Ramachandran V, Komoroski BJ, Zhang S, Schiff PL, Strom SC. Milk thistle, a herbal supplement, decreases the activity of CYP3A4 and uridine diphosphoglu-

- 346 curonosyl transferase in human hepatocyte cultures. Drug Metab 37531.
347 Dispos 2000; 28:1270-3. 376
34824. Ishihara K, Kushida H, Yuzurihara M, Wakui Y, Yanagisawa T, 377
349 Kamei H *et al.* Interaction of drugs and Chinese herbs: Pharma-37832.
350 cokinetic changes of tolbutamide and diazepam caused by ex-379
351 tract of *Angelica dahurica*, J Pharm Pharmacol 2000; 52:1023-9. 380
35225. Guo LQ, Taniguchi M, Chen QY, Baba K, Yamazoe Y. Inhibi-38133.
353 tory potential of herbal medicines on human cytochrome P450-382
354 mediated oxidation: Properties of umbelliferous or citrus crude383
355 drugs and their relative prescriptions. Jpn J Pharmacol 2001; 85:384
356 399-408.
35726. Budzinski JW, Foster BC, Vandenhoeck S, Arnason JT. An *in*
358 *vitro* evaluation of human cytochrome P450 3A4 inhibition by
359 selected commercial herbal extracts and tinctures. Phytomedi-385
360 cine 2000; 7: 273-82.
36127. Chearwae W, Anuchapreeda S, Nandigama K, Ambudkar SV,386
362 Limtrakul P. Biochemical mechanism of modulation of human387
363 P-glycoprotein (ABCB1) by curcumin I, II, and III purified from388
364 Turmeric powder, Biochem Pharmacol 2004; 68: 2043-52. 389
36528. Tankanow R, Tamer HR, Streetman DS, Smith SG, Welton J L,390
366 Annesley T *et al.* Interaction study between digoxin and a prepa-391
367 ration of hawthorn (*Crataegus oxyacantha*), J Clin Pharmacol392
368 2003; 43: 637-42.
36929. Kim SW, Kwon HY, Chi DW, Shim JH, Park JD, Lee YH *et al.*393
370 Reversal of P-glycoprotein-mediated multidrug resistance by394
371 ginsenoside Rg(3). Biochem Pharmacol 2003; 65: 75-82. 395
37230. Sadzuka Y, Sugiyama T, Sonobe T. Efficacies of tea compo-396
373 nents on doxorubicin induced antitumor activity and reversal of397
374 multidrug resistance. Toxicol Lett 2000; 114:155-62. 398
- 399
- Zhang S, Morris M E. Effects of the flavonoids biochanin A, morin, phloretin, and silymarin on P-glycoprotein-mediated transport. J Pharmacol Exp Ther 2003; 304:1258-67.
- Velpandian T, Jasuja R, Bhardwaj RK, Jaiswal J, Gupta SK. Piperine in food: interference in the pharmacokinetics of phenytoin. Eur J Drug Metab Pharmacokinet 2001; 26: 241-7.
- Westphal K, Weinbrenner A, Giessmann T, Stuhr M, Franke G, Zschiesche M *et al.* Oral bioavailability of digoxin is enhanced by talinolol: Evidence for involvement of intestinal P-glycoprotein. Clin Pharmacol Ther 2000; 68: 6-12.

CURRENT AUTHOR ADDRESSES

Vinod Thomas, Department of Pharmacology, Jawaharlal Institute of Post-graduate Medical Education and Research Pondicherry-605 006, India. E-mail: drvinodthomas7@rediffmail.com (Corresponding author)

Kesavan Ramasamy, Department of Pharmacology, Jawaharlal Institute of Post-graduate Medical Education and Research Pondicherry-605 006, India.

Rajan Sundaram, Department of Pharmacology, Jawaharlal Institute of Post-graduate Medical Education and Research Pondicherry-605 006, India.

Adithan Chandrasekaran, Department of Pharmacology, Jawaharlal Institute of Post-graduate Medical Education and Research Pondicherry-605 006, India.