

1 RESEARCH ARTICLE

 2 Blood Biochemical Studies of Enrofloxacin in Yak
 3 after Intravenous Administration Article Title

4 SANJIB KHARGHARIA, CHANDANA CHOUDHURY BARUA, NABIN NATH and M. BHATTACHARYA

5 For author affiliations, see end of text.

6 Received August 7, 2006; Accepted July 10, 2007

7 This paper is available online at <http://ijpt.iums.ac.ir>

8 ABSTRACT

9 The blood biochemical parameters were investigated in yak (*Bos grunniens* L) after intravenous admini-
 10 stration of 5 mg/kg enrofloxacin. Blood samples were collected from the jugular vein daily for 5 days after
 11 the initiation of drug administration. Heparinized blood samples were collected for determination of
 12 mean corpuscular volume (MCV), erythrocyte sedimentation rate (ESR), hemoglobin (Hb) estimation and
 13 serum was collected for estimation of total protein, albumin, globulin, urea and blood glucose. Results of
 14 hematological and blood chemistry profile revealed that enrofloxacin did not change above mentioned
 15 parameters significantly. Therefore, the drug might be used safely in the yaks.

16 **Keywords:** *Enrofloxacin, Yak, Blood chemistry, Hematology*

17 Enrofloxacin, a fluoroquinolone antimicrobial is ap- 44 clinically to evaluate health status and to rule out the
 18 proved exclusively for veterinary use and has a broad 45 possibility of any diseases. They were housed in the
 19 spectrum of antibacterial activity with MIC values rang- 46 animal shed with concrete floor and were maintained on
 20 ing from 0.008 to 0.06 µg/ml. It has wide spread distri- 47 green fodder, dry grass and concentrate. Water was pro-
 21 bution to most tissues and body fluids with a potential 48 vided ad libitum.

23 Hence, it is effectively used in the treatment of septic- 49 *Experimental design*

24 mia, respiratory tract, urinary tract, skin, soft tissues, 50 The study was conducted in six clinically healthy
 25 bone and joint infections. Yak (*Bos grunniens* L), being 51 male yaks. Blood samples (12 ml) were collected by
 26 the most ecologically sustainable animal resource of 52 jugular vein puncture just before drug administration
 27 Indian Himalayas, is the mainstay for highlanders, pro- 53 (normal/control), 30 minutes after drug administration
 28 viding basic needs in terms of meat, milk, hair wool 54 and on 24 hours intervals for 5 days. 4 ml aliquots of
 29 and transportation in hilly terrain. For safety of admini- 55 collected blood were transferred to vials containing
 30 stration in various animals, biochemical properties of 56 heparin as anticoagulant in order to estimate haemoglo-
 31 enrofloxacin have been carried out in cattle [1], buffalo 57 bin, PCV and ESR. Remaining 8ml of blood samples
 32 [2] and goat [3]. However, available data are inade- 58 were kept in clean dry wide mouthed tubes in slanting
 33 quate to warrant effective clinical use of enrofloxacin in 59 position at room temperature to separate serum for esti-
 34 yak. In the light of the above reports, it was decided to 60 mation of total serum protein, albumin, globulin, urea,
 35 study possible biochemical and hematological changes 61 and blood glucose.

62 *Estimation of biochemical and hematological prop-*
63 *erties:*

37 MATERIALS AND METHODS

38 *Experimental animals*

39 The study was conducted in six male yaks (*Bos* 64 Standard procedures were followed for quantitative
 40 *grunniens* L) reared in the National Research Centre on 65 determination of biochemical and hematological proper-
 41 Yak, Nykmadung Farm, Dirang, Arunachal Pradesh, 66 ties. Haemoglobin was determined by acid hematin
 42 India. The animals were weighing between 270-330 kg 67 method described by Barker et al in 1965 [4], PCV by
 43 and 2 ½- 3 ½ years of age. The animals were examined 68 micro-haematocrit method as described by Prasad by
 69 1992 [5], and ESR by standard Wintrobe method. Bio-
 70 chemical analyses of samples were performed to meas-
 71 ure glucose, total protein, albumin, blood urea nitrogen

Table 1. Mean (n = 5) hematological and biochemical parameters after i.v. administration of enrofloxacin (5mg/kg) in Yak

Day	Hb (g%)	ESR (mm/24 h)	PCV (%)	Protein (g %)	Albumin (g/dl)	Globulin (g/dl)	Glucose (mg/dl)	Urea (mg/dl)
Normal	12.17±0.55 ^a	6.97±0.35 ^a	46.50±0.64 ^a	5.03±0.06 ^a	2.69±0.14 ^a	2.14±0.09 ^a	42.82±1.01 ^a	30.24±2.02 ^a
1 st	12.05±0.45 ^a	6.67±0.57 ^a	46.37±1.79 ^a	5.79±0.39 ^a	3.20±0.11 ^a	2.59±0.10 ^a	124.60±3.01 ^b	49.01±2.46 ^b
2 nd	12.57±0.42 ^a	7.08±0.56 ^a	43.90±1.49 ^a	5.63±0.34 ^a	3.15±0.11 ^a	2.44±0.09 ^a	78.91±1.71 ^c	89.04±3.11 ^c
3 rd	12.03±0.47 ^a	6.75±0.50 ^a	45.00±1.42 ^a	5.17±0.34 ^a	2.91±0.14 ^a	2.27±0.16 ^a	57.97±2.43 ^b	60.36±2.99 ^b
4 th	12.66±0.46 ^a	6.25±0.40 ^a	46.66±2.13 ^a	5.27±0.42 ^a	2.87±0.13 ^a	2.39±0.14 ^a	51.32±2.35 ^a	47.39±4.68 ^a
5 th	13.46±0.40 ^a	7.42±0.40 ^a	40.71±1.11 ^a	5.63±0.38 ^a	3.01±0.13 ^a	2.62±0.12 ^a	45.12±2.02 ^a	38.00±3.96 ^a

Having same superscripts no significant difference ($p < 0.05$)

Having different superscripts significant difference ($p < 0.05$)

72 and globulin as described in literature supplied with the 113tion of enrofloxacin. The mean level of protein, albu-
73 kits. For estimation of blood glucose, albumin, total 114min, and globulin in present study did not show signifi-
74 serum protein and blood urea nitrogen, a UV spectro- 115cant difference and was consistent with data reported by
75 photometer (Systronic) was used at wavelength 630, 116Nath et al., 2000 [9].

76 555 and 480 nm respectively in accordance with pro- 117 The present study suggests that administration of en-
77 vided instructions. Results were calculated according to 118rofloxacin intravenously at a rate of 5 mg/kg body
78 the formula provided in kit. 119weight in yak does not change the hematology and bio-
79 120chemical profile of the animal; hence it is suggested for
121 clinical use.

80 Statistical analysis

81 Analysis of variance was used to detect differences 122
82 between means for hematological and biochemical pa- 123
83 rameters. p values < 0.05 were considered to be statisti- 124
84 cally significant. 125

85 RESULTS

86 The results obtained in the study are presented in the 130
87 Table 1. 131

88 DISCUSSION

89 Mean (n=5) hematological and biochemical results 132
90 after intravenous administration of enrofloxacin 133
91 (5mg/kg) in yak are presented in Table 1. In the present 134
92 study, it is observed that blood glucose level increases 135
93 after intravenous injection of enrofloxacin in yak. This 136
94 might be due to stress which is developed due to re- 137
95 peated puncture of the jugular vein which in turn trig- 138
96 gers the release of epinephrine from adrenal medulla, 139
97 stimulating glycogen breakdown and also inhibits gly- 140
98 cogen synthesis in the liver, thus directing all available 141
99 glucose residue and precursors into the production of 142
100 free blood glucose [6]. It was observed that blood urea 143
101 level increased significantly. This was probably due to 144
102 intravenous administration of enrofloxacin, which inhib- 145
103 its the growth of enteric organisms that synthesize en- 146
104 zyme urease responsible for breakdown of urea to am- 147
105 monia in the G.I. tract. As a result, urea passes from 148
106 intestine to blood and elevates the blood urea level [7]. 149

107 The normal level of Hb, ESR and PCV of adult yak 150
108 are 12.8 g/dl, 6-10 mm/24 h and 36.0% respectively as 151
109 reported by Nivasarkar et al., 1998 [8]. In the present 152
110 study Hb, ESR and PCV values are consistent with the 153
111 previous report and there was no significance effect on 154
112 Hb, ESR and PCV values after intravenous administra- 155

122 REFERENCES

1231. Davidson, J; Lechtenberg, K.; Apley, M.D. and Bechtol, D.T.,
124 1998. Hematological profile after administration of enrofloxacin
125 in cattle. Freedom of Information Summary NADA 141-068.
1262. Javid, M.A.; Saeed, M.A.; Aleem, M. and Bashie, I.N., 2004.
127 Effect of gentamicin and enrofloxacin on uterine characteristics,
128 hematology, cervical mucus and pregnancy rate in postpartum
129 endometritic nili-ravi buffaloes. Buffalo Journal , 3, 279-288.
1303. Rao, G.S.; Ramesh, S.; Ahmad, A.H.; Tripathi, H.C.; Sharma,
131 L.D. and Malik, J.K., 2000. Effect of endotoxin -induced fever
132 and probencid on disposition of enrofloxacin and its metabolite
133 ciprofloxacin after intravenous administration of enrofloxacin in
134 goats. J Vet Pharm & Therapeutics , 23(6), 365.
1354. Prasad, B., 1992. Veterinary Clinical Diagnostic Technology. 1st
136 edn. CBS Publishers and Distributors, Nazia Printers, Lal Quan,
137 Delhi
1385. Barker, S. and Simsaung, P., 1965. Practical Clinical Biochemis-
139 try. 4th edn, Gulab Vazirani Pvt Ltd., New Delhi.
1406. Lehninger, A.L., 1978. Biochemistry 2nd edition, Kalyani Pub-
141 lishers, New Delhi.
1427. Kaneko, J.J., 1989. Clinical Biochemistry of Domestic Animal.
143 4th edition, Academic Press, London.
1448. Nivsarkar, A.E.; Gupta, S.C. and Gupta, N., 1997. Anatomy and
145 Physiology of Yak. Yak Production. Directorate of Information
146 and Publication on agriculture, ICAR, New Delhi: 171.
1479. Nath, N.C.; Aziz, A. and Bhattacharya, M., 2000. Comparative
148 serum protein pattern of mithun ((Bos frontalis) , yak (Bos
149 grunniens) and zebu cattle (Bos indicus). Indian J. Anim. Sci,
150 70 (8), 843-844.

151 CURRENT AUTHOR ADDRESSES

152 Sanjib Khargharia, Department of Pharmacology and Toxicology,
153 West Bengal University of Animal and Fishery Sciences, Mohan-
154 pur, Nadia-741252. E-mail: kharukharu@sify.com (Correspond-
155 ing author)

156 Chandana Choudhury Barua, Department of Pharmacology and Toxi-
157 cology, College of Veterinary Science, Khanapara, AAU, Guwa-
158 hati-781022, Assam, India.

159 Nabin Nath, Department of Animal Physiology and Biochemistry,
160 College of Veterinary Science, Khanapara, AAU, Guwahati-
161 781022, Assam, India

162 M. Bhattacharya, Director NRC Yak, Dirang, Arunachal Pradesh, India.