

# Effect of Aqueous Extract of *Syzygium cumini* Pulp on Antioxidant Defense System in Streptozotocin Induced Diabetic Rats

NAMASIVAYAM REKHA, RAMACHANDRAN BALAJI and MUNUSWAMY DEECARAMAN

For author affiliations, see end of text.

Received January 6, 2008; Revised September 30, 2008; Accepted October 29, 2008

This paper is available online at <http://ijpt.iuims.ac.ir>

## ABSTRACT

The aqueous extract of *Syzygium cumini* (Pulp) was investigated for its possible hypoglycemic and antioxidant potentials in streptozotocin (STZ) induced diabetic wistar female rats. A single intraperitoneal injection of STZ at a dose of 55 mg/kg body weight elevated the glucose levels more than 230 mg/dl after 3 days. Treatment with the aqueous extract of *Syzygium cumini* (Pulp) at 100 mg/kg and 200 mg/kg body weight resulted in significant reduction ( $p < 0.001$ ) in blood glucose levels. Body weights were significantly reduced ( $p < 0.001$ ) in STZ Induced diabetic rats when compared to normal rats, while in diabetic rats *S. cumini* extract ( $p < 0.001$ ) prevented significantly the decrease in body weight in a dose dependant manner. Total cholesterol, phospholipids, triglycerides, LDL-c, HDL-c levels were altered in STZ induced diabetic rats, which were considerably restored to near normal in animals treated with *S. cumini* extract. The administration of aqueous extract of *Syzygium cumini* resulted in significant ( $p < 0.001$ ) increase in the levels of superoxide dismutase, catalase, glutathione peroxidase, glutathione-S-transferase and reduced glutathione (GSH), thus resulting in reduced free radical formation in liver tissues of the diabetic rats. The increased levels of lipid peroxidation and hydroperoxides in diabetic rats were reverted back to near normal levels after the treatment with aqueous extract of *Syzygium cumini*. These observations demonstrated that aqueous extract of *Syzygium cumini* (Pulp) have strong hypoglycemic effect and *in vivo* antioxidant activity in STZ-Induced diabetic rats and was dose dependent.

**Keywords:** Free radical, *Syzygium cumini*, Diabetic rats

Diabetes mellitus is a syndrome characterized by abnormal insulin secretion, derangement in carbohydrate and lipid metabolism, and is diagnosed by the presence of hyperglycemia [1]. The currently suggested mechanism underlying diabetes and diabetic complications is Oxidative stress [2]. In recent years, much attention has been focused on the role of oxidative stress, and it has been reported that oxidative stress may constitute the key and common event in the pathogenesis of secondary diabetic complications [3]. The role of free radical reactions in disease pathology is well established, suggesting that these reactions are necessary for normal metabolism but can be detrimental to health as well. During metabolism, oxygen consumption involves the constant generation of free radicals and reactive oxygen species. Free radicals are either generated by cellular metabolisms such as glycolysis, mitochondrial respiration, and xenobiotic detoxification or by exogenous factors such as redox reactions. Some are extremely reac-

tive and therefore interact with some "vital" macromolecules including lipids, nucleic acids and proteins [4]. There are many enzymatic and nonenzymatic antioxidant defence systems in the body that remove these toxic species. Enzymes such as superoxide dismutase, catalase, glutathione peroxidase, etc. are involved in this detoxification process [5]. Moreover, diabetes also induces changes in the activity of the antioxidant enzymes [6]. Accumulating evidence suggests that oxidative cellular injury caused by free radicals contributes to the development of diabetes mellitus [7]. Thus oxidative stress has been shown to have a role in the causation of diabetes type I and II and as such antioxidants may have a role in the alleviation of diabetes and related problems [8, 9].

Currently, there is growing interest in herbal remedies due to the side effects associated with the therapeutic agents (oral hypoglycemic agents and insulin) for the treatment of diabetes mellitus [10].

Recently, some synthetic antioxidants such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) have been suspected to be dangerous to human health [11]. Therefore there is an urgent need to search for novel antioxidants from natural sources, which could be used in medicine and additives to nutraceuticals [12]. Plants often contain substantial amounts of antioxidants, such as carotenoids, flavonoids and tannins [13] and we suggest that antioxidant action may be an important property of plant medicines for treatment of diabetes. Many indigenous Indian medicinal plants have been found to be useful to successfully manage diabetes [14].

Out of a large number of herbal drugs stated to possess anti-diabetic activity in the Auyurvedic system of medicine of India, *Syzygium cumini* of family *Myrtaceae* is being widely used to treat diabetes by the traditional practitioners over many centuries [15]. *Syzygium cumini* is commonly called as Jamun, Black plum or Indian Black berry. It is a large tree found in all forests over the greater part of India from the sub-Himalayan tract to extreme south. It is also found in Thailand and Philippines. The fruits of *Syzygium cumini* are oval to elliptical, 1.5–3.5 cm long, dark purple or nearly black, luscious, fleshy and edible [16]. The fruits are reported to contain vitamin C, gallic acid, tannins, anthocyanins, includes cyanidin-, petunidin, malvidin glucoside and other components. The juice of unripe fruits is used for preparing vinegar that is considered to be a stomachic, carminative and diuretic [17, 18]. So far the antihyperglycemic activity of seeds of *Syzygium cumini* has been well established [19–25] but there are only very few reports regarding the antihyperglycemic activity of fruit-pulp of *Syzygium cumini* [19, 22, 26]. Also studies' regarding the antioxidant potentials of *Syzygium cumini* pulp has not yet been reported. Thus an attempt was made to investigate the effect of *Syzygium cumini* pulp on antioxidant enzymes in liver of streptozotocin-induced diabetic rats. The efficacy was compared with the standard hypoglycemic drug, glyburide.

## MATERIALS AND METHODS

### Chemical

Streptozotocin (STZ) and glyburide were obtained from Sigma Chemical Co (St Louis, MO-USA). Biochemical kits and all other chemicals utilized were of analytical grade.

### Plant Materials

The plant material of *Syzygium cumini* fruit was procured from local market during the month of June. This was identified and authenticated by Dr. P. Jayaraman, Director, Plant Anatomy Research Centre (PARC), Chennai. The voucher sample (No. PARC/2006/11) was kept in our laboratory for reference.

### Preparation of Plant Extract

The ripe fruits of *Syzygium cumini* (1 Kg) were first washed well and the pulp was separated from the seeds. The pulp was ground for 10 min in a mixer along with

the distilled water (500 ml). It was allowed to stand overnight and then filtered through several layers of muslin cloth. The whole procedure was carried out in the cold condition at 4 °C. The filtrate was centrifuged in a refrigerated centrifuge at 10,000 rpm. The supernatant was lyophilized to get a thick paste of water extract. The yield of lyophilized water extract was about 8.9 g from 650 g of pulp, obtained from 1 kg fruits of *Syzygium cumini*.

### Animals

Female Wistar rats weighing 170±10 g obtained from Tamil Nadu Veterinary and Animal Sciences University, Chennai, India were used for the present investigation. The animals were maintained on standard rat feed supplied by Hindustan Lever Ltd., Bangalore, India. The experiments were conducted according to the ethical norms approved by Ministry of Social Justice and Empowerment, Government of India and Institutional Animal Ethics Committee Guidelines. Animals were housed in stainless steel cages and kept in a room where a 12 h light/dark cycle was maintained. Rats had free access to water and standard feed throughout the period of the experiment.

### Induction of Diabetes in Rats

After one week of acclimatization, the rats were subjected to overnight fasting. Diabetes was induced with a single intraperitoneal injection of STZ at a dose of 55 mg/kg body weight. The STZ was freshly dissolved in citrate buffer (0.01M, pH 4.5) [27]. The injection volume was prepared to contain 1.0 ml/kg [28]. The animals were allowed to drink 5% glucose solution overnight to overcome the drug induced hypoglycemia. After 3 days, blood glucose levels were measured and the animals with a glucose concentration of more than 230 mg/dL were classified as diabetic [29] and taken for the experiment. Administration of the aqueous extract of *Syzygium cumini* was started on 4<sup>th</sup> day after STZ injection and this was considered to be the 1<sup>st</sup> day of treatment, which was continued for 15 days.

### Experimental Design

Thirty female Wistar rats were used in this study. The rats were randomized and divided into five groups of six animals each.

Group I: Control rats, received citrate buffer (0.01M, pH 4.5).

Group II: Diabetic controls, received STZ (55 mg/kg body weight, i.p.) once.

Group III: Diabetic rats, receiving aqueous extract of *Syzygium cumini* (100 mg/kg body weight) orally for 15 days.

Group IV: Diabetic rats, receiving aqueous extract of *Syzygium cumini* (200 mg/kg body weight) Orally for 15 days.

Group V: Diabetic rats receiving 5 mg/kg body weight of Glyburide orally for 15 days.

At the end of the experiment, all the rats were decapitated after fasting for 16 hours. Blood was

**Table 1.** Effect of aqueous extract of *Syzygium cumini* on body weight in STZ induced diabetic rats.

Treatment	Body weight (g)			
	Initial	5 <sup>th</sup> day	10 <sup>th</sup> day	15 <sup>th</sup> day
Vehicle control (group I)	174.38 ± 8.2	179.47 ± 6.8	183.26 ± 7.3	191.73 ± 9.4
Diabetic control (group II)	176.83 ± 7.1a*	142.14 ± 6.2a*	139.41 ± 5.8a*	132.14 ± 5.1a*
Diabetic + 100mg/kg <i>S.cumini</i> (group III)	175.37 ± 6.4b*	148.21 ± 7.6b*	157.6 ± 7.4b*	163.49 ± 6.1b*
Diabetic + 200mg/kg <i>S.cumini</i> (group IV)	175.23 ± 7.8c*	151.26 ± 5.1c*	161.91 ± 9.7c*	168.28 ± 7.2c*
Diabetic + 5mg/kg glyburide (group V)	178.28 ± 9.2d*	156.73 ± 6.3d*	164.14 ± 7.2d*	173.38 ± 5.4d*

Values are given as Mean ± SEM of six animals in each group.

a Comparison of Group I vs Group II.

b,c, d Comparison of Group III, Group IV & Group V vs Group II.

\**p*<0.001 statistically significant (ANOVA followed by Dunnett's t-test), when diabetic control was compared with the vehicle control and extract treated groups were compared with the diabetic control.

collected without anticoagulant to separate serum for various biochemical estimations. The liver was dissected out and cleared off blood. This was immediately transferred to ice-cold containers containing 0.9% NaCl and homogenized in 0.1N Tris-HCl buffer (pH 7.4). The tissue homogenates were used for the following estimations: Thiobarbituric acid reactive substances (TBARS) and hydroperoxides were estimated according to method of Ohkawa et al [30] and Jiang et al [31] respectively. Reduced glutathione (GSH) was estimated by the method of Ellman [32]. Protein was estimated by the method of Lowry et al [33]. The activity of superoxide dismutase (SOD) was assayed by the method of Misra et al [34]. The activity of glutathione peroxidase (GPx) was assayed according to the method of Rotruck et al [35]. Catalase (CAT) activity was assayed by the method of Sinha [36]. Glutathione-S-transferase (GST) was estimated by the method of Habig et al [37].

**Measurement of Body weight & Blood Glucose Level**

The body weight and blood glucose level were measured at about every 5 days interval. Blood samples were obtained by tail vein puncture of both the normal and STZ induced diabetic rats. Blood glucose level was measured by single touch glucometer.

**Serum Total Cholesterol**

Serum total cholesterol (TC) was quantified by spec-

trophotometric method [38] by the addition of enzyme present in reagent kit. The absorbance of red quinoneimine complex was determined at 505 nm. The value of TC present in serum was expressed in mg/dL.

**Serum Lipoprotein Cholesterol**

Serum LDL-c was measured according to protocol of Friendswald et al [39]. Serum HDL-c was measured by the method of Waenic et al [40].

**Serum Triglyceride**

Serum triglyceride was measured by using kit. The absorbance was noted at 520 nm. The value was expressed in the unit of mg/dL [41].

**Serum Phospholipids**

Serum phospholipids was determined by the method of Zilversmit et al [42]

**Statistical analysis**

Statistical evaluation of data was performed by using one-way analysis of variance (ANOVA) followed by Dunnett's t-test [43]. *p*-values < 0.05 were considered as significant.

**RESULTS**

**Blood Glucose Level and Body Weight Changes**

The body weight changes in diabetic group was significantly decreased (*p* <0.001) when compared with the

**Table 2.** Effect of aqueous extract of *Syzygium cumini* on blood glucose levels in STZ induced diabetic rats.

Treatment	Blood glucose level (mg/dL)				
	Initial day	1 <sup>st</sup> day	5 <sup>th</sup> day	10 <sup>th</sup> day	15 <sup>th</sup> day
Vehicle control (group I)	88.32 ± 5.2	86.91 ± 6.4	87.51 ± 4.8	89.26 ± 3.2	86.72 ± 4.8
Diabetic control (group II)	87.21 ± 5.6a*	239.78 ± 9.4a*	256.32 ± 8.1a*	274.28 ± 6.2a*	281.63 ± 6.9a*
Diabetic + 100mg/kg <i>S.cumini</i> (group III)	85.61 ± 4.8b*	236.81 ± 7.2b*	192.23 ± 4.1b*	159.23 ± 8.2b*	127.16 ± 5.4b*
Diabetic + 200mg/kg <i>S.cumini</i> (group IV)	87.30 ± 4.2c*	234.91 ± 6.2c*	166.27 ± 5.3c*	127.42 ± 5.8c*	114.56 ± 5.1c*
Diabetic + 5mg/kg glyburide (group V)	86.49 ± 5.8d*	198.38 ± 4.1d*	142.29 ± 6.2d*	112.3 ± 5.4d*	98.68 ± 4.7d*

Values are given as Mean ± SEM of six animals in each group.

a Comparison of Group I vs Group II.

b,c, d Comparison of Group III, Group IV & Group V vs Group II.

\**p*<0.001 statistically significant (ANOVA followed by Dunnett's t-test), when diabetic control was compared with the vehicle control and extract treated groups were compared with the diabetic control.

**Table 3.** Effect of aqueous extract of *Syzygium cumini* on serum lipid profile in STZ induced diabetic rats.

Treatment	Serum Total cholesterol (mg/dL)	Serum HDL (mg/dL)	Serum LDL (mg/dL)	Serum Triglycerides (mg/dL)	Serum Phospholipid (mg/dL)
Vehicle control (group I)	128.72 ± 4.8	45.73 ± 3.2	46.56 ± 5.2	78.63 ± 6.1	104.61 ± 5.9
Diabetic control (group II)	233.61 ± 7.2a*	27.91 ± 4.8a*	126.62 ± 4.7a*	183.16 ± 9.2a*	63.28 ± 4.7a*
Diabetic + 100mg/kg <i>S.cumini</i> (group III)	188.82 ± 6.1b*	34.57 ± 5.3b*	83.38 ± 4.2b*	131.56 ± 4.8b*	85.41 ± 4.9b*
Diabetic + 200mg/kg <i>S.cumini</i> (group IV)	164.48 ± 5.2c*	40.22 ± 4.6c*	48.26 ± 3.8c*	98.23 ± 5.1c*	91.58 ± 5.6c*
Diabetic + 5mg/kg glyburide (group V)	136.28 ± 5.9d*	41.27 ± 3.8d*	42.81 ± 3.2d*	70.22 ± 4.6d*	99.23 ± 5.4d*

Values are given as Mean ± SEM of six animals in each group.

a Comparison of Group I vs Group II.

b,c, d Comparison of Group III, Group IV & Group V vs Group II.

\* $p < 0.001$  statistically significant (ANOVA followed by Dunnett's t-test), when diabetic control was compared with the vehicle control and extract treated groups were compared with the diabetic control.

normal control, which then returned to near normal in diabetic rats treated with aqueous extract of *Syzygium cumini* at a dose of 100 and 200 mg/kg body weight. The results were shown in (Table 1).

The results of blood glucose level changes in normal, STZ induced diabetic rats and *Syzygium cumini* treated diabetic rats were shown in (Table 2). There was a significant ( $p < 0.001$ ) increase in blood glucose levels in STZ induced diabetic rats (group II) when compared with normal rats. Administration of aqueous extract of *Syzygium cumini* at a dose of 100 and 200 mg/kg body weight significantly ( $p < 0.001$ ) decreased blood glucose level in STZ induced diabetic rats (group III and IV). The results were found to be in a dose dependent manner, comparable with that of standard glyburide.

### Serum Lipids

The changes in the level of serum lipids in control and experimental rats are illustrated in (Table 3). The total-cholesterol, LDL-Cholesterol and triacylglycerol significantly increased and HDL-Cholesterol, phospholipids significantly decreased in STZ induced diabetic rats (group II) ( $p < 0.001$ ) when compared with the normal (group I) rats. The aqueous extract of *Syzygium cumini* (100 mg/kg and 200 mg/kg body weight) offered a significant protection against alteration in the serum

lipids of diabetic rats. The results were also dose dependent. The results were comparable with that of standard glyburide.

### Lipid peroxidation and Hydroperoxides

The levels of MDA in liver were significantly ( $p < 0.001$ ) increased in STZ induced diabetic rats as compared to normal rats. Treatment with aqueous extract of *Syzygium cumini* resulted in significant decrease in the level of lipid peroxidation products (MDA) and Hydroperoxides in diabetic rats. The results were shown in (Fig. 1 & 2).

### Antioxidant enzymes

A significant decrease ( $p < 0.001$ ) in the activities of antioxidant enzymes such as SOD and CAT were observed in the liver of STZ induced diabetic rats when compared with that of normal rats. Upon administration of 100 and 200mg/kg body weight of aqueous extract of *Syzygium cumini*, the activities of both SOD and CAT were significantly reversed to near normal. The levels of reduced glutathione, Glutathione peroxidase, Glutathione-S-transferase were significantly depleted in STZ induced diabetic rats. Treatment with aqueous extract of *Syzygium cumini* significantly increased the levels of these antioxidant enzymes in diabetic rats. The results were shown in (Table 4).

**Table 4.** Effect of aqueous extract of *Syzygium cumini* on antioxidant enzymes in STZ induced diabetic rat liver.

Treatment	SOD	CAT	GPx	GSH-Rase	GST
Vehicle control (group I)	7.31 ± 0.13	54.62 ± 6.2	7.63 ± 0.21	0.971 ± 0.06	8.15 ± 0.61
Diabetic control (group II)	3.72 ± 0.08a*	18.59 ± 6.8a*	5.19 ± 0.34a*	0.492 ± 0.05a*	4.58 ± 0.47a*
Diabetic + 100mg/kg <i>S.cumini</i> (group III)	4.87 ± 0.25b*	36.38 ± 7.3b*	6.25 ± 0.25b*	0.710 ± 0.02b*	6.24 ± 0.43b*
Diabetic + 200mg/kg <i>S.cumini</i> (group IV)	5.94 ± 0.16c*	49.26 ± 5.6c*	6.92 ± 0.38c*	0.823 ± 0.04c*	7.53 ± 0.53c*
Diabetic + 5mg/kg glyburide (group V)	6.98 ± 0.15d*	51.83 ± 5.9d*	7.12 ± 0.12d*	0.91 ± 0.05d*	7.89 ± 0.58d*

Values are given as Mean ± SEM of six animals in each group.

a Comparison of Group I vs Group II.

b,c, d Comparison of Group III, Group IV & Group V vs Group II.

\* $p < 0.001$  statistically significant (ANOVA followed by Dunnett's t-test), when diabetic control was compared with the vehicle control and extract treated groups were compared with the diabetic control.

### Units

SOD - Units/mg protein.


1 unit - The amount of enzyme required bring about 50% of inhibition of autooxidation of epinephrine.

GPx -  $\mu$  moles of GSH oxidized/min/mg protein.

GSH - n moles of DTNB conjugated /mg protein.

CAT - n moles of hydrogen peroxide decomposed/min/mg protein.

GST - n moles of CDNB conjugate formed/min/mg protein.



**Fig 1.** Effect of aqueous extract of *Syzygium cumini* on lipid peroxidation in STZ induced diabetic rat liver


**DISCUSSION AND CONCLUSION**

The present study is assessment of antioxidant and antihyperglycemic activity of aqueous extract of pulp of *Syzygium cumini* on female wistar rats. STZ is toxic to pancreatic β-cells and is thus widely used for induction of experimental diabetes mellitus in animals, resulting in the production of ROS [44]. STZ causes a significant elevation in the level of blood glucose in rats. Administration of 100 and 200 mg/kg body weight of aqueous extract of *Syzygium cumini* Pulp significantly decreased the blood glucose level in these rats suggesting that it has hypoglycemic properties. The decreased body weight in diabetic rats is due to excessive breakdown of tissue proteins [45]. Treatment with *Syzygium cumini* improved body weight significantly in a dose dependent manner, indicating prevention of muscle wasting due to hyperglycemic condition.

The rise in blood sugar is accompanied with the increase in TC, LDL-c, TG and fall of HDL-c & phospholipids. Diabetes is also known to be associated with an increase in the synthesis of cholesterol, which may be due to the increased activity of HMG CoA reductase [46]. Increased LDL-cholesterol may arise from glycosylation of the lysyl residues of apoprotein B as well as from decreasing affinity for the LDL receptor and hence, decreased metabolism [47]. A number of observations indicate that plasma HDL cholesterol is low in untreated insulin-deficient diabetics [48], which was associated with a decline in HDL-turnover rate. Further the HDL-cholesterol levels correlate with lipoprotein lipase (LPL) levels in IDDM patients [49]. Hypertriglyceridemia is a common finding in patients with diabetes mellitus and is responsible for vascular compli-

cations [50]. It has been reported that deficiency of lipoprotein lipase (LPL) activity may contribute significantly to the elevation of triglycerides in diabetes [51]. However, Oral administration of aqueous extract of pulp exhibited hypocholesterolemic and hypotriglyceridemic effects while at the same time increasing HDL-c, possibly by controlling the hydrolysis of certain lipoproteins and their selective uptake and metabolism by different tissues. This implies that *Syzygium cumini* pulp can prevent or be helpful in reducing the complications of lipid profile seen in some diabetics in whom hyperglycaemia and hypercholesterolemia coexist quite often.

Induction of diabetes in rats with STZ uniformly results in an increase in lipid peroxidation (TBARS), an indirect evidence of intensified free radical production [52]. Most of the tissue damage is considered to be mediated by these free radicals by attacking membranes through peroxidation of unsaturated fatty acids [53]. The concentration of lipid peroxidation products may reflect the degree of oxidative stress in diabetes [54]. In the present study the concentrations of lipid peroxides and hydroperoxides were significantly increased in liver of diabetic rats, indicating an increase in the generation of free radicals. An observed increase in the level of TBARS in liver may be due to increased susceptibility of the tissue of diabetic rats to lipid peroxidation [55]. Administration with *Syzygium cumini* protects the cells through attenuation of lipid peroxidation and decreased the production of free radical derivatives, as evident from the decreased levels of liver MDA and hydroperoxides. This suggests protective role of fruit pulp, which could be due to the antioxidative effect of polyphenols [56] present in the fruit pulp, which may act as strong superoxide radical and singlet oxygen quenchers.



**Fig 2.** Effect of aqueous extract of *Syzygium cumini* on hydroperoxides in STZ induced diabetic rat liver.

Values are given as Mean  $\pm$  SEM of six animals in each group.

a Comparison of Group I vs Group II.

b, c, d Comparison of Group III, Group IV & Group V vs Group II.

\* $p < 0.001$  statistically significant (ANOVA followed by Dunnet's t-test), when diabetic control was compared with the vehicle control and extract treated groups were compared with the diabetic control.

Reduced activities of SOD and CAT in liver of diabetic rats have been observed in our study. The decreased activities of SOD and CAT in liver during diabetes may be due to increased production of reactive oxygen radicals that can themselves reduce the activity of these enzymes [57]. SOD is an important defense enzyme, which converts superoxide radicals to hydrogen peroxide [58]. Increase in SOD activity could be due to its induction by increased production of superoxide, which has been implicated in cell dysfunction.  $H_2O_2$  has been reported to act as an inducer of tissue SOD [59]. CAT is a heme protein, which decomposes hydrogen peroxide and protects the tissues from highly reactive hydroxyl radicals [60]. The reduction in the activity of these enzymes may result in a number of deleterious effects. Any compound, natural or synthetic, with antioxidant properties, might contribute towards the partial or total alleviation of this damage. Therefore, removing  $O_2$  and  $OH^\cdot$  probably one of the most effective defenses of a living body against diseases [61]. Administration of *Syzygium cumini* increased the activity of enzymes and may help to avoid the free radicals generated during diabetes. The result of SOD and CAT activity clearly shows that the *Syzygium cumini* contains a free radical-scavenging activity, which could exert a beneficial action against pathological alterations caused by the presence of  $O_2$  and  $OH^\cdot$ . This action, predominantly due to the extract, could involve mechanisms related to scavenging activity.

Reduced glutathione (GSH) is a major endogenous antioxidant which counterbalance free radical mediated damage and it is well known that GSH is involved in the protection of normal cells structure and function by maintaining the redox homeostasis, quenching of free radicals and by participating in detoxification reactions [62]. GSH is also known to protect the cellular system against the toxic effects of lipid peroxidation [63]. The decrease in GSH level in liver during diabetes is proba-

bly due to its increased utilization by the hepatic cells which could be the result of decreased synthesis or increased degradation of GSH by oxidative stress in diabetes [64]. A relative depletion of NADPH due to aldose reductase activation and secondary to reduced production through the pentose cycle impairs GSH regeneration and leads to depletion of this free radical scavenger [65]. GSH reacts with free radicals and is a crucial substrate for GPx and GST, which takes part in the cellular defense mechanisms against intermediate oxygenated products of metabolism. GPx, an enzyme with selenium and GST, catalyzes the reduction of hydrogen peroxide to non-toxic compounds [66]. GPx metabolizes hydrogen peroxide to water by using GSH as a hydrogen donor [67]. The reduced activity of GPx may result in the accumulation of toxic products due to oxidative damage. The activities of GPx and GST were observed to decrease significantly in diabetic rats. Treatment with *Syzygium cumini* significantly recovered the GSH content and increase in the levels of GSH dependent enzymes GPx, GST, which clearly indicates the protective effect of *Syzygium cumini* on antioxidants.

Hyperglycemia is the prominent cause for the production of free radicals, which leads to development of diabetic complications. Oxidative stress is currently suggested as mechanism underlying diabetes and diabetic complications. Many of the complications of diabetes including retinopathy and atherosclerotic vascular disease, the leading cause of mortality in diabetes have been linked to oxidative stress and antioxidants have been considered as treatments [68, 69]. Flavonoids, tannins, anthocyanins,  $\beta$ -carotene and other phenolic constituents present in food of plant origin are potential antioxidants and may play a beneficial role in the prevention of several chronic disorders [70, 71].

The fruits of *Syzygium cumini* are reported to contain vitamin C, gallic acid, tannins, anthocyanins cyanidin,

glucoside, petunidin, malvidin and other components [18]. Several authors reported that, saponins, flavonoids, phenolic compounds, glycosides, triterpenoids and vitamin C have potent antioxidant, hypolipidemic and hypocholesterolemic effects [72, 73, 74, 75]. *Syzygium cumini* kernels have been reported to ameliorate the oxidative stress in diabetic condition [76] and also the fruit skin of *Syzygium cumini* has been reported to possess *invitro* antioxidant activity [77]. In our present investigation treatment with *Syzygium cumini* pulp in rats induced with STZ restored the fatty acid composition to near normal level and this may be due to the decreased toxicity of reactive oxygen species by its anti-oxidant property. Further it can resist, to an extent, the formation of lipid peroxy radicals in a number of tissues, thereby protecting these tissues. In this context a number of other plants have also been observed to have hypoglycaemic and insulin-release stimulatory effects [78]. The results obtained suggest that pulp of *Syzygium cumini* possesses potent antioxidant and antidiabetic activity; this may be due to the antioxidant phytochemicals present in the fruit of *Syzygium cumini*, which can scavenge free radicals and prevent the depletion of endogenous antioxidants. Thus consumption of *Syzygium cumini* pulp may be considered as a potential source of natural antioxidant and antihyperglycemic activity that helps in reducing oxidative stress in diabetic condition, which may provide beneficial role in the management of diabetes.

**REFERENCES**

1. Zimmet PZ, McCarty DJ, DeCourten M.P. The global epidemiology of non-insulin-dependent diabetes mellitus and the metabolic syndrome. *J Diabet Complications*. 1997;11:60-8.
2. Halliwell B, Gutteridge JMC. Free radicals in biology and medicine. 2<sup>nd</sup> ed. Oxford: Clarendon Press; 1989.
3. Ceriello A. Oxidative stress and glycemic regulation. *Metabolism*. 2000;49:27-9.
4. Nia R, Paper DH, Essien EE, Oladimoji OH, Iyadi KC, Franz B. Investigation into *in vitro* radical scavenging and *in vivo* anti-inflammatory potential of *Tridax procumbens*. *Nig J Physiol Sci*. 2003;18(1-2):39-43.
5. Clark IA, Cowden WB, Hunt NH. Free radical induced pathology. *Med Res Rev*. 1985;5:297-332.
6. Genet S, Kale RK, Baguez NZ. Alteration on antioxidant enzymes and oxidative damage in experimental diabetic rat tissue: effect of vanadate and fenugreek (*Trigonella faenum graeoum*). *Mol Cell Biochem*. 2002;236:7-125.
7. Baimbolkar S, Sainani GS. Evaluation of oxidative stress in diabetic with or without vascular complications. *J Assoc Phys India*. 1995;43:10-12.
8. John WB. Role of oxidative stress in development of complications in diabetes. *Diabetes*. 1991;40:405.
9. Hunt JV, Smith CCT, Wolff SP. Autooxidative glycosylation and possible involvement of peroxides and free radicals in LDL modification by glucose. *Diabetes*. 1990;39:1420-4.
10. Kameswara Rao B, Giri R, Kesavulu MM, Apparao C. Herbal medicine: In the management of diabetes mellitus. *Manphar Vaidhya Patrika*. 1997;I: p. 33-5.
11. Safer AM, Al-Naghamigh AJ. Hepatotoxicity induced by the antioxidant food additive BHT in rats. An electromicroscopic study. *Histol Histopathol*. 1999;14:391-406.
12. Thomes PG, Wade AM. Histidine containing nutraceuticals. Official Gazette of the United States and Trade mark office patents. 2001;2:1253.
13. Larson RA. The antioxidants of higher plants. *Phytochemistry*. 1988;27:969-78.
14. Joy KL, Kuttan R. Anti-diabetic activity of *Picrorrhiza kurroa* extract. *J Ethnopharmacol*. 1999;167:143-8.
15. Nadkarni AK. In: Indian Materia Medica. Vol I Popular Prakashan. Bombay, India, 1954. p. 1331.
16. Chopra RN, Chopra IC, Handa KL, Kapur LD. Indigenous Drugs of India. 2<sup>nd</sup> ed. U.N. Dhur and Sons Pvt. Ltd, Calcutta. p. 1958686-9.
17. Martinez SB, Del Valle MJ. Storage stability and sensory quality of duhat (*Syzygium cumini* Linn.) anthocyanins as food colorant. In: Mazza G, Miniati E. Anthocyanins in fruits, vegetables and grains. London: CRC Press; 1993. p. 139-47.
18. Wealth of India. Raw materials Vol X, CSIR New Delhi, 1976. p. 100-4.
19. Shrotri DS, Kelkar M, Deshmukh VK, Aiman R. Investigations of the hypoglycemic properties of *Vinca rosea*, *Cassia auriculata* and *Eugenia jambolana*. *Indian J Med Res*. 1963;51:464-7.
20. Bansal R, Ahmad N, Kidwai JR. Effect of oral administration of *Eugenia jambolana* seeds and chlorpropamide on blood glucose level and pancreatic cathepsin B in rats. *Indian J Biochem Biophys*. 1981;18:377-81.
21. Kohli KR. A study on Kriyakala of diabetes mellitus and its treatment with *Eugenia jambolana* MD (Ay) [dissertation]. Institute of Medical Sciences: Banaras Hindu University, Varanasi (India); 1983.
22. Achrekar S, Kaklij GS, Pote MS, Kelkar SM. Hypoglycemic activity of *Eugenia jambolana* and *Ficus bengalensis*: Mechanism of action. *In Vivo*. 1991;5:133-48.
23. Grover JK, Vats V, Rathi SS. Antihyperglycemic effect of *Eugenia jambolana* and *tinospora cordifolia* in experimental diabetes and their effects on key metabolic enzymes involved in carbohydrate metabolism. *J Ethnopharmacol*. 2000;73:461-70.
24. Vikrant V, Grover JK, Tandon N, Rathi SS, Gupta N. Treatment with extracts of *Momordica charantia* and *Eugenia jambolana* prevents hyperglycemia and hyperinsulinemia in fructose fed rats. *J Ethnopharmacol*. 2001;76:139-43.
25. Sharma SB, Nasir A, Prabhu KM, Dev G, Murthy PS. Hypoglycemic and hypolipidemic effect of ethanolic extracts of seeds of *Eugenia jambolana* in alloxan-induced diabetic model of rabbits. *J Ethnopharmacol*. 2003;85:201-06.
26. Suman BS, Afreena N, Krishna MP, Pothapragada SM. Antihyperglycemic effect of the fruit-pulp of *Eugenia jambolana* in experimental diabetes mellitus. *J Ethnopharmacol*. 2006;104:367-73.
27. Ozsoy-Sacan O. Effects of chard (*Beta vulgaris Lvarcicla*) extract on pancreatic  $\beta$ -cells in STZ-diabetic rats: morphological and biochemical study. *J Ethnopharmacol*. 2000;73:251-9.
28. Murali B, Upadhyaya UM, Goyal RK. Effect of chronic treatment with *Enicostemma littorale* in non-insulin dependent diabetic (NIDDM) rats. *J Ethnopharmacol*. 2002;81:199-204.
29. Cetto AA, Weidonfeld H, Revilla MC, Sergio IA. Hypoglycemic effect of *Equisetum mriochaetum* aerial parts on STZ-diabetic rats. *J Ethnopharmacol*. 2000;72:129-33.
30. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem*. 1979;95:351-8.
31. Jiang ZY, Hunt JV, Wolff SD. Ferrous ion oxidation in the presence of xylenol orange for detection of lipid hydroperoxides in low-density lipoprotein. *Anal Biochem*. 1992;202:384-9.
32. Ellman GC. Tissue sulfhydryl groups. *Arch Biochem Biophys*. 1959;82:70-7.
33. Lowry OH, Rosenbrough NJ, Farr AL, Randall R. Protein measurement with Folin phenol reagent. *J Biol Chem*. 1951;193:265-9.

34. Misra HP, Fridovich I. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem.* 1972;247:3170-5.
35. Rotruck JT, Pope AL, Gaspher HE, Hafeman DG, Hoekstra WG. Selenium biochemical role as a component of glutathione peroxidase. *Science.* 1973;179:588-90.
36. Sinha KA. Colorimetric assay of catalase. *Anal Biochem.* 1972;47:389-94.
37. Habig WH, Pabst MJ, Jakoby WB. Glutathione-transferases. The first enzymatic step in mercapturic acid formation. *J Biol Chem.* 1974;249(22):7130-9.
38. Allain CC, Poon LS, Chan CSG. Enzymatic determination of total serum cholesterol. *Clin Chem.* 1974;20:470-5.
39. Friendwald WT, Levy KJ, Frederickson DS. Estimation of concentration of LDL in plasma without use of preparative ultracentrifuge. *Clin Chem.* 1972;18:499-502.
40. Waenic RG, Albers JJ. A comprehensive evaluation of the heparin manganese precipitation procedure for estimating high density lipoprotein cholesterol. *J Lipid Res.* 1978;19:65-76.
41. McGowan MW, Aritiss ZE, Strandbergh DR, Zak B. A peroxidase-coupled method for the colorimetric determination serum triglycerides. *Clin Chem.* 1983;29:538-42.
42. Zilversmit DB, Davies AK. Micromethod for the determination of plasma phospholipid by trichloroacetic acid precipitation. *J Lab Clin Med.* 1950;35:155-160.
43. Duncan BD. Multiple range tests for correlated and heteroscedastic means. *Biometrics.* 1957;13:359-64.
44. Mazunder UK, Gupta M, Rajeshwar Y. Antihyperglycaemic effect and antioxidant potential of *Phyllanthus nuriri* (Euphorbiaceae) in streptozotocin induced diabetic rats. *Eur Bull Drug Res.* 2005;13(1):15-23.
45. Chatterjea MN, Shinde R. Text Book of Medical Biochemistry. 5th ed. New Delhi: Jaypee brothers, Medical publishers Ltd; 2002. p. 317.
46. Goodman MW, Michels LD, Keane WF. Intestinal and hepatic cholesterol synthesis in the alloxan diabetic rats. *Proc Soc Exp Biol Med.* 1982;170:286-90.
47. Golay A, Chen YD, Reaven GM. Effect of differences in glucose tolerance on insulin stability regulate carbohydrate and free fatty acid metabolism in obese individuals. *J Clin Endocrinol Metab.* 1986;62:1081-8.
48. Glasgow AM, August GP, Hung W. Relationship between control and serum lipids in juvenile-onset diabetes. *Diabetes Care.* 1981;4:76-80.
49. Nikkila EA, Huttunen JK, Ehnholm C. Postheparin plasma lipoprotein lipase and hepatic lipase in diabetes mellitus. Relationship to plasma triglyceride metabolism. *Diabetes.* 1977;26:11-21.
50. Kudchodkar BJ, Lee JC, Lee SM, DiMarco NM, Lacko AG. Effect on cholesterol homeostasis in diabetic rats. *J Lipid Res.* 1988;29:1272-87.
51. Bruan JEA, Severson DL. Lipoprotein lipase release from cardiac myocytes is increased by decavandate but not insulin. *Am J Physiol.* 1992;262 :E663-E70.
52. Maritim AC, Sanders RA, Watkins JB. Diabetes, Oxidative Stress and antioxidants: A review. *J Biochem Mol Toxicol.* 2003;17(1):24-38.
53. Stringer MD, Gorgo PG, Freeman A, Kakkar VV. Lipid peroxides and atherosclerosis. *Br Med J.* 1989;298:281-4.
54. Baynes JW. Role of oxidative stress in development of complications in diabetes. *Diabetes.* 1991;40:405-12.
55. Matkovic B, Kotorman M, Sz Varga I, Quy Hai D, Varga C. Oxidative stress in experimental diabetes induced by streptozotocin. *Acta Physiologica Hungarica.* 1998;85(1):29-38.
56. Banerjee A, Dasgupta N, De B. In vitro study of antioxidant activity of *Syzygium cumini* fruit. *Food Chem.* 2005;90:727-33.
57. Wohaieb SA, Godin DV. Alterations in free radical tissue-defense mechanisms in streptozotocin-induced diabetes in rats. *Diabetes.* 1987;36:1014-8.
58. McCrod JM, Keele BB, Fridovich J Jr. An enzyme based theory of obligate anaerobiosis: The physiological functions of superoxide dismutase. *Proc Natl Acad Sci U.S.A.* 1996;68:1024-7.
59. Matkovic B. Effect of plant and animal lesions on superoxide dismutase activities. In: McCord, Michelson AM, Fridovich I, eds. Superoxide and superoxide dismutase. New York: Academic press; 1977.p. 501-15.
60. Chance B, Green Stein DS, Roughton RJW. The mechanism of catalase action I-steady state analysis. *Arch Biochem Biophys.* 1952;37:301-9.
61. Lin JM, Lin CC, Fengchen M, Ujji T, Takadu A. Scavenging effects of *Mallotus repandus* on active oxygen species. *J Ethnopharmacol.* 1995;46:175-181.
62. Pari L, Amadi R. Protective role of tetrahydrocurcumin (THC) an active principle of turmeric on chloroquine induced hepatotoxicity in rats. *J Pharm Pharmaceut Sci.* 2005; 8(1):115-23.
63. Nicotera P, Orrenius S. Role of thiols in protection against biological reactive intermediates. *Adv Exp Med Bio.* 1986;197:41-9.
64. Loven D, Schedl H, Wilson H, Daabees TT, Stegink LD, Diekus M, Oberley L. Effect of insulin and oral glutathione on glutathione levels and superoxide dismutase activities in organs of rats with streptozotocin induced diabetes. *Diabetes.* 1986;35:503-7.
65. Dominguez Z, Ruiz E, Gussinye M, Carrascosa A. Oxidative stress at onset and in early stages of type 1 diabetes in children and adolescents. *Diabetes Care.* 1998;21:1736-42.
66. Illing EKB, Gray CH, Lawrence RD. Blood glutathione and non glucose reducing substances in diabetes. *Biochem J.* 1991;48:637.
67. Sies H. Damage to plasmid DNA by singlet oxygen and its protection. *Mut Res.* 1993;299:183-91.
68. Cunningham JJ. Micronutrients as nutraceutical intervention in diabetes mellitus. *J Am Coll Nutr.* 1998;17:7-10.
69. Reaven PD, Herold DA, Barnett J, Edelman S. Effects of vitamin E on susceptibility of low-density lipoprotein and low-density lipoprotein subfractions to oxidation and on protein glycation in NIDDM. *Diabetes Care.* 1995;8:807-16.
70. Diplock AT, Charleux JL, Crozier-Willi G, Kok FG, Rice-Evans C, Roberfroid M, Stahl W, Vifia-Ribes J. Functional food science and defence against reactive oxidative species. *Br J Nutr.* 1998;80(1):S77-112.
71. Saskia ABE, Van Acker S, Van de Berg D, Tromp M, Griffioen D, Van Bennekom W, Van der vijgh W, Bast A. Structural aspect of antioxidant activity of flavonoids. *Free Radic Biol Med.* 1996;3:331-42.
72. Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB. The relative antioxidant activity of plant derived polyphenolic flavonoids. *Free Radical Res.* 1995;22:375- 83.
73. Sroka Z, Cisowski W. Hydrogen peroxide scavenging, antioxidant and antiradical activity of some phenolic acids. *Food Chem. Toxicol.* 2003;41:753-58.
74. Rupasinghe HP, Jackson CJ, Poysa V, DiBerado C, Bewely JD, Jenkinson J. Soyasapogenol A and B distribution in Soybean (Glycine Max L.Merr) in relation to seed physiology, genetic variability and growing location. *J. Agric. Food Chem.* 2003;5(20):5888-94.
75. Leontowicz H, Gorinstein S, Lojek A, Leontowicz M, Ciz M, Soliva-Fortuny R. Comparative content of some bioactive compounds in apples, peaches, and pears and their influence on lipids and antioxidant capacity in rats. *J Nutr Biochem.* 2002;13:603-10.
76. Kasiappan R, Balasubramanian R, Sorimuthu S. Effect of *Eugenia Jambolana* seed kernel on antioxidant defense system in streptozotocin-induced diabetes in rats. *Life Sci.* 2004;75:2717-31.
77. Archana B, Nabasree D, Bratati D. In vitro study of antioxidant activity of *Syzygium cumini* fruit. *Food Chem.* 2005;90:727-33.
78. Gupta SS. Prospects and perspectives of natural plant products in medicine. *Indian J Pharmacol.* 1994;26:5-9.



**CURRENT AUTHOR ADDRESSES**

Namasivayam Rekha, Dr.MGR Educational & Research Institute,  
Maduravoyal, Chennai-95. E-mail:  
[rekhanamachivayam@gmail.com](mailto:rekhanamachivayam@gmail.com) (Corresponding author)

Ramachandran Balaji, Dr.MGR Educational & Research Institute,  
Maduravoyal, Chennai-95.

Munuswamy Deecaraman, Dr.MGR Educational & Research Institute,  
Maduravoyal, Chennai-95.