

Review Article

IRANIAN JOURNAL OF PHARMACOLOGY & THERAPEUTICS Copyright © 2018 by Iran University of Medical Sciences



Iranian J Pharmacol Ther. 2019 (September);17:1-10.

Biomarker efficacy in multiple sclerosis diagnosis and treatment

Fatemeh Ghasemi Sakha¹, Seyed Mohammad Moazzeni², Farnaz Etesam³, Amirreza Azimi Saeen^{4*}, Gholamhassan Vaezi⁵

- ¹ Department of Biology, Damghan Branch, Islamic Azad University, Damghan, Iran
- ² Department of immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran
- ³ Psychosomatic Medicine Research Center, Imam Khomeini Hospital, Tehran University of Medical Sciences, Tehran, Iran
- ⁴ MS Research Center, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran
- ⁵ Department of Biology, Damghan Branch, Islamic Azad University, Damghan, Iran

Please cite this article as:

Ghasemi Sakha F, Moazzeni SM, Etesam F, Azimi Saeen A, Vaezi Gh. Biomarker efficacy in multiple sclerosis diagnosis and treatment. Iranian J Pharmacol Ther. 2019 (September);17:1-10.

ABSTRACT

The goal in studies concerning biomarkers in autoimmune diseases is finding a marker which fluctuates in correlation with the disease's severity and settles within normal borders after effective treatment. This marker would later be used as an efficient tool in diagnosis and analysis of medicine clout. It seems the most cogent biomarkers are those measurable in serum or plasma. MS is a neurological disease common within the young adult population with a predictable course, often leading to life-immersing disability. The prognosis, however difficult and limited, is currently possible via diagnostic tests (brain MRI) or clinical information (severity and degrees of disability). Many studies have been conducted in an effort to detect an adequate biomarker. It is still often overlooked that a reliable biomarker should both be clinically applicable and functional is prognosis; as this is what could lead to timely treatment.

Conflicts of Interest: Declared None Funding: None

Keywords

MS biomarkers, Autoimmune disease, Demyelination, Micrornas

Corresponding to:

Amirreza Azimi Saeen, Multiple Sclerosis Research Center, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran

Email: amirreza_azimi@yahoo.com

Received: 8 Apr 2019 Published:30 Sep 2019

INTRODUCTION

MS (Multiple Sclerosis) is an autoimmune disorder of unknown etiology which affects the central nervous system. This disease is usually observed in persons aged 20-45 [1-3] The disease is most prevalent in European countries and possesses average prevalence in Iran. There are four separate phenotypes identified for the disease; clinically isolated syndrome (CIS), relapsing-remitting MS (RPMS), primary progressive MS (PPMS), secondary progressive MS (SPMS) [4]. Different treatment programs are available for decelerating the disease's progress, disabling each attack and alleviating disturbances; namely the prescription of interferon. However, at this point, there is no definite cure for MS [5]. MS is a complex disorder with different factors impinging its mechanism [6].

We have categorized how biomarkers can be beneficial to the MS patient below:

- 1) Phenotype and severity diagnosis
- 2) Determining the course of the disease

3) Choosing a treatment method and predicting its success rate

4) Analyzing new treatment options

Studying biomarkers relevant to MS is heavily supported by MS International Federation and the Autoimmune Disease Committee. Many breakthroughs have been made in the field and the positive effects within the aforementioned categories are immediately visible. Studying biomarkers bears similarity in significance to the changes of gene expression, the disease's dependence on different biological molecules such as free radicals, lipids and peptides which are not perceptible via magnetic resonance imaging [7].

Biomarker Types in Multiple Sclerosis

We have grouped MS biomarkers in seven categories:

- 1. Biomarkers affecting the immune system
- a. Cytokines and their receptors
- b. Chemokines and their receptors
- c. Antibodies
- d. Biomarkers affecting the complement system
- e. adhesion molecules
- f. molecules affecting cell death
- 2. Biomarkers damaging the blood-brain barrier
- 3. Biomarkers damaging myelin
- 4. Stress oxidative biomarkers
- 5. Biomarkers damaging axon or nerve cells

6. Biomarkers increasing the production of astrocytes in the brain

7. Biomarkers restoring myelin

Alternatively, we can differentiate between the biomarkers by categorizing them into positive biomarkers and negative biomarkers. Positive biomarkers are either not produced in a physiological environment or produced in small amounts and in particular pathological circumstances. Negative biomarkers are produced in a physiological environment in certain tissues and are targeted at the manifestation of the disease. Namely, oligodendrocytes and myelin belong to this category. The aforementioned biomarkers are ideal for analysis before and after the manifestation or before and after treatments[8].

Immunological markers, markers contributing to the de-

ation of axon and also stress markers a

Ghasemi Sakha et al.

struction and restoration of axon and also stress markers are ideal as well. We can prescribe more specific medical treatments and assess their performance using these biomarkers. Table 1 expands on the prominence of biomarkers [9].

Considerations and Challenges in the Credibility of the Clinical Use of Biomarkers

The biomarker must apply to a set of basic rules when used in a clinical environment. First, it must be reliably and easily measured through a number of precise test. Secondly, the biomarker must be unambiguous and sensitive to the biology of the disease or its pathogenesis such as its inflammatory properties, its degree of nerve damage and ability to damage or restore myelin.

Biomarker Classification

Biomarkers can be classified based on their ability in prediction, diagnosis and correlation with the disease, as elaborated in as elaborated in Tables 1, 2, and 3.

Diagnostic Biomarkers

A diagnostic biomarker makes distinctions between different autoimmune diseases and differentiates between the healthy and unhealthy patient. This biomarker recognizes the probability of occurrence of a destructive nervous disease such as clinically isolated syndromes (CIS) [10] or radiologically isolated syndromes (RTS) [11]. Ideally, assimilated with the clinical disorder scale and radiological examination, the biomarker can contribute to an increase in sensitivity and appropriation. Most studies conducted belong to this class of biomarkers (Table 1). They are especially helpful in patients exposed to CIS which might later transform into MS.

Biomarkers Related to the Course of the Disease

Biomarkers differentiate between relapsing patients and patients diagnosed with secondary-progressive MS (SPMS).

Biomarker	CIS	RRMS	SPMS	PPMS	NMO
CSF OCB	Predicts	Present		Less	Less
	RRMS	in 90%		frequent	frequent
CD19+ B cells in CSF	Common	Common	Absent	Absent	
CD39+ Treg		Decreased	Normal		
CXCL3	Elevated	Elevated	Norma	Norma	
BDNF (CSF)		Highest	Elevated	Reduced	
Fetuin A			Active SPMS		
Neurofilament	Predicts				
chains (CSF)	RRMS				
NAA (CSF)		Increased	Lower		
Vitamin D		High		Low	
binding protein				High if	
Neopterin (urine)				stable	
	Predicts NMO				
Aquaporin-4	in LETM	Abcont			Present
antibody	severe or	Ausein			in 60-80%
	bilateral ON				

Table 1. Biomarkers associated with disease stages and clinical phenotypes

Table 2. Potential biomarkers of disease activity in MS			
Correlate	Biomarker		
	Cytokines		
Relapse (+), Disability (+)	TNF- α (serum and CSF)		
Relapse (-), Treatment Response	IL-10 (serum and CSF)		
Relapse (+)	IL-12 (serum and CSF)		
Relapse (+), Treatment Response	IL-17 (serum and CSF)		
Relapse (+), Treatment Response	IFN-γ		
Disease activity (+)	Osteopontin (CSF)		
NMO severity (+)	IL-6 (CSF)		
	IFN=interferon.		
	IL=interleukin.		
	TNF=tumor necrosis factor.		

Table 3. Recently discovered biomarkers

K2P5.1	Potassium channel affecting T cells		
CCL2	Chemokine		
CXCL10	Chemokine		
IL-8	Chemokine		
Complement C4 fragment	Marker of complement activation		
MMP-8	Affects inflammatory cell migration through tissue		
CD56bright NK cells	Regulate activated T cells		
25-Hydroxy-vitamin D	Has immunomodulatory effects		
Urinary neopterin	Marker of macrophage activity		
ILT3	Downregulates immune activity		
Myoinositol	MR marker of tissue integrity		
Bri2-23	Neuronal protein		
Fetuin-A	Immunoregulatory protease inhibitor		
Pentosidine	Marker of tissue damage and inflammation		
Leptin receptor	Metabolic and immune pathways		
IL=interleukin.			
ILT=immunoglobulin-like transcript.			
MMP=matrix metalloprotease			

In MS patients, these biomarkers contribute to the pathophysiological mechanism of the disease[12]. Hitherto, inflammatory and stress oxidative pointers are expected to play a more significant role in the relapse of the disease. On the other hand, inactivity of the glia, the restoration of the myelin and axon damage point to progressive and therefore, destructive syndromes. These biomarkers can otherwise differentiate between an MS patient with moderate symptoms or severe symptoms (Table 2).

Treatment Response Biomarkers

Treatment response biomarkers evaluate how effective the treatment for the MS patient has been. They also determine the imminence of treatment failure, making it possible to move on to alternative treatment options beforehand. Kinetic and dynamic medical biomarkers also belong to this class of biomarkers, helping the regulation of the patient's dosage. Another supposed use of these biomarkers is the appraisal of the patient's response to different personalized medical treatment. With over ten treatment options available for MS, it is crucial that we find the proper blood-based selection to avoid unnecessary side-effects.

Side-Effect Scrutiny Biomarkers

These biomarkers are advantageous for patients prone to

progressive, pathological or inflammatory diseases. For instance, the antibodies for human polyomavirus II, formerly known as John Cunningham virus, is a biomarker convenient for allocating the danger of progressive multifocal leukoencephalopathy (PML) in patients treated with Natalizumab[13, 14].

Advantages and Disadvantages of Biomarker Assessment

MS biomarkers are found in different bodily fluids such as urine, blood, cerebrospinal fluid (CSF) and tears. There are advantages and disadvantages to each sample. MS waste products are seldom inspected, making CSF a fitting choice for pathology. There are different approaches to CSF sampling; I. e. markers soluble in both solutions, like cell aggregations. However, CSF sampling is an invasive procedure and only applicable for a limited number of times.

Blood sampling is a fairly simple procedure. However, there is an abundance of biomarkers in the blood which can be a result of different systematic infections in the body, heme catabolism and kidney waste.

Urine is also another non-invasive sampling procedure. The issue with this method is the chronic infections of the urethra and bacterial colonizations of the bladder, common in most disabled MS patients, which manipulates the results. Plus, some patients with bladder dysfunction use artificial hypo-hydration for ejection, which restricts examination [15, 16].

Immune Change Biomarker Feedback Cytokines and Their Receptors

Cytokines are intensely examined in MS. As a result of inflammation in MS waste, there is an increase in different types of cytokines in MS patients. These changes are not exclusive to MS patients and can be detected in other inflammatory CNS disorders. TH1 inflammatory cytokines such as INF Gamma, tumor necrosis factor alpha (TNF alpha) and interleukin 12 increase in MS relapse phases [16], while anti-inflammatory cytokines like TH2, IL-4, IL-10 and TGF Beta correlate with clinical recovery and RRMS patients [17]. TH17 is a proinflammatory cytokine which stands as the gnomon for the IL-17 cell line, able to strongly affect a wide spectrum of cells, consequently releasing a couple types of inflammatory mediators, the CXL10 and CXL1. GN-CSF chemokines, the IL-8 and IL-6[18]. It is reported that there is a connection between MS patients and the level of active plaques. The IL-17 inflammatory cells produce a range of IFN- λ , which makes them pathogens for MS and EAE [19, 20]. In fact, the IFN- λ produced by the TH1 cells, increase the interference of leukocyte adhesion molecules with the endothelial cells of the blood-brain barrier, making this region more impenetrable [21]. This cytokine also increases the activity of MHCII and MHCI molecules on the surface of astrocytes and oligodendrocytes, enhancing the reactivity of these cells with immunized cells, probably causing cell death [22]. Furthermore, macrophages and microglia cooperate in the destruction of myelin in MS and EAE. We can basically reaffirm that the IFN- λ kills oligodendrocytes using apoptosis or necrosis (depending on the cell's maturity) [23]. Hence, most MS treatments focus on the suppression of inflammatory responses; I.E. IFN- λ applied for the inhibition of MHCII for its effects on antigen cell surface and the reduction of T cells, via stopping the production of IL-2 [24]. It has been conducted that MS treatments owe part of their competence to the ability of taming the TH17 cells [25, 26]. Sweeney et al. have concluded that the prescription of Interferon Beta for EAE reduces IL-17 while increasing IL-27 [27].

IL-10 operates as a regulatory cytokine, especially valuable in the limitation of inflammatory responses and tissue damage and adjusts the immune system. Studies have shown that CD4-T cells produce IL-10, making them quite profitable in MS treatment [28, 29].

Tumor Necrosis Factor Alpha (TNF α) is another essential cytokine is modifying the immune system for the better[30]. This cytokine is built by TH1, inflammatory macrophages, natural killing cells, mast cells, eosinophils and neurons themselves [31]. The TNF α counts as a pathogen for EAE and MS [32, 33]. This cytokine breaks the blood-brain barrier, activates the microglia cells and inducts apotosis cells in MS and EAE [32]. There is a direct correlation between TNF α levels and severity of MS symptoms [33, 34]. Ozensy et al. have demonstrated that there is a higher number of cells producing TNF α in MS patients than the average individual. In another research work by Rentzos et al. in 1996, it has been established the TNF α witnesses higher levels in the cerebrospinal fluid in MS patients.

Chemokines and Their Receptors

Chemokines are chemotactic cytokines which summon immune cells to the lymphoid organs and inflammation points. The CCR5 chemokine receptor relates to the relapse of the disease accompanied by symptoms [35, 36]; also CXCR3 increases in T lymphocytes during relapse as the immunohistochemistry autopsy report of some brain parts contains active MS waste. This addresses the existence of CXCR3 in all impenetrable T lymphocytes [35, 37].

In brain samples, the CXCR3+ is abnormal. It has been expressed that the retention of T cells and CXCR3+ in MS patients leads to the appearance of its ligands (IP-10) [38]. In the absence of ligands, the CXCR3 cells go back to their normal cycle. Active inflammatory MS waste containing CCR5+/CCR1 is evidence for hematogen monocytes abundantly found in perivascular cell spaces (crowding point for varied leucocytes) and demyelination borders despite their omission in non-inflammatory brain cuts [39]. Chemokines and their receptors might be important in the heterogynous study of the disease, but more research is in order for certainty.

Oligoclonal Bands and Antibodies

Recently, the only biomarker accepted apart from MRI in MS diagnosis are called CSF oligoclonal bands (OCB), which occur as isolated immunoglobulins via isoelectric focusing in 90-95% of MS patients [40]. The persistent and remitting presence of OCB in MS patients is a telltale sign of an immune response to B lymphocytes in the spine [41]. The infusion of the oligoclonal IEF immunoglobulin G (IgG) and immune recognition (antibody) of alkali phosphatase is a more sensitive and specific approach to MS symptoms [42]. This method enables the prediction of a second attack in patients with isolated demyelination syndrome and the plausibility of transformation into Clinically Definite Multiple Sclerosis (CDMS). The oligoclonal IgM is similarly detectable in IEF [43]. IgM is the strongest activator for complementation in demyelination areas in MS and NMO patients [44, 45]. Hitherto, the first reports of IgM do not stand as a strong prophecy on early conversion into CDMS and its invasion period [46-51].

The role of auto-antibody pathology in autoimmune disorders is widely affirmed. Specifically, we highlight MBL and glycoprotein oligodendrocyte (MOG) out of the plausible MS autoantigens responsible for experimental autoimmune encephalomyelitis (EAE) because of their positioning in the dense MBP myelin and the outer surface of the myelin sheath of MOG. Recently, it has been suggested that the MBP-exclusive antibodies and IgM antibodies exclusive to the outer cell domain of MOG present in CIS patients' serums are extremely inclined to turn into CDMS [52]. AntiMOG antibodies are more prevalent in MS patients than the healthy individual. Akin to the axon damage perceived in rats after injection, this antibody causes a boost in demyelination and cytotoxicity in the laboratory environment. It is not yet known whether this peculiar production of antibodies in MS patients is feedback to myelin damage or the cause of it [52-54]. More research is needed to actuate the correct relationship between these antibodies and pathological or clinical parameters [55, 56].

Lately, it has become apparent that the Epstein-Barr virus (EBV) is a possible environmental factor contributing to the development of MS [57]. Although the host range specificity of OCB is not yet fully understood and a clear link between OCB and MS' pathogenesis, EBV seems to be a target of OCB [58]. In an effort to discover OCB's specificity range, the layout of 37,000 protein expressions were surveyed. Out of these, two highly amenable epitopes composed of EBV proteins were discovered. In blood samples collected before the manifestation of MS, there were higher aggregations of antibodies for EBNA and EBNA1 complexes than that of healthy people and these aggregations maintained their high numbers after MS [59]. There is generally a higher risk for MS with antibody aggregations. Teenagers and young adults with a history of infectious mononucleosis (a fever-inducing illness caused by EBV) are more exposed to MS than individuals who contract this illness in later years of their lives [60]. Apart from EBV [61, 62], HLA-DR15 which is the main genetic component of MS also induce EBNA1 antibody aggregations [63].

Nueromyelitis Optica (NMO) syndrome, is a heterogeneous MS subgroup consisting of demyelination and inflammation in the spinal and optical nerves. Based on serologic and clinical evidence, it has been concluded that antibody autoimmune responses bear greater numbers in NMO patients [64, 65]. Confirmed immunomodulatory MS ttreatments are ineffective for NMO patients, making treatment plans for these patients tremendously exclusive. Precise primary diagnosis and a strong invasive interference are crucial in order to avoid an NMO relapse, as this increases mortality rates and disability in MS patients [45, 66].

Aquaporin-4 has been identified as the target antigen of NMO-IgG. Aquaporin-4 acts in the elemental astrocytic processes of the blood-brain barrier and is also the main executioner of the brain's hemostasis [67, 68]. NMO is found in MS waste in various studies done on different patient populations [69, 70]. NMO-IgG antibodies are the most-recently discovered exclusive biomarkers for MS, first distinguished while investigating demyelination diseases affecting CNS. They greatly aid in differentiating between NMO and MS patients [54, 71].

The YKL-40 biomarker is a secreted glycoprotein, also known as chitinase-3-like protein (CHI3L1), is a gliaactivation marker mainly produced via astrocyte reaction [72, 73]. It can be alternatively be produced by activated macrophages, vascular cells, epithelial airways and chondrocytes [73]. The serum volume of YKL-14 increases with the occurrence inflammatory conditions such as rheumatoid arThe CXL13 biomarker is a B lymphocyte chemoattractant, making adjustments in secondary lymphoid organs in both homeostatic and inflammatory contingencies [75]. They are produced by B cell follicles and assist in the formation of the germinal center. Its receptor is CXCR5, since it uses active B cells and CD4 T cell follicles in its job [76-78].

Many studies have shown that the expression of CXCL13 witnesses an increase in CIS, RRMS, SPMS and PPMS patients in comparison to the healthy control group and other neurological patients [79]. This increase is often accompanied by the relapse of EDSS and nerve damage [80, 81]. The CXCL13 is likewise more ubiquitous in RRMS patients than other MS patients [81].

MicroRNAs

A microRNA is a noncoding RNA molecule containing no more than 22 nucleotides serving in RNA silencing, posttranscriptional gene expression regulation and protein translation [82, 83]. MicroRNAs play an imperative part in many biological processes such as metabolism, apoptosis and angiogenesis[84, 85]. Almost 1/3 of all human genes utilize the indirect encoding of these molecules [86, 87].

MicroRNA profiles have been investigated at length in MS patients in many samples such as the peripheral blood mononuclear cell (PBMC) [88, 89], blood samples [90] and brain damage [91]; all of which have presented the same result: the microRNA profile is correlated with Multiple Sclerosis [89].

A research work was conducted, calculating microRNAs expression rates like miR-21, miR-14ba, miR-146a, miR-146b, miR-150 and miR155 in MS patients as opposed to that of the healthy control group. The conclusion was that RRMS patients had more notable expression rates of miR-21, miR146a and miR146b [88].

Another study calibrated the expression rate of miR-326 in PBMC and deduced that the rates correlate with the severity of MS and the symptoms of rats suffering from experimental autoimmune encephalomyelitis (EAE) [6].

In the laboratory, containing the miR-126 resulted in lower numbers of TH17 [92].

The microRNA can be present in many bodily fluids including plasma, serum, urine and saliva [93] and is scrutinized as a possible biomarker in many diseases [94, 95].

A recent study conducted by the Gandhi group has assessed the miRNA-320a to be the most significant altering microRNA in MS patients' serum. They have also found hsa-miR-27a-3p to be the most significant microRNA connected with MS relapses. The miRNA-19 correlates with EDSS, with miR-199-5p correlating with associated disabilities [96].

KIR4.1 Antibodies

A detailed study examining various antibody responses in patients with MS [97] identified IgG1 and IgG3 antibod-

ies that bind to glial cells in the brain tissues. This gliaspecific immunoreactivity was not found in the sera of patients who had other neurological disorders. KIR4.1, a potassium channel, was the molecular target of these antibodies. This channels helps in maintaining potassium and balancing water levels in the body. Antibodies moving against KIR4.1 were detected in the serum of 186 out of 397 patients with MS (about 46%), whereas they were present in the sera of less than 1% of patients with other neurological disorders (n=329) making it insignificant, and not observed at all in healthy individuals. This study determined that KIR4.1 is the target of an autoantibody response in some patients with MS [97]. KIR4.1 expressed on astrocytes localizes with another channel, the aquaporin-4 (AQP4) which was previously argued to be the target for NMO. NMO-IgGis present in 73% of patients with NMO, but absent in the serum of patients with MS [64]

TOB1

Studies have suggested that the polymorphism of TOB1 is an individual supporting the development of CIS into a clinical diagnosis of MS. The TOB1 gene contains the multiplication of T cells. This gene keeps inclined cells in a non-reactive state and its decrease leads to a stringent immune response [98].

Osteopontin

OPN, also known as bone sialoprotein I is a protein encoded by the SPP1 gene, expressed from a crowding of cells and tissues. OPN operates in a range of physiopathological duties, as its seen in the nucleus and cytoplasm of many different types of cells [99, 100].

OPN plays a paramount role in nonbone affairs; i.e. monocyte adhesion, phagocytosis and cell migration [101, 102]. It also polarizes TH cells into TH1 and TH2, modifying cytokine expression [103].

OPN engages in the advancement of many autoimmune diseases such as MS [104-107], rheumatoid arthritis [108, 109], psoriasis [110] and fungus infections [111]. In MS patients, OPN expression accompanies MS damage [112, 113]. Many studies have also shown that OPN also correlates with severity and relapse of PPMS

Microbiota-Related Lipopeptides

Gut microbiota intervenes with many immunological diseases including MS, inflammatory bowel disease (IBD), type I diabetes and rheumatoid arthritis [114-116].

When a potent 16s ribosomal RNA was studied with the purpose of analyzing gut microbiota's association with MS (n=16) and healthy control groups (n=44), how the RNAs change in the presence of MS was revealed. Methanobrevibacter archaea and Akkermansia genera increase and the Butyricimonas genera decrease. These changes in the gut interfere with the expression of other genes involved in dendritic cell maturity, interferon signaling and NF. Patients in treatment present signs of an increase in Prevotella and Sutterella genera and a decrease in Sarcina genera as opposed to

patients not being treated. However, in order to confidently comment on how gut microbiota acts in MS pathogenesis, more studies must be in order [117].

Complement Biomarkers Profile

There is no shortage of studies on how the complement system could aid in the pathogenesis of MS and therefore, how various proteins can carry out as an MS biomarker. Their conclusions, however, contradict one another. There is a proteomic analysis conducted, scrutinizing complementexclusive proteins (I Factor, C3 and Clusterin) and their inconstancies, usually occurring in the CSF samples of MS patients. Lately, the H factor, a complement system regulator, is considered a serum biomarker for detection of the disease's activities. The H factor's levels are meaningful in progressive diseases and patients with a higher rate of relapse [118].

Adhesion Molecules as Blood-Barrier Disorder Biomarkers

These adhesion molecules exploring the channels for endothelial cells regulating leukocyte transmigration are pathogens for MS [119, 120]. Adhesion molecules can be found in the form of activated endothelial cell solution, coenocytes, platelets, serum and CSF [121]. It has been determined that the plasmatic levels of soluble adhesion molecules (sP-Selectin, sPECAM-1, sE-Selectin) have increased in RRMS patients [122], in comparison to chronic progressive patients. Plus, this increase in relapse signifies these molecules as clinical markers for the activity of MS. The escalation of these intercellular molecules' levels (sICAM-1) has been perceived in MS patients during relapse. A correlation between sICAM-1 in the cerebrovascular fluid and igG traces and RRMS patients has been addressed [123]. A surge in circulating vascular cell adhesion molecule-1 (SVCAM-1) leads to a decrease in MS waste in patients in treatment, alternatively linked with IFN- β -1a[124]. The first regulation of sVCAM-1 happening in the first one to six months of SPMS clinically interferes with patients being treated with IFN-β-1b in months 19-24. The same thing happened in IFN- β -1a treatments [125, 126].

VLA-4 is more sensitive to predicting treatment results, as compared with the boost in VCAM-1 and overall increases in each relapse period [126, 127].

MMP9 enzymes work in the destruction of the outer cell matrix and the proteolysis of myelin. The MMP9 enzyme around the arteries breaks collagen and opens the bloodbrain barrier. This breakage causes inflammatory cells to spill into the white section of CNS. When T cells are distributed in the brain's white matter, an immune response happens to almost every molecule in the myelin structure. Migrating B cells are discharged against proteins and lipids present in the myelin shell. T cells usually mark regularlyoccurring proteins in the myelin, like MOG, MBP and PLP[24]. T cells also secret TNF- α and LT- α cytokines. These cytokines produce nitric oxide and Osteopontin, employing the macrophages, microglia cells and astrocytes Biomarker efficacy in multiple sclerosis diagnosis and treatment

7

[96].

The NO free radical is an important mediator in autoimmune diseases. NO kills the oligodendrocytes via the microglia. iNOS which catalyzes NO production is found in demyelinated areas of the body. TNF- α and IFN- γ encode iNOS in the astrocytes, microglia and macrophages. The final result of antibodies, the complement system, NO and TNF- α is an intensive damage to the myelin and further stimulation of the macrophages moving to the phagocytosis of larger parts of the myelin shell. Osteopontin output by macrophages and T cells regulate the increasing of T-helper-1 cytokines such as IFN- γ and IL-12 and regulate the reduction of IL-10, a Thelper-2. T-helper-1 cytokines worsen MS symptoms, while T-helper-2 cytokines probably stop platelets from growing [128, 129].

Karabudak et al. conducted a study in 2004, investigating the effects of IFN- β on MMP-9 and TIMP-1 in RRMS patients in a one year period. MMP-9 levels did not seem to present anything meaningful, but TIMP-1 rates increased in the course of the disease in a significant manner [130].

Garcia Montojo et al. studied 50 MS patients in 2010 who were being treated with INF-beta within a two-year period. MMP9 and TIMP1 levels almost doubled in this time. It is thus believed that these two are highly inhabitable for INF- β [131].

Liuzzi et al. discerned a reverse meaningful relation between MMP-9 and its containment endogens, TIMP-1 in RRMS. This finding confirms that the increase of MMP-9 and decrease of TIMP-1 in sera levels of MS patients participate in BBB destruction and CNS invasion by lymphocyte T [132].

CONCLUSION

Biomarkers present a wide array of biological information concerning MS symptoms. Their analysis acts as a guiding light when looking for the right treatment and investigating its efficiency, in case the researcher is willing to closely examine it before, during and after treatment. We hope molecular techniques result in more specific and sensitive biomarkers. It is certain that with the engagement of biomarkers signifying the disease's course and activity, we are able to prescribe more peculiar medical treatment and assess the patient's response adequately.

CONFLICTS OF INTEREST

The author(s) declare(s) that there is no conflict of interest regarding the publication of this article.

REFERENCES

- 1. Das Sarma J. A mechanism of virus-induced demyelination. Interdiscipl perspect Infect Dis 2010;2010.
- Kakalacheva K, Münz C, Lünemann JD. Viral triggers of multiple sclerosis. Biochim Biophysic Acta (BBA)-Mol Bas Dis 2011;1812(2):132-40.
- 3. McFarlin DE, McFarland HF. Multiple sclerosis. New Eng J Med 1982;307(20):1246-51.
- Hemmer B, Archelos JJ, Hartung H-P. New concepts in the immunopathogenesis of multiple sclerosis. Nature Rev Neurosci

2002;3(4):291.

- Haghjooy SJ, Saadatnia M, Homayouni V, Nikoogoftar M, Maghzi A, Etemadifar M, et al. Interferon-beta-1b protects against multiple sclerosis-induced endothelial cells apoptosis. Front Biosci (Elite edition). 2012;4:1368-74.
- Zahednasab H, Balood M. The role of miR-326 and miR-26a in MS disease activity. Gene 2014;548(1):158.
- Villoslada P. Biomarkers for multiple sclerosis. Drug News Perspect 2010;23(9):585-95.
- Bielekova B, Howard T, Packer AN, Richert N, Blevins G, Ohayon J, et al. Effect of anti-CD25 antibody daclizumab in the inhibition of inflammation and stabilization of disease progression in multiple sclerosis. Arch Neurol 2009;66(4):483-9.
- Beilekova B. Regulatory CD56[<] bright> natural killer cells mediate immunomodulatory effects of IL2Rα-targeted therapy (daclizumab) in multiple sclerosis. Proc Natl Acad Sci USA 2006;103:5941-6.
- Miller D, Weinshenker BG, Filippi M, Banwell B, Cohen J, Freedman M, et al. Differential diagnosis of suspected multiple sclerosis: a consensus approach. Multiple Scleros J 2008;14(9):1157-74.
- Nielsen JM, Korteweg T, Barkhof F, Uitdehaag BM, Polman CH. Overdiagnosis of multiple sclerosis and magnetic resonance imaging criteria. Ann Neurol 2005;58(5):781-3.
- Lassmann H, Brück W, Lucchinetti CF. The immunopathology of multiple sclerosis: an overview. Brain Pathol 2007;17(2):210-8.
- Antoniol C, Stankoff B. Immunological markers for PML prediction in MS patients treated with natalizumab. Front Immunol 2015;5:668.
- Antoniol C, Stankoff B. Immunological Markers for PML Prediction in MS Patients Treated with Natalizumab. Front Immunol 2014;5:668.
- Sospedra M, Martin R. Immunology of multiple sclerosis. Ann Rev Immunol 2005;23:683-747.
- Link H. The cytokine storm in multiple sclerosis. Multiple Scleros J 1998;4(1):12-5.
- Balashov KE, Smith DR, Khoury SJ, Hafler DA, Weiner HL. Increased interleukin 12 production in progressive multiple sclerosis: induction by activated CD4+ T cells via CD40 ligand. Proceed Natl Acad Sci 1997;94(2):599-603.
- Korn T, Oukka M, Kuchroo V, Bettelli E, editors. Th17 cells: effector T cells with inflammatory properties. Seminars in immunology; 2007: Elsevier.
- 19. Balasa R. T helper 17 cells in multiple sclerosis and experimental autoimmune encephalomyelitis. Roman J Neurol. 2010;9(4).
- Jadidi-Niaragh F, Mirshafiey A. Th17 cell, the new player of neuroinflammatory process in multiple sclerosis. Scand J Immunol 2011;74(1):1-13.
- 21. Cayrol R, Wosik K, Berard JL, Dodelet-Devillers A, Ifergan I, Kebir H, et al. Activated leukocyte cell adhesion molecule promotes leukocyte trafficking into the central nervous system. Nature Immunol 2008;9(2):137.
- Rodriguez M. Effectors of demyelination and remyelination in the CNS: implications for multiple sclerosis. Brain Pathol 2007;17(2):219-29.
- Balabanov R, Strand K, Kemper A, Lee JY, Popko B. Suppressor of cytokine signaling 1 expression protects oligodendrocytes from the deleterious effects of interferon-γ. J Neurosci 2006;26(19):5143-52.
- Javaid MA, Abdallah M-N, Ahmed AS, Sheikh Z. Matrix metalloproteinases and their pathological upregulation in multiple sclerosis: an overview. Acta Neurol Belgic 2013;113(4):381-90.
- Durelli L, Conti L, Clerico M, Boselli D, Contessa G, Ripellino P, et al. T-helper 17 cells expand in multiple sclerosis and are inhibited by interferon-β. Ann Neurol 2009;65(5):499-509.
- Ramgolam VS, Sha Y, Jin J, Zhang X, Markovic-Plese S. IFN-β inhibits human Th17 cell differentiation. J Immunol 2009;183(8):5418-27.
- 27. Sweeney CM, Lonergan R, Basdeo SA, Kinsella K, Dungan LS, Higgins SC, et al. IL-27 mediates the response to IFN- β therapy in multiple sclerosis patients by inhibiting Th17 cells. Brain Behav Immun 2011;25(6):1170-81.
- Boonstra A, Barrat FJ, Crain C, Heath VL, Savelkoul HF, O'Garra A. 1α, 25-Dihydroxyvitamin D3 has a direct effect on naive CD4+ T cells to enhance the development of Th2 cells. J Immunol 2001;167(9):4974-80
- 29. Perrella O, Sbreglia C, Perrella M, Spetrini G, Gorga F, Pezzella M, et

This paper is available online at: http://ijpt.iums.ac.ir

al. Interleukin-10 and tumor necrosis factor-alpha: model of

- immunomodulation in multiple sclerosis. Neurol Res 2006;28(2):193-5. 30. Locksley RM, Killeen N, Lenardo MJ. The TNF and TNF receptor
- superfamilies: integrating mammalian biology. Cell 2001;104(4):487-501.
- 31. Kant S, Swat W, Zhang S, Zhang Z-Y, Neel BG, Flavell RA, et al. TNF-stimulated MAP kinase activation mediated by a Rho family GTPase signaling pathway. Genes Develop 2011;25(19):2069-78.
- 32. Li R, Rezk A, Healy LM, Muirhead G, Prat A, Gommerman JL, et al. Cytokine-defined B cell responses as therapeutic targets in multiple sclerosis. Front Immunol 2016;6:626.
- Patejdl R, Penner IK, Noack TK, Zettl UK. Multiple sclerosis and fatigue: a review on the contribution of inflammation and immunemediated neurodegeneration. Autoimmun Rev 2016;15(3):210-20.
- 34. Sharief MK, Hentges R. Association between tumor necrosis factor-α and disease progression in patients with multiple sclerosis. New Eng J Med 1991;325(7):467-72.
- 35. Sørensen TL, Tani M, Jensen J, Pierce V, Lucchinetti C, Folcik VA, et al. Expression of specific chemokines and chemokine receptors in the central nervous system of multiple sclerosis patients. J Clin Invest 1999;103(6):807-15.
- 36. Strunk T, Bubel S, Mascher B, Schlenke P, Kirchner H, Wandinger KP. Increased numbers of CCR5+ interferon- γ -and tumor necrosis factor- α secreting T lymphocytes in multiple sclerosis patients. Ann Neurol 2000;47(2):269-73.
- 37. Balashov KE, Rottman JB, Weiner HL, Hancock WW. CCR5+ and CXCR3+ T cells are increased in multiple sclerosis and their ligands MIP-1α and IP-10 are expressed in demyelinating brain lesions. Proceed Natl Acad Sci 1999;96(12):6873-8.
- 38. Sørensen TL, Trebst C, Kivisäkk P, Klaege KL, Majmudar A, Ravid R, et al. Multiple sclerosis: a study of CXCL10 and CXCR3 co-localization in the inflamed central nervous system. J Neuroimmunol 2002;127(1-2):59-68.
- Trebst C, Sørensen TL, Kivisäkk P, Cathcart MK, Hesselgesser J, Horuk R, et al. CCR1+/CCR5+ mononuclear phagocytes accumulate in the central nervous system of patients with multiple sclerosis. Am J Pathol 2001;159(5):1701-10.
- 40. Kabat EA, Glusman M, Knaub V. Quantitative estimation of the albumin and gamma globulin in normal and pathologic cerebrospinal fluid by immunochemical methods. Am J Med 1948;4(5):653-62.
- Bourahoui A, De Seze J, Guttierez R, Onraed B, Hennache B, Ferriby D, et al. CSF isoelectrofocusing in a large cohort of MS and other neurological diseases. Eur J Neurol 2004;11(8):525-9.
- 42. Masjuan J, Alvarez-Cermeno J, Garcia-Barragan N, Díaz-Sánchez M, Espiño M, Sadaba M, et al. Clinically isolated syndromes: a new oligoclonal band test accurately predicts conversion to MS. Neurology 2006;66(4):576-8.
- 43. Sharief MK, Keir G, Thompson EJ. Intrathecal synthesis of IgM in neurological diseases: a comparison between detection of oligoclonal bands and quantitative estimation. J Neurol Sci 1990;96(2-3):131-42.
- 44. Lucchinetti C, Brück W, Parisi J, Scheithauer B, Rodriguez M, Lassmann H. Heterogeneity of multiple sclerosis lesions: implications for the pathogenesis of demyelination. Ann Neurol 2000;47(6):707-17.
- 45. Lucchinetti CF, Mandler RN, McGavern D, Bruck W, Gleich G, Ransohoff RM, et al. A role for humoral mechanisms in the pathogenesis of Devic's neuromyelitis optica. Brain 2002;125(7):1450-61.
- 46. Schneider R, Euler B, Rauer S. Intrathecal IgM-synthesis does not correlate with the risk of relapse in patients with a primary demyelinating event. Eur J Neurol 2007;14(8):907-11.
- 47. Villar LM, Masjuan J, González-Porqué P, Plaza J, Sádaba MC, Roldán E, et al. Intrathecal IgM synthesis is a prognostic factor in multiple sclerosis. Ann Neurol 2003;53(2):222-6.
- Villar LM, Sádaba MC, Roldán E, Masjuan J, González-Porqué P, Villarrubia N, et al. Intrathecal synthesis of oligoclonal IgM against myelin lipids predicts an aggressive disease course in MS. J Clin Invest 2005;115(1):187-94.
- Villar L, Garcia-Barragan N, Espino M, Roldán E, Sádaba M, Gómez-Rial J, et al. Influence of oligoclonal IgM specificity in multiple sclerosis disease course. Multiple Scleros J 2008;14(2):183-7.
- Perini P, Ranzato F, Calabrese M, Battistin L, Gallo P. Intrathecal IgM production at clinical onset correlates with a more severe disease course

- in multiple sclerosis. J Neurol Neurosurg Psychiatry 2006;77(8):953-5.
- 51. Koch M, Heersema D, Mostert J, Teelken A, Keyser JD. Cerebrospinal fluid oligoclonal bands and progression of disability in multiple sclerosis. Eur J Neurol 2007;14(7):797-800.
- 52. Berger T, Rubner P, Schautzer F, Egg R, Ulmer H, Mayringer I, et al. Antimyelin antibodies as a predictor of clinically definite multiple sclerosis after a first demyelinating event. New Eng J Med 2003;349(2):139-45.
- 53. Tomassini V, De Giglio L, Reindl M, Russo P, Pestalozza I, Pantano P, et al. Anti-myelin antibodies predict the clinical outcome after a first episode suggestive of MS. Multiple Scleros J 2007;13(9):1086-94.
- 54. Greeve I, Sellner J, Lauterburg T, Walker U, Rösler KM, Mattle H. Anti-myelin antibodies in clinically isolated syndrome indicate the risk of multiple sclerosis in a Swiss cohort. Acta Neurol Scand 2007;116(4):207-10.
- 55. Kuhle J, Lindberg RL, Regeniter A, Mehling M, Hoffmann F, Reindl M, et al. Antimyelin antibodies in clinically isolated syndromes correlate with inflammation in MRI and CSF. J Neurol 2007;254(2):160-8.
- 56. Rauer S, Euler B, Reindl M, Berger T. Antimyelin antibodies and the risk of relapse in patients with a primary demyelinating event. J Neurol Neurosurg Psychiatry 2006;77(6):739-42.
- Giovannoni G, Cutter GR, Lunemann J, Martin R, Münz C, Sriram S, et al. Infectious causes of multiple sclerosis. Lancet Neurol 2006;5(10):887-94.
- 58. Sundström P, Juto P, Wadell G, Hallmans G, Svenningsson A, Nyström L, et al. An altered immune response to Epstein-Barr virus in multiple sclerosis: a prospective study. Neurology 2004;62(12):2277-82.
- DeLorenze GN, Munger KL, Lennette ET, Orentreich N, Vogelman JH, Ascherio A. Epstein-Barr virus and multiple sclerosis: evidence of association from a prospective study with long-term follow-up. Arch Neurol 2006;63(6):839-44.
- 60. Levin LI, Munger KL, Rubertone MV, Peck CA, Lennette ET, Spiegelman D, et al. Temporal relationship between elevation of Epstein-Barr virus antibody titers and initial onset of neurological symptoms in multiple sclerosis. JAMA 2005;293(20):2496-500.
- Thacker EL, Mirzaei F, Ascherio A. Infectious mononucleosis and risk for multiple sclerosis: a meta-analysis. Ann Neurol 2006;59(3):499-503.
- Hernán MA, Zhang SM, Lipworth L, Olek MJ, Ascherio A. Multiple sclerosis and age at infection with common viruses. Epidemiology 2001;12(3):301-6.
- 63. De Jager P, Simon K, Munger K, Rioux J, Hafler D, Ascherio A. Integrating risk factors: HLA-DRB1* 1501 and Epstein–Barr virus in multiple sclerosis. Neurology 2008;70(13 Part 2):1113-8.
- Lennon VA, Kryzer TJ, Pittock SJ, Verkman A, Hinson SR. IgG marker of optic-spinal multiple sclerosis binds to the aquaporin-4 water channel. J Experim Med 2005;202(4):473-7.
- Lennon VA, Wingerchuk DM, Kryzer TJ, Pittock SJ, Lucchinetti CF, Fujihara K, et al. A serum autoantibody marker of neuromyelitis optica: distinction from multiple sclerosis. Lancet 2004;364(9451):2106-12.
- Weinshenker BG, Wingerchuk DM, Vukusic S, Linbo L, Pittock SJ, Lucchinetti CF, et al. Neuromyelitis optica IgG predicts relapse after longitudinally extensive transverse myelitis. Ann Neurol 2006;59(3):566-9.
- 67. Misu T, Fujihara K, Kakita A, Konno H, Nakamura M, Watanabe S, et al. Loss of aquaporin 4 in lesions of neuromyelitis optica: distinction from multiple sclerosis. Brain 2007;130(5):1224-34.
- Roemer SF, Parisi JE, Lennon VA, Benarroch EE, Lassmann H, Bruck W, et al. Pattern-specific loss of aquaporin-4 immunoreactivity distinguishes neuromyelitis optica from multiple sclerosis. Brain 2007;130(5):1194-205.
- Jarius S, Franciotta D, Bergamaschi R, Wright H, Littleton E, Palace J, et al. NMO-IgG in the diagnosis of neuromyelitis optica. Neurology 2007;68(13):1076-7.
- 70. Takahashi T, Fujihara K, Nakashima I, Misu T, Miyazawa I, Nakamura M, et al. Anti-aquaporin-4 antibody is involved in the pathogenesis of NMO: a study on antibody titre. Brain 2007;130(5):1235-43.
- Keegan M, Pineda A, McClelland R, Darby C, Rodriguez M, Weinshenker BG. Plasma exchange for severe attacks of CNS demyelination: predictors of response. Neurology 2002;58(1):143-6.
- Bonneh-Barkay D, Wang G, Starkey A, Hamilton RL, Wiley CA. In vivo CHI3L1 (YKL-40) expression in astrocytes in acute and chronic

neurological diseases. J Neuroinflam 2010;7(1):34.

- Canto E, Tintoré M, Villar LM, Costa C, Nurtdinov R, Álvarez-Cermeño JC, et al. Chitinase 3-like 1: prognostic biomarker in clinically isolated syndromes. Brain 2015;138(4):918-31.
- Peltomaa R, Paimela L, Harvey S, Helve T, Leirisalo-Repo M. Increased level of YKL-40 in sera from patients with early rheumatoid arthritis: a new marker for disease activity. Rheumatol Int 2001;20(5):192-6.
- Cyster JG. Chemokines, sphingosine-1-phosphate, and cell migration in secondary lymphoid organs. Ann Rev Immunol 2005;23:127-59.
- 76. Zotos D, Coquet JM, Zhang Y, Light A, D'Costa K, Kallies A, et al. IL-21 regulates germinal center B cell differentiation and proliferation through a B cell–intrinsic mechanism. J Experim Med 2010;207(2):365-78.
- 77. Crotty S. The 1-1-1 fallacy. Immunol Rev 2012;247(1):133-42.
- Victora GD, Nussenzweig MC. Germinal centers. Annual review of immunology. 2012;30:429-57.
- 79. Stilund M, Gjelstrup MC, Petersen T, Møller HJ, Rasmussen PV, Christensen T. Biomarkers of inflammation and axonal degeneration/damage in patients with newly diagnosed multiple sclerosis: contributions of the soluble CD163 CSF/serum ratio to a biomarker panel. PloS One 2015;10(4):e0119681.
- Festa ED, Hankiewicz K, Kim S, Skurnick J, Wolansky LJ, Cook SD, et al. Serum levels of CXCL13 are elevated in active multiple sclerosis. Multiple Scleros J 2009;15(11):1271-9.
- Khademi M, Kockum I, Andersson ML, Iacobaeus E, Brundin L, Sellebjerg F, et al. Cerebrospinal fluid CXCL13 in multiple sclerosis: a suggestive prognostic marker for the disease course. Multiple Scleros J 2011;17(3):335-43.
- Ota K, Matsui M, Milford EL, Mackin GA, Weiner HL, Hafler DA. Tcell recognition of an immuno-dominant myelin basic protein epitope in multiple sclerosis. Nature 1990;346(6280):183.
- Pette M, Fujita K, Kitze B, Whitaker J, Albert E, Kappos L, et al. Myelin basic protein-specific T lymphocyte lines from MS patients and healthy individuals. Neurology 1990;40(11):1770-.
- 84. Liu B, Yang X, Liang X, Wang L, Shao M, Han W, et al. Expressions of MiR-132 in patients with chronic hepatitis B, posthepatitic cirrhosis and hepatitis B virus-related hepatocellular carcinoma. Eur Rev Med Pharmacol Sci 2018;22(23):8431-7.
- Qu Z, Li W, Fu B. MicroRNAs in autoimmune diseases. BioMed Res Int 2014;2014.
- Lewis BP, Shih IH, Jones-Rhoades MW, Bartel DP, Burge CB. Prediction of mammalian microRNA targets. Cell 2003;115(7):787-98.
- 87. Martinelli-Boneschi F, Fenoglio C, Brambilla P, Sorosina M, Giacalone G, Esposito F, et al. MicroRNA and mRNA expression profile screening in multiple sclerosis patients to unravel novel pathogenic steps and identify potential biomarkers. Neurosci Lett 2012;508(1):4-8.
- 88. Fenoglio C, Cantoni C, De Riz M, Ridolfi E, Cortini F, Serpente M, et al. Expression and genetic analysis of miRNAs involved in CD4+ cell activation in patients with multiple sclerosis. Neurosci Lett 2011;504(1):9-12.
- Otaegui D, Baranzini SE, Armañanzas R, Calvo B, Muñoz-Culla M, Khankhanian P, et al. Differential micro RNA expression in PBMC from multiple sclerosis patients. PloS One 2009;4(7):e6309.
- 90. Keller A, Leidinger P, Lange J, Borries A, Schroers H, Scheffler M, et al. Multiple sclerosis: microRNA expression profiles accurately differentiate patients with relapsing-remitting disease from healthy controls. PloS One 2009;4(10):e7440.
- Junker A, Krumbholz M, Eisele S, Mohan H, Augstein F, Bittner R, et al. MicroRNA profiling of multiple sclerosis lesions identifies modulators of the regulatory protein CD47. Brain 2009;132(12):3342-52.
- 92. Zhang J, Cheng Y, Cui W, Li M, Li B, Guo L. MicroRNA-155 modulates Th1 and Th17 cell differentiation and is associated with multiple sclerosis and experimental autoimmune encephalomyelitis. J Neuroimmunol 2014;266(1-2):56-63.
- 93. Park NJ, Zhou H, Elashoff D, Henson BS, Kastratovic DA, Abemayor E, et al. Salivary microRNA: discovery, characterization, and clinical utility for oral cancer detection. Clin Cancer Res 2009;15(17):5473-7.
- 94. Mitchell PS, Parkin RK, Kroh EM, Fritz BR, Wyman SK, Pogosova-Agadjanyan EL, et al. Circulating microRNAs as stable blood-based markers for cancer detection. Proceed Natl Acad Sci

2008;105(30):10513-8.

- Redell JB, Moore AN, Ward III NH, Hergenroeder GW, Dash PK. Human traumatic brain injury alters plasma microRNA levels. J Neurotrauma 2010;27(12):2147-56.
- Regev K, Paul A, Healy B, von Glenn F, Diaz-Cruz C, Gholipour T, et al. Comprehensive evaluation of serum microRNAs as biomarkers in multiple sclerosis. Neurol Neuroimmunol Neuroinflam 2016;3(5):e267.
- Srivastava R, Aslam M, Kalluri SR, Schirmer L, Buck D, Tackenberg B, et al. Potassium channel KIR4. 1 as an immune target in multiple sclerosis. New Eng J Med 2012;367(2):115-23.
- Corvol JC, Pelletier D, Henry RG, Caillier SJ, Wang J, Pappas D, et al. Abrogation of T cell quiescence characterizes patients at high risk for multiple sclerosis after the initial neurological event. Proceed Natl Acad Sci 2008;105(33):11839-44.
- Chellaiah MA, Kizer N, Biswas R, Alvarez U, Strauss-Schoenberger J, Rifas L, et al. Osteopontin deficiency produces osteoclast dysfunction due to reduced CD44 surface expression. Mol Biol Cell 2003;14(1):173-89.
- 100. Scatena M, Liaw L, Giachelli CM. Osteopontin: a multifunctional molecule regulating chronic inflammation and vascular disease. Arterioscleros Thrombos Vasc Biol 2007;27(11):2302-9.
- 101. Crawford HC, Matrisian LM, Liaw L. Distinct roles of osteopontin in host defense activity and tumor survival during squamous cell carcinoma progression in vivo. Cancer Res 1998;58(22):5206-15.
- 102. Giachelli CM, Lombardi D, Johnson RJ, Murry CE, Almeida M. Evidence for a role of osteopontin in macrophage infiltration in response to pathological stimuli in vivo. Am J Pathol 1998;152(2):353.
- 103. Ashkar S, Weber GF, Panoutsakopoulou V, Sanchirico ME, Jansson M, Zawaideh S, et al. Eta-1 (osteopontin): an early component of type-1 (cell-mediated) immunity. Science 2000;287(5454):860-4.
- 104. Comi C, Cappellano G, Chiocchetti A, Orilieri E, Buttini S, Ghezzi L, et al. The impact of osteopontin gene variations on multiple sclerosis development and progression. Clin Develop Immunol 2012;2012.
- 105. Kivisäkk P, Healy BC, Francois K, Gandhi R, Gholipour T, Egorova S, et al. Evaluation of circulating osteopontin levels in an unselected cohort of patients with multiple sclerosis: relevance for biomarker development. Multiple Scleros J 2014;20(4):438-44.
- 106. Shimizu Y, Ota K, Ikeguchi R, Kubo S, Kabasawa C, Uchiyama S. Plasma osteopontin levels are associated with disease activity in the patients with multiple sclerosis and neuromyelitis optica. J Neuroimmunol 2013;263(1-2):148-51.
- 107. Wen SR, Liu GJ, Feng RN, Gong FC, Zhong H, Duan SR, et al. Increased levels of IL-23 and osteopontin in serum and cerebrospinal fluid of multiple sclerosis patients. J Neuroimmunol 2012;244(1-2):94-6.
- 108. Ji HI, Lee SH, Song R, Yang HI, Lee YA, Hong SJ, et al. Serum level of osteopontin as an inflammatory marker does not indicate disease activity or responsiveness to therapeutic treatments in patients with rheumatoid arthritis. Clin Rheumatol 2014;33(3):397-402.
- 109. Iwadate H, Kobayashi H, Kanno T, Asano T, Saito R, Sato S, et al. Plasma osteopontin is correlated with bone resorption markers in rheumatoid arthritis patients. Int J Rheumat Dis 2014;17(1):50-6.
- 110. Chen YJ, Shen JL, Wu CY, Chang YT, Chen CM, Lee FY. Elevated plasma osteopontin level is associated with occurrence of psoriasis and is an unfavorable cardiovascular risk factor in patients with psoriasis. J Am Acad Dermatol 2009;60(2):225-30.
- 111. Xu L, Ma X, Wang Y, Li X, Qi Y, Cui B, et al. The expression and pathophysiological role of osteopontin in Graves' disease. J Clin Endocrinol Metabol 2011;96(11):E1866-E70.
- 112. Chabas D, Baranzini SE, Mitchell D, Bernard CC, Rittling SR, Denhardt DT, et al. The influence of the proinflammatory cytokine, osteopontin, on autoimmune demyelinating disease. Science 2001;294(5547):1731-5.
- 113. Sinclair C, Mirakhur M, Kirk J, Farrell M, McQuaid S. Up-regulation of osteopontin and αB-crystallin in the normal-appearing white matter of multiple sclerosis: an immunohistochemical study utilizing tissue microarrays. Neuropathol Appl Neurobiol 2005;31(3):292-303.
- 114. Scher JU, Sczesnak A, Longman RS, Segata N, Ubeda C, Bielski C, et al. Expansion of intestinal Prevotella copri correlates with enhanced susceptibility to arthritis. Elife 2013;2:e01202.
- 115. Alkanani AK, Hara N, Gottlieb PA, Ir D, Robertson CE, Wagner BD, et al. Alterations in intestinal microbiota correlate with susceptibility to type 1 diabetes. Diabetes 2015;64(10):3510-20.

- 116. Kostic AD, Xavier RJ, Gevers D. The microbiome in inflammatory bowel disease: current status and the future ahead. Gastroenterology 2014;146(6):1489-99.
- 117. Jangi S, Gandhi R, Cox LM, Li N, Von Glehn F, Yan R, et al. Alterations of the human gut microbiome in multiple sclerosis. Nature Commun 2016;7:12015.
- 118. Ingram G, Hakobyan S, Hirst CL, Harris CL, Pickersgill TP, Cossburn MD, et al. Complement regulator factor H as a serum biomarker of multiple sclerosis disease state. Brain 2010;133(6):1602-11.
- 119. Minagar A, Alexander JS. Blood-brain barrier disruption in multiple sclerosis. Multiple Sclerosis Journal. 2003;9(6):540-9.
- 120. Minagar A, Jy W, Jimenez JJ, Sheremata W, Mauro L, Mao W, et al. Elevated plasma endothelial microparticles in multiple sclerosis. Neurology 2001;56(10):1319-24.
- 121. Hartung HP, Reiners K, Archelos JJ, Michels M, Seeldrayers P, Heidenreich F, et al. Circulating adhesion molecules and tumor necrosis factor receptor in multiple sclerosis: correlation with magnetic resonance imaging. Ann Neurol 1995;38(2):186-93.
- 122. Kuenz B, Lutterotti A, Khalil M, Ehling R, Gneiss C, Deisenhammer F, et al. Plasma levels of soluble adhesion molecules sPECAM-1, sPselectin and sE-selectin are associated with relapsing-remitting disease course of multiple sclerosis. J Neuroimmunol 2005;167(1-2):143-9.
- 123. Acar G, İdiman F, Kirkali G, Özakbaş S, Oktay G, Çakmakçi H, et al. Intrathecal sICAM-1 production in multiple sclerosis correlation with triple dose Gd-DTPA MRI enhancement and IgG index. J Neurol 2005;252(2):146-50.
- 124. Calabresi PA, Tranquill LR, Dambrosia JM, Stone LA, Maloni H, Bash CN, et al. Increases in soluble VCAM-1 correlate with a decrease in MRI lesions in multiple sclerosis treated with interferon β -1b. Ann

Neurol 1997;41(5):669-74.

- 125. Rieckmann P, Kruse N, Nagelkerken L, Beckmann K, Miller D, Polman C, et al. Soluble vascular cell adhesion molecule (VCAM) is associated with treatment effects of interferon beta-1b in patients with secondary progressive multiple sclerosis. J Neurol 2005;252(5):526-33.
- 126. Soilu-Hänninen M, Laaksonen M, Hänninen A, Erälinna J-P, Panelius M. Downregulation of VLA-4 on T cells as a marker of long term treatment response to interferon beta-1a in MS. J Neuroimmunol 2005;167(1-2):175-82.
- 127. Muraro P, Leist T, Bielekova B, McFarland H. VLA-4/CD49d downregulated on primed T lymphocytes during interferon-β therapy in multiple sclerosis. Journal of neuroimmunology. 2000;111(1-2):186-94.
- 128. Nikbin B, Bonab MM, Khosravi F, Talebian F. Role of B cells in pathogenesis of multiple sclerosis. Int Rev Neurobiol 2007;79:13-42.
- 129. Steinman L. Multiple sclerosis: a two-stage disease. Nature Immunol 2001;2(9):762.
- 130. Karabudak R, Kurne A, Guc D, Sengelen M, Canpinar H, Kansu E. Effect of interferon β-1a on serum matrix metalloproteinase—9 (MMP-9) and tissue inhibitor of matrix metalloproteinase (TIMP-1) in relapsing remitting multiple sclerosis patients. J Neurol 2004;251(3):279-83.
- 131. Garcia-Montojo M, Dominguez-Mozo M, De Las Heras V, Bartolome M, Garcia-Martinez A, Arroyo R, et al. Neutralizing antibodies, MxA expression and MMP-9/TIMP-1 ratio as markers of bioavailability of interferon-beta treatment in multiple sclerosis patients: a two-year follow-up study. Eur J Neurol 2010;17(3):470-8.
- 132. Liuzzi G, Trojano M, Fanelli M, Avolio C, Fasano A, Livrea P, et al. Intrathecal synthesis of matrix metalloproteinase-9 in patients with multiple sclerosis: implication for pathogenesis. Multiple Scleros J 2002;8(3):222-8.